

Lynne Turner-Stokes

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3977170/publications.pdf>

Version: 2024-02-01

187
papers

7,572
citations

50276

46
h-index

71685

76
g-index

198
all docs

198
docs citations

198
times ranked

6049
citing authors

#	ARTICLE	IF	CITATIONS
1	Goal attainment scaling (GAS) in rehabilitation: a practical guide. <i>Clinical Rehabilitation</i> , 2009, 23, 362-370.	2.2	583
2	THE NORTHWICK PARK NECK PAIN QUESTIONNAIRE, DEVISED TO MEASURE NECK PAIN AND DISABILITY. <i>Rheumatology</i> , 1994, 33, 469-474.	1.9	250
3	Shoulder pain after stroke: a review of the evidence base to inform the development of an integrated care pathway. <i>Clinical Rehabilitation</i> , 2002, 16, 276-298.	2.2	238
4	The UK FIM+FAM: development and evaluation. <i>Clinical Rehabilitation</i> , 1999, 13, 277-287.	2.2	200
5	Multidisciplinary rehabilitation for adults with multiple sclerosis. <i>The Cochrane Library</i> , 2007, , CD006036.	2.8	188
6	Multi-disciplinary rehabilitation for acquired brain injury in adults of working age. <i>The Cochrane Library</i> , 2015, 2015, CD004170.	2.8	187
7	Botulinum toxin assessment, intervention and aftercare for upper limb hypertonicity in adults: international consensus statement. <i>European Journal of Neurology</i> , 2010, 17, 74-93.	3.3	174
8	Multi-disciplinary rehabilitation for acquired brain injury in adults of working age. , 2005, , CD004170.		145
9	Botulinum toxin A for treatment of upper limb spasticity following stroke: A multi-centre randomized placebo-controlled study of the effects on quality of life and other person-centred outcomes. <i>Journal of Rehabilitation Medicine</i> , 2009, 41, 536-544.	1.1	138
10	Depression after stroke: a review of the evidence base to inform the development of an integrated care pathway. Part 1: Diagnosis, frequency and impact. <i>Clinical Rehabilitation</i> , 2002, 16, 231-247.	2.2	137
11	Evidence for the effectiveness of multi-disciplinary rehabilitation following acquired brain injury: a synthesis of two systematic approaches. <i>Journal of Rehabilitation Medicine</i> , 2008, 40, 691-701.	1.1	133
12	Does reducing spasticity translate into functional benefit? An exploratory meta-analysis. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2004, 75, 1547-1551.	1.9	132
13	The pain beliefs questionnaire: an investigation of beliefs in the causes and consequences of pain. <i>Pain</i> , 1992, 51, 267-272.	4.2	119
14	Goal attainment scaling: Does it provide added value as a person-centred measure for evaluation of outcome in neurorehabilitation following acquired brain injury?. <i>Journal of Rehabilitation Medicine</i> , 2009, 41, 528-535.	1.1	118
15	Goal attainment scaling in the evaluation of treatment of upper limb spasticity with botulinum toxin: A secondary analysis from a double-blind placebo-controlled randomized clinical trial. <i>Journal of Rehabilitation Medicine</i> , 2010, 42, 81-89.	1.1	109
16	Goal attainment for spasticity management using botulinum toxin. <i>Physiotherapy Research International</i> , 2006, 11, 24-34.	1.5	104
17	Evaluation of functional outcome measures for the hemiparetic upper limb: A systematic review. <i>Journal of Rehabilitation Medicine</i> , 2008, 40, 787-795.	1.1	103
18	Multidisciplinary rehabilitation programmes following joint replacement at the hip and knee in chronic arthropathy. <i>The Cochrane Library</i> , 2010, 2010, CD004957.	2.8	103

#	ARTICLE	IF	CITATIONS
19	The use of standardized outcome measures in rehabilitation centres in the UK. <i>Clinical Rehabilitation</i> , 1997, 11, 306-313.	2.2	100
20	From diagnosis to death: exploring the interface between neurology, rehabilitation and palliative care in managing people with long-term neurological conditions. <i>Clinical Medicine</i> , 2007, 7, 129-136.	1.9	100
21	The Northwick Park Dependency Score (NPDS): a measure of nursing dependency in rehabilitation. <i>Clinical Rehabilitation</i> , 1998, 12, 304-318.	2.2	99
22	Complex regional pain syndrome in adults: concise guidance. <i>Clinical Medicine</i> , 2011, 11, 596-600.	1.9	99
23	The Depression Intensity Scale Circles (DISCs): a first evaluation of a simple assessment tool for depression in the context of brain injury. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2005, 76, 1273-1278.	1.9	81
24	Multiple sclerosis: disability profile and quality of life in an Australian community cohort. <i>International Journal of Rehabilitation Research</i> , 2006, 29, 87-96.	1.3	77
25	Efficiency of specialist rehabilitation in reducing dependency and costs of continuing care for adults with complex acquired brain injuries. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2006, 77, 634-639.	1.9	73
26	Use of Goal Attainment Scaling in Inpatient Rehabilitation for Persons With Multiple Sclerosis. <i>Archives of Physical Medicine and Rehabilitation</i> , 2008, 89, 652-659.	0.9	73
27	The profile of patients and current practice of treatment of upper limb muscle spasticity with botulinum toxin type A: an international survey. <i>International Journal of Rehabilitation Research</i> , 2010, 33, 199-204.	1.3	70
28	Depression after stroke: a review of the evidence base to inform the development of an integrated care pathway. Part 2: Treatment alternatives. <i>Clinical Rehabilitation</i> , 2002, 16, 248-260.	2.2	67
29	Long-term neurological conditions: management at the interface between neurology, rehabilitation and palliative care. <i>Clinical Medicine</i> , 2008, 8, 186-191.	1.9	64
30	Results from the Upper Limb International Spasticity Study-II (ULIS-II): a large, international, prospective cohort study investigating practice and goal attainment following treatment with botulinum toxin A in real-life clinical management. <i>BMJ Open</i> , 2013, 3, e002771.	1.9	64
31	Factors contributing to work-ability for injured workers: literature review and comparison with available measures. <i>Disability and Rehabilitation</i> , 2010, 32, 1173-1183.	1.8	63
32	Multidisciplinary rehabilitation following botulinum toxin and other focal intramuscular treatment for post-stroke spasticity. <i>The Cochrane Library</i> , 2013, , CD009689.	2.8	62
33	The Northwick Park Care Needs Assessment (NPCNA): a directly costable outcome measure in rehabilitation. <i>Clinical Rehabilitation</i> , 1999, 13, 253-267.	2.2	61
34	Outpatient cognitive behavioral pain management programs: a randomized comparison of a group-based multidisciplinary versus an individual therapy model ¹¹ No commercial party having a direct financial interest in the results of the research supporting this article has or will confer a benefit upon the author(s) or upon any organization with which the author(s) is/are associated.. <i>Archives of Physical Medicine and Rehabilitation</i> , 2003, 84, 781-788.	0.9	61
35	Factor analysis and Rasch analysis of the Zarit Burden Interview for acquired brain injury carer research. <i>Journal of Rehabilitation Medicine</i> , 2010, 42, 302-309.	1.1	61
36	Rasch analysis of the Beck Depression Inventory-II in a neurological rehabilitation sample. <i>Disability and Rehabilitation</i> , 2010, 32, 8-17.	1.8	61

#	ARTICLE	IF	CITATIONS
37	A comprehensive psychometric evaluation of the UK FIM and FAM. Disability and Rehabilitation, 2013, 35, 1885-1895.	1.8	60
38	Patient engagement and satisfaction with goal planning: Impact on outcome from rehabilitation. International Journal of Therapy and Rehabilitation, 2015, 22, 210-216.	0.3	60
39	Effectiveness of brain injury rehabilitation. Clinical Rehabilitation, 1999, 13, 7-24.	2.2	59
40	Goal attainment scaling: a direct comparison of alternative rating methods. Clinical Rehabilitation, 2010, 24, 66-73.	2.2	59
41	Prophylaxis and treatment of infective endocarditis in adults: a concise guide. Clinical Medicine, 2004, 4, 545-550.	1.9	58
42	Cost-efficiency of specialist inpatient rehabilitation for working-aged adults with complex neurological disabilities: a multicentre cohort analysis of a national clinical data set. BMJ Open, 2016, 6, e010238.	1.9	58
43	The use of standardized outcome measures in rehabilitation centres in the UK. Clinical Rehabilitation, 2006, 20, 609-615.	2.2	55
44	Three-dimensional motion analysis of upper limb movement in the bowing arm of string-playing musicians. Clinical Biomechanics, 1999, 14, 426-433.	1.2	52
45	Managing depression in brain injury rehabilitation: the use of an integrated care pathway and preliminary report of response to sertraline. Clinical Rehabilitation, 2002, 16, 261-268.	2.2	52
46	Effectiveness of vocational rehabilitation intervention on the return to work and employment of persons with multiple sclerosis. The Cochrane Library, 2009, , CD007256.	2.8	51
47	Cost-efficiency of longer-stay rehabilitation programmes: Can they provide value for money?. Brain Injury, 2007, 21, 1015-1021.	1.2	50
48	Acquired brain injury and dementia: A comparison of carer experiences. Brain Injury, 2009, 23, 433-444.	1.2	50
49	Self-referential selective memory in pain patients. British Journal of Clinical Psychology, 1993, 32, 365-374.	3.5	48
50	Initial psychometric evaluation of the Arm Activity Measure (ArMA): a measure of activity in the hemiparetic arm. Clinical Rehabilitation, 2013, 27, 728-740.	2.2	47
51	Rehabilitation interventions for foot drop in neuromuscular disease. , 2009, , CD003908.		44
52	The Rehabilitation Complexity Scale: a simple, practical tool to identify "complex specialised" services in neurological rehabilitation. Clinical Medicine, 2007, 7, 593-599.	1.9	43
53	Can a Barthel score be derived from the FIM?. Clinical Rehabilitation, 1999, 13, 56-63.	2.2	41
54	The Rehabilitation Complexity Scale "extended version: detection of patients with highly complex needs. Disability and Rehabilitation, 2012, 34, 715-720.	1.8	39

#	ARTICLE	IF	CITATIONS
55	Botulinum toxin in the management of spasticity in adults. <i>Clinical Medicine</i> , 2002, 2, 128-130.	1.9	38
56	Management of shoulder and proximal upper limb spasticity using botulinum toxin and concurrent therapy interventions: A preliminary analysis of goals and outcomes. <i>Disability and Rehabilitation</i> , 2009, 31, 220-226.	1.8	38
57	Conceptualisation and development of the arm activity measure (ArmA) for assessment of activity in the hemiparetic arm. <i>Disability and Rehabilitation</i> , 2013, 35, 1513-1518.	1.8	38
58	A comprehensive person-centered approach to adult spastic paresis: a consensus-based framework. <i>European Journal of Physical and Rehabilitation Medicine</i> , 2018, 54, 605-617.	2.2	38
59	Can a Barthel score be derived from the FIM?. <i>Clinical Rehabilitation</i> , 1999, 13, 56-63.	2.2	37
60	Upper limb international spasticity study: rationale and protocol for a large, international, multicentre prospective cohort study investigating management and goal attainment following treatment with botulinum toxin A in real-life clinical practice. <i>BMJ Open</i> , 2013, 3, e002230.	1.9	37
61	Reflex sympathetic dystrophy--a complex regional pain syndrome. <i>Disability and Rehabilitation</i> , 2002, 24, 939-947.	1.8	35
62	Screening for ability to complete a questionnaire: a preliminary evaluation of the AbilityQ and ShoulderQ for assessing shoulder pain in stroke patients. <i>Clinical Rehabilitation</i> , 2003, 17, 150-157.	2.2	35
63	The evidence for the cost-effectiveness of rehabilitation following acquired brain injury. <i>Clinical Medicine</i> , 2004, 4, 10-12.	1.9	35
64	Development of a pictorial scale of pain intensity for patients with communication impairments: initial validation in a general population. <i>Clinical Medicine</i> , 2006, 6, 580-585.	1.9	34
65	Common goal areas in the treatment of upper limb spasticity: a multicentre analysis. <i>Clinical Rehabilitation</i> , 2016, 30, 617-622.	2.2	34
66	An examination of the factor structure of the Beck Depression Inventory-II in a neurorehabilitation inpatient sample. <i>Journal of the International Neuropsychological Society</i> , 2009, 15, 142-147.	1.8	33
67	Engaging the hearts and minds of clinicians in outcome measurement – the UK rehabilitation outcomes collaborative approach. <i>Disability and Rehabilitation</i> , 2012, 34, 1871-1879.	1.8	33
68	International casemix and funding models: lessons for rehabilitation. <i>Clinical Rehabilitation</i> , 2012, 26, 195-208.	2.2	32
69	Assessment of shoulder pain in hemiplegia: Sensitivity of the ShoulderQ. <i>Disability and Rehabilitation</i> , 2006, 28, 389-395.	1.8	31
70	Generating the evidence base for the National Service Framework for Long Term Conditions: a new research typology. <i>Clinical Medicine</i> , 2006, 6, 91-97.	1.9	31
71	Work sampling: a quantitative analysis of nursing activity in a neurorehabilitation setting. <i>Journal of Advanced Nursing</i> , 2009, 65, 2097-2107.	3.3	31
72	Clinical standards for inpatient specialist rehabilitation services in the UK. <i>Clinical Rehabilitation</i> , 2000, 14, 468-480.	2.2	29

#	ARTICLE	IF	CITATIONS
73	Multidisciplinary care for Guillain-Barré syndrome. The Cochrane Library, 2010, , CD008505.	2.8	29
74	The Neurological Impairment Scale: reliability and validity as a predictor of functional outcome in neurorehabilitation. Disability and Rehabilitation, 2014, 36, 23-31.	1.8	29
75	Systematic review of patient-reported outcome measures for functional performance in the lower limb. Journal of Rehabilitation Medicine, 2015, 47, 9-17.	1.1	29
76	Poststroke depression: getting the full picture. Lancet, The, 2003, 361, 1757-1758.	13.7	28
77	The Rehabilitation Complexity Scale version 2: a clinimetric evaluation in patients with severe complex neurodisability. Journal of Neurology, Neurosurgery and Psychiatry, 2010, 81, 146-153.	1.9	28
78	The post-ICU presentation screen (PICUPS) and rehabilitation prescription (RP) for intensive care survivors part I: Development and preliminary clinimetric evaluation. Journal of the Intensive Care Society, 2022, 23, 253-263.	2.2	28
79	Symptomatic Therapy and Rehabilitation in Primary Progressive Multiple Sclerosis. Neurology Research International, 2011, 2011, 1-22.	1.3	27
80	Impact of integrated upper limb spasticity management including botulinum toxin A on patient-centred goal attainment: rationale and protocol for an international prospective, longitudinal cohort study (ULIS-III). BMJ Open, 2016, 6, e011157.	1.9	27
81	Can young severely disabled stroke patients regain the ability to walk independently more than three months post stroke?. Clinical Rehabilitation, 2000, 14, 538-547.	2.2	26
82	Development of an integrated care pathway for the management of hemiplegic shoulder pain. Disability and Rehabilitation, 2002, 24, 390-398.	1.8	26
83	Politics, policy and payment – facilitators or barriers to person-centred rehabilitation?. Disability and Rehabilitation, 2007, 29, 1575-1582.	1.8	26
84	Goal Attainment Scaling and its relationship with standardized outcome measures: A commentary. Journal of Rehabilitation Medicine, 2011, 43, 70-72.	1.1	26
85	Estimated Life-Time Savings in the Cost of Ongoing Care Following Specialist Rehabilitation for Severe Traumatic Brain Injury in the United Kingdom. Journal of Head Trauma Rehabilitation, 2019, 34, 205-214.	1.7	26
86	The UK FIM+FAM Extended Activities of Daily Living module: evaluation of scoring accuracy and reliability. Disability and Rehabilitation, 2009, 31, 825-830.	1.8	25
87	Investigating muscle selection for botulinum toxin-A injections in adults with post-stroke upper limb spasticity. Journal of Rehabilitation Medicine, 2011, 43, 1032-1037.	1.1	25
88	Healthcare tariffs for specialist inpatient neurorehabilitation services: rationale and development of a UK casemix and costing methodology. Clinical Rehabilitation, 2012, 26, 264-279.	2.2	25
89	A matter of life and death: controversy at the interface between clinical and legal decision-making in prolonged disorders of consciousness. Journal of Medical Ethics, 2017, 43, 469-475.	1.8	25
90	Systematic Review of Upper Limb Function Measurement Methods in Botulinum Toxin Intervention for Focal Spasticity. Physiotherapy Research International, 2013, 18, 178-189.	1.5	24

#	ARTICLE	IF	CITATIONS
91	Psychometric evaluation of the Northwick Park Dependency Scale. <i>Journal of Rehabilitation Medicine</i> , 2010, 42, 936-943.	1.1	23
92	The Needs and Provision Complexity Scale: a multicentre prospective cohort analysis of met and unmet needs and their cost implications for patients with complex neurological disability. <i>BMJ Open</i> , 2013, 3, e002353.	1.9	23
93	Cost-efficiency of specialist hyperacute in-patient rehabilitation services for medically unstable patients with complex rehabilitation needs: a prospective cohort analysis. <i>BMJ Open</i> , 2016, 6, e012112.	1.9	23
94	The Associated Reaction Rating Scale: a clinical tool to measure associated reactions in the hemiplegic upper limb. <i>Clinical Rehabilitation</i> , 2002, 16, 726-735.	2.2	22
95	Standardized outcome assessment in brain injury rehabilitation for younger adults. <i>Disability and Rehabilitation</i> , 2002, 24, 383-389.	1.8	22
96	Rehabilitation following acquired brain injury: concise guidance. <i>Clinical Medicine</i> , 2004, 4, 61-65.	1.9	21
97	Multidisciplinary rehabilitation for adults with multiple sclerosis. <i>Postgraduate Medical Journal</i> , 2008, 84, 385-385.	1.8	21
98	Multiple sclerosis rehabilitation outcomes: analysis of a national casemix data set from Australia. <i>Multiple Sclerosis Journal</i> , 2009, 15, 869-875.	3.0	21
99	Clinical practice improvement approach in multiple sclerosis rehabilitation: a pilot study. <i>International Journal of Rehabilitation Research</i> , 2010, 33, 238-247.	1.3	21
100	The UK Functional Assessment Measure (UK FIM+FAM): Psychometric Evaluation in Patients Undergoing Specialist Rehabilitation following a Stroke from the National UK Clinical Dataset. <i>PLoS ONE</i> , 2016, 11, e0147288.	2.5	21
101	Functional Outcomes and Efficiency of Rehabilitation in a National Cohort of Patients with Guillain - Barré Syndrome and Other Inflammatory Polyneuropathies. <i>PLoS ONE</i> , 2014, 9, e110532.	2.5	20
102	Prolonged disorders of consciousness: new national clinical guidelines from the Royal College of Physicians, London. <i>Clinical Medicine</i> , 2014, 14, 4-5.	1.9	20
103	Rasch analysis of the UK Functional Assessment Measure in patients with complex disability after stroke. <i>Journal of Rehabilitation Medicine</i> , 2018, 50, 420-428.	1.1	20
104	The Post-ICU presentation screen (PICUPS) and rehabilitation prescription (RP) for intensive care survivors part II: Clinical engagement and future directions for the national Post-Intensive care Rehabilitation Collaborative. <i>Journal of the Intensive Care Society</i> , 2022, 23, 264-272.	2.2	20
105	Clinical standards for specialist community rehabilitation services in the UK. <i>Clinical Rehabilitation</i> , 2001, 15, 611-623.	2.2	19
106	Serial injection of botulinum toxin for muscle imbalance due to regional spasticity in the upper limb. <i>Disability and Rehabilitation</i> , 2007, 29, 1806-1812.	1.8	19
107	Chronic spinal cord injury: management of patients in acute hospital settings. <i>Clinical Medicine</i> , 2008, 8, 70-74.	1.9	19
108	Multidisciplinary rehabilitation for follow-up of women treated for breast cancer. <i>The Cochrane Library</i> , 2021, 2021, CD009553.	2.8	19

#	ARTICLE	IF	CITATIONS
109	Serial measurement of Wessex Head Injury Matrix in the diagnosis of patients in vegetative and minimally conscious states: a cohort analysis. <i>BMJ Open</i> , 2015, 5, e006051-e006051.	1.9	19
110	Engaging patients with moderate to severe acquired brain injury in goal setting. <i>International Journal of Therapy and Rehabilitation</i> , 2016, 23, 20-31.	0.3	19
111	Costs of caring for adults with long-term neurological conditions. <i>Journal of Rehabilitation Medicine</i> , 2013, 45, 653-661.	1.1	18
112	Autoantibody and idiotype profile of lung involvement in autoimmune rheumatic disease.. <i>Annals of the Rheumatic Diseases</i> , 1990, 49, 160-162.	0.9	17
113	The Northwick Park Care Needs Assessment (NPCNA): a measure of community care needs: sensitivity to change during rehabilitation. <i>Clinical Rehabilitation</i> , 1999, 13, 482-491.	2.2	17
114	Can brain-injured patients participate in an aerobic exercise programme during early inpatient rehabilitation?. <i>Clinical Rehabilitation</i> , 2001, 15, 535-544.	2.2	17
115	Rehabilitation interventions for foot drop in neuromuscular disease. , 2007, , CD003908.		17
116	Development and initial validation of the Northwick Park Therapy Dependency Assessment. <i>Clinical Rehabilitation</i> , 2009, 23, 922-937.	2.2	17
117	Factors Influencing Goal Attainment in Patients with Post-Stroke Upper Limb Spasticity Following Treatment with Botulinum Toxin A in Real-Life Clinical Practice: Sub-Analyses from the Upper Limb International Spasticity (ULIS-II) Study. <i>Toxins</i> , 2015, 7, 1192-1205.	3.4	17
118	Longitudinal goal attainment with integrated upper limb spasticity management including repeat injections of botulinum toxin A: Findings from the prospective, observational Upper Limb International Spasticity (ULIS-III) cohort study. <i>Journal of Rehabilitation Medicine</i> , 2021, 53, jrm00157.	1.1	17
119	Comparison of Rehabilitation Outcomes for Long Term Neurological Conditions: A Cohort Analysis of the Australian Rehabilitation Outcomes Centre Dataset for Adults of Working Age. <i>PLoS ONE</i> , 2015, 10, e0132275.	2.5	17
120	A completed audit cycle and integrated care pathway for the management of depression following brain injury in a rehabilitation setting. <i>Clinical Rehabilitation</i> , 2002, 16, 534-540.	2.2	16
121	Use of antidepressant medication following acquired brain injury: concise guidance. <i>Clinical Medicine</i> , 2005, 5, 268-274.	1.9	16
122	A longitudinal, multicentre, cohort study of community rehabilitation service delivery in long-term neurological conditions. <i>BMJ Open</i> , 2014, 4, e004231.	1.9	16
123	Staged residential post-acute rehabilitation for adults following acquired brain injury: A comparison of functional gains rated on the UK Functional Assessment Measure (UK FIM+FAM) and the Mayo-Portland Adaptability Inventory (MPAI-4). <i>Brain Injury</i> , 2017, 31, 1405-1413.	1.2	16
124	Life Expectancy after Stroke Based On Age, Sex, and Rankin Grade of Disability: A Synthesis. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2019, 28, 104450.	1.6	16
125	Outcome Measures for Inpatient Neurorehabilitation Settings. <i>Neuropsychological Rehabilitation</i> , 1999, 9, 329-343.	1.6	15
126	Arm Activity measure (ArmA) application for recording functional gain following focal spasticity treatment. <i>International Journal of Therapy and Rehabilitation</i> , 2014, 21, 10-17.	0.3	15

#	ARTICLE	IF	CITATIONS
127	Goal setting, using goal attainment scaling, as a method to identify patient selected items for measuring arm function. <i>Physiotherapy</i> , 2015, 101, 88-94.	0.4	15
128	Introduction. <i>Clinical Rehabilitation</i> , 1999, 13, 3-6.	2.2	14
129	Outcome measurement in brain injury rehabilitation - towards a common language. <i>Clinical Rehabilitation</i> , 1999, 13, 273-275.	2.2	14
130	An evaluation of aftercare following discharge from a specialist in-patient rehabilitation service. <i>Disability and Rehabilitation</i> , 2003, 25, 1281-1288.	1.8	14
131	Deriving a Barthel Index from the Northwick Park Dependency Scale and the Functional Independence Measure: are they equivalent?. <i>Clinical Rehabilitation</i> , 2010, 24, 1121-1126.	2.2	14
132	A cost analysis of specialist inpatient neurorehabilitation services in the UK. <i>Clinical Rehabilitation</i> , 2012, 26, 256-263.	2.2	13
133	Service Use and Costs for People with Long-Term Neurological Conditions in the First Year following Discharge from In-Patient Neuro-Rehabilitation: A Longitudinal Cohort Study. <i>PLoS ONE</i> , 2014, 9, e113056.	2.5	13
134	Updated NICE guidance on chronic fatigue syndrome. <i>BMJ, The</i> , 2020, 371, m4774.	6.0	13
135	Measurement of haematological indices of chronic rheumatic disease with two newer generation automated systems, the H1 and H6000 (Technicon).. <i>Annals of the Rheumatic Diseases</i> , 1991, 50, 583-587.	0.9	12
136	Assessing the effectiveness of upper-limb spasticity management using a structured approach to goal-setting and outcome measurement: First cycle results from the ULIS-III Study. <i>Journal of Rehabilitation Medicine</i> , 2021, 53, jrm00133.	1.1	11
137	The national service framework for long term conditions: a novel approach for a "new style" NSF. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2005, 76, 901-902.	1.9	10
138	The Needs and Provision Complexity Scale: a first psychometric analysis using multicentre data. <i>Clinical Rehabilitation</i> , 2014, 28, 687-695.	2.2	10
139	Time between acquired brain injury and admission to community-based rehabilitation: differences in cognitive and functional gains. <i>Brain Injury</i> , 2020, 34, 713-722.	1.2	10
140	Use of an integrated care pathway: a third round audit of the management of shoulder pain in neurological conditions. <i>Journal of Rehabilitation Medicine</i> , 2003, 35, 265-270.	1.1	9
141	Complex Specialized Rehabilitation Following Severe Brain Injury. <i>Journal of Head Trauma Rehabilitation</i> , 2007, 22, 239-247.	1.7	9
142	Using the UKROC dataset to make the case for resources to improve cost-efficiency in neurological rehabilitation. <i>Disability and Rehabilitation</i> , 2012, 34, 1900-1906.	1.8	9
143	The Work-ability Support Scale: Evaluation of Scoring Accuracy and Rater Reliability. <i>Journal of Occupational Rehabilitation</i> , 2014, 24, 511-524.	2.2	9
144	Categorisation of goals set using Goal Attainment Scaling for treatment of leg spasticity: a multicentre analysis. <i>Disability and Rehabilitation</i> , 2019, 41, 1925-1930.	1.8	9

#	ARTICLE	IF	CITATIONS
145	Northwick Park Care Needs Assessment: adaptation for inpatient neurological rehabilitation settings. <i>Journal of Advanced Nursing</i> , 2007, 59, 612-622.	3.3	8
146	Screening for pain in patients with cognitive and communication difficulties: evaluation of the SPIN-screen. <i>Clinical Medicine</i> , 2008, 8, 393-398.	1.9	8
147	fMRI for vegetative and minimally conscious states. <i>BMJ, The</i> , 2012, 345, e8045-e8045.	6.0	8
148	Development of a new tool to evaluate work support needs and guide vocational rehabilitation: the Work-ability Support Scale (WSS). <i>Disability and Rehabilitation</i> , 2015, 37, 247-258.	1.8	8
149	Management of Neurogenic Heterotopic Ossification. <i>Physiotherapy</i> , 2003, 89, 471-477.	0.4	7
150	The National Service Framework for Long Term Conditions: the challenges ahead. <i>Clinical Medicine</i> , 2005, 5, 203-206.	1.9	7
151	The patient categorisation tool: psychometric evaluation of a tool to measure complexity of needs for rehabilitation in a large multicentre dataset from the United Kingdom. <i>Disability and Rehabilitation</i> , 2019, 41, 1101-1109.	1.8	7
152	Validation of the Memory and Behavior Problems Checklist-1990R for use in acquired brain injury. <i>Brain Injury</i> , 2007, 21, 817-824.	1.2	6
153	Accuracy of prediction of walking for young stroke patients by use of the FIM. <i>Physiotherapy Research International</i> , 2001, 6, 1-14.	1.5	5
154	The development of clinical governance in the UK: its implications for rehabilitation medicine. <i>Clinical Medicine</i> , 2003, 3, 135-141.	1.9	5
155	Diagnosis and treatment of depression following acquired brain injury: a comparison of practice in the UK and the Netherlands. <i>Clinical Rehabilitation</i> , 2007, 21, 805-811.	2.2	5
156	A systematic review protocol to evaluate the psychometric properties of measures of function within adult neuro-rehabilitation. <i>Systematic Reviews</i> , 2015, 4, 86.	5.3	5
157	Conceptualization and Development of the Leg Activity Measure (LegA) for Patient and Carer Reported Assessment of Activity in the Paretic Leg. <i>Physiotherapy Research International</i> , 2017, 22, e1660.	1.5	5
158	Goal-setting and attainment in prolonged disorders of consciousness – development of a structured approach. <i>Brain Injury</i> , 2020, 34, 78-88.	1.2	5
159	Secondary safety of car adaptations for disabled motorists. <i>Disability and Rehabilitation</i> , 1996, 18, 317-327.	1.8	4
160	Can the Northwick Park Dependency Scale be translated to a Barthel Index?. <i>Clinical Rehabilitation</i> , 2010, 24, 1112-1120.	2.2	4
161	Unmet need for specialised rehabilitation following neurosurgery: can we maximise the potential cost-benefits?. <i>British Journal of Neurosurgery</i> , 2017, 31, 249-253.	0.8	4
162	Abobotulinumtoxina injections in shoulder muscles to improve adult upper limb spasticity: Results from a phase 4 real-world study and a phase 3 open-label trial. <i>Journal of Rehabilitation Medicine</i> , 2020, 52, jrm00068.	1.1	4

#	ARTICLE	IF	CITATIONS
163	Psychometric properties of measures of upper limb activity performance in adults with and without spasticity undergoing neurorehabilitationâ€“A systematic review. PLoS ONE, 2021, 16, e0246288.	2.5	4
164	A pictorial scale of pain intensity (SPIN) for patients with communication impairments. International Journal of Therapy and Rehabilitation, 2006, 13, 457-463.	0.3	3
165	The Northwick Park Therapy Dependency Assessment scale: a psychometric analysis from a large multicentre neurorehabilitation dataset. Disability and Rehabilitation, 2015, 37, 1976-1983.	1.8	3
166	Dimensionality and scaling properties of the Patient Categorisation Tool in patients with complex rehabilitation needs following acquired brain injury. Journal of Rehabilitation Medicine, 2018, 50, 435-443.	1.1	3
167	The Leg Activity measure, a new measure of passive and active function and impact on quality of life; informing goal setting and outcome evaluation in leg spasticity. Disability and Rehabilitation, 2019, 43, 1-9.	1.8	3
168	Psychometric evaluation of the leg activity measure (LegA) for outcome measurement in people with brain injury and spasticity. Disability and Rehabilitation, 2021, 43, 976-987.	1.8	3
169	Functional outcomes and cost-efficiency of specialist in-patient rehabilitation following spinal cord injury: a multi-centre national cohort analysis from the UK Rehabilitation Outcomes Collaborative (UKROC). Disability and Rehabilitation, 2022, 44, 5603-5611.	1.8	3
170	Prolonged disorders of consciousness: identification using the UK FIM + FAM and cohort analysis of outcomes from a UK national clinical database. Disability and Rehabilitation, 2023, 45, 620-629.	1.8	3
171	Development of a measure of therapy provision for spasticity management in the paretic lower limb â€“ the Leg Therapy recording Schedule (LegTS). Disability and Rehabilitation, 2018, 40, 479-485.	1.8	2
172	Rasch analysis of the UK Functional Assessment Measure in a sample of patients with traumatic brain injury from the UK national clinical database. Journal of Rehabilitation Medicine, 2019, 51, 566-574.	1.1	2
173	Clinical governance in rehabilitation medicine. The state of the art in 2002. A British Society of Rehabilitation Medicine report. Clinical Rehabilitation, 2002, 16 Suppl 1, 1-60.	2.2	2
174	MRI and arthroscopic surgery: a combined breakthrough in management of shoulder pain.. Annals of the Rheumatic Diseases, 1996, 55, 405-406.	0.9	1
175	Can the Northwick Park Care Needs Assessment be used to estimate nursing staff requirements in an inpatient rehabilitation setting?. Clinical Rehabilitation, 2007, 21, 535-544.	2.2	1
176	Rehabilitation interventions for foot drop in neuromuscular disease. The Cochrane Library, 2015, , CD003908.	2.8	1
177	Spasticity management with botulinum toxin: A comparison of UK physiotherapy and rehabilitation medicine injectors. International Journal of Therapy and Rehabilitation, 2018, 25, 215-222.	0.3	1
178	Staff experiences of working with patients with prolonged disorders of consciousness: a focus group analysis. International Journal of Therapy and Rehabilitation, 2018, 25, 602-612.	0.3	1
179	Cost-efficiency of specialist inpatient rehabilitation for adults with multiple sclerosis: A multicentre prospective cohort analysis of the UK Rehabilitation Outcomes Collaborative national clinical dataset for rehabilitation centres in England. Multiple Sclerosis Journal - Experimental, Translational and Clinical, 2020, 6, 205521732091278.	1.0	1
180	Stratified management of hemiplegic shoulder pain using an integrated care pathway: an 18-year clinical cohort analysis. Disability and Rehabilitation, 2021, , 1-10.	1.8	1

#	ARTICLE	IF	CITATIONS
181	The spasticity-related quality of life 6-dimensions instrument in upper-limb spasticity: Development and responsiveness. <i>Journal of Rehabilitation Medicine</i> , 2021, , .	1.1	1
182	The spasticity-related quality of life 6-dimensions instrument in upper-limb spasticity: A first psychometric evaluation. <i>Journal of Rehabilitation Medicine</i> , 2021, , .	1.1	1
183	Rheumatoid Arthritis, Impairment and Mobility. <i>Rheumatology</i> , 1994, 33, 408-408.	1.9	0
184	Poster 373â€œ Integrated Upper Limb Spasticity Management Including Botulinum Toxin A (BoNTâ€œA) on Patientâ€œCentered Goal Attainment: Methodology for ULISâ€œIII and Initial Goalâ€œSetting Data. <i>PM and R</i> , 2016, 8, S331.	1.6	0
185	Poster 313 Relationship Between AbobotulinumtoxinA Injections into Shoulder Muscles and Patient Centred Primary Goal Selection and Achievement: Sub-Analyses from the Upper Limb International Spasticity (ULIS)-II Study. <i>PM and R</i> , 2016, 8, S262.	1.6	0
186	Poster 57: Time to Retreatment with Botulinum Toxin A in Upper Limb Spasticity Management: Initial Data from the Upper Limb International Spasticity (ULIS)â€œIII Study. <i>PM and R</i> , 2017, 9, S155.	1.6	0
187	Health Policy: United States and International Perspectives. , 2012, , .		0