

# Lynne Turner-Stokes

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3977170/publications.pdf>

Version: 2024-02-01

187  
papers

7,572  
citations

50276

46  
h-index

71685

76  
g-index

198  
all docs

198  
docs citations

198  
times ranked

6049  
citing authors

#	ARTICLE	IF	CITATIONS
1	Prolonged disorders of consciousness: identification using the UK FIM + FAM and cohort analysis of outcomes from a UK national clinical database. <i>Disability and Rehabilitation</i> , 2023, 45, 620-629.	1.8	3
2	The Post-ICU presentation screen (PICUPS) and rehabilitation prescription (RP) for intensive care survivors part II: Clinical engagement and future directions for the national Post-Intensive care Rehabilitation Collaborative. <i>Journal of the Intensive Care Society</i> , 2022, 23, 264-272.	2.2	20
3	The post-ICU presentation screen (PICUPS) and rehabilitation prescription (RP) for intensive care survivors part I: Development and preliminary clinimetric evaluation. <i>Journal of the Intensive Care Society</i> , 2022, 23, 253-263.	2.2	28
4	Functional outcomes and cost-efficiency of specialist in-patient rehabilitation following spinal cord injury: a multi-centre national cohort analysis from the UK Rehabilitation Outcomes Collaborative (UKROC). <i>Disability and Rehabilitation</i> , 2022, 44, 5603-5611.	1.8	3
5	Multidisciplinary rehabilitation for follow-up of women treated for breast cancer. <i>The Cochrane Library</i> , 2021, 2021, CD009553.	2.8	19
6	Assessing the effectiveness of upper-limb spasticity management using a structured approach to goal-setting and outcome measurement: First cycle results from the ULIS-III Study. <i>Journal of Rehabilitation Medicine</i> , 2021, 53, jrm00133.	1.1	11
7	Psychometric evaluation of the leg activity measure (LegA) for outcome measurement in people with brain injury and spasticity. <i>Disability and Rehabilitation</i> , 2021, 43, 976-987.	1.8	3
8	Longitudinal goal attainment with integrated upper limb spasticity management including repeat injections of botulinum toxin A: Findings from the prospective, observational Upper Limb International Spasticity (ULIS-III) cohort study. <i>Journal of Rehabilitation Medicine</i> , 2021, 53, jrm00157.	1.1	17
9	Psychometric properties of measures of upper limb activity performance in adults with and without spasticity undergoing neurorehabilitation – A systematic review. <i>PLoS ONE</i> , 2021, 16, e0246288.	2.5	4
10	Stratified management of hemiplegic shoulder pain using an integrated care pathway: an 18-year clinical cohort analysis. <i>Disability and Rehabilitation</i> , 2021, , 1-10.	1.8	1
11	The spasticity-related quality of life 6-dimensions instrument in upper-limb spasticity: Development and responsiveness. <i>Journal of Rehabilitation Medicine</i> , 2021, , .	1.1	1
12	The spasticity-related quality of life 6-dimensions instrument in upper-limb spasticity: A first psychometric evaluation. <i>Journal of Rehabilitation Medicine</i> , 2021, , .	1.1	1
13	Goal-setting and attainment in prolonged disorders of consciousness – development of a structured approach. <i>Brain Injury</i> , 2020, 34, 78-88.	1.2	5
14	Cost-efficiency of specialist inpatient rehabilitation for adults with multiple sclerosis: A multicentre prospective cohort analysis of the UK Rehabilitation Outcomes Collaborative national clinical dataset for rehabilitation centres in England. <i>Multiple Sclerosis Journal - Experimental, Translational and Clinical</i> , 2020, 6, 205521732091278.	1.0	1
15	AbobotulinumtoxinA injections in shoulder muscles to improve adult upper limb spasticity: Results from a phase 4 real-world study and a phase 3 open-label trial. <i>Journal of Rehabilitation Medicine</i> , 2020, 52, jrm00068.	1.1	4
16	Time between acquired brain injury and admission to community-based rehabilitation: differences in cognitive and functional gains. <i>Brain Injury</i> , 2020, 34, 713-722.	1.2	10
17	Updated NICE guidance on chronic fatigue syndrome. <i>BMJ</i> , The, 2020, 371, m4774.	6.0	13
18	Rasch analysis of the UK Functional Assessment Measure in a sample of patients with traumatic brain injury from the UK national clinical database. <i>Journal of Rehabilitation Medicine</i> , 2019, 51, 566-574.	1.1	2

#	ARTICLE	IF	CITATIONS
19	Life Expectancy after Stroke Based On Age, Sex, and Rankin Grade of Disability: A Synthesis. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2019, 28, 104450.	1.6	16
20	The Leg Activity measure, a new measure of passive and active function and impact on quality of life; informing goal setting and outcome evaluation in leg spasticity. <i>Disability and Rehabilitation</i> , 2019, 43, 1-9.	1.8	3
21	Estimated Life-Time Savings in the Cost of Ongoing Care Following Specialist Rehabilitation for Severe Traumatic Brain Injury in the United Kingdom. <i>Journal of Head Trauma Rehabilitation</i> , 2019, 34, 205-214.	1.7	26
22	Categorisation of goals set using Goal Attainment Scaling for treatment of leg spasticity: a multicentre analysis. <i>Disability and Rehabilitation</i> , 2019, 41, 1925-1930.	1.8	9
23	The patient categorisation tool: psychometric evaluation of a tool to measure complexity of needs for rehabilitation in a large multicentre dataset from the United Kingdom. <i>Disability and Rehabilitation</i> , 2019, 41, 1101-1109.	1.8	7
24	Spasticity management with botulinum toxin: A comparison of UK physiotherapy and rehabilitation medicine injectors. <i>International Journal of Therapy and Rehabilitation</i> , 2018, 25, 215-222.	0.3	1
25	Development of a measure of therapy provision for spasticity management in the paretic lower limb – the Leg Therapy recording Schedule (LegTS). <i>Disability and Rehabilitation</i> , 2018, 40, 479-485.	1.8	2
26	Staff experiences of working with patients with prolonged disorders of consciousness: a focus group analysis. <i>International Journal of Therapy and Rehabilitation</i> , 2018, 25, 602-612.	0.3	1
27	Rasch analysis of the UK Functional Assessment Measure in patients with complex disability after stroke. <i>Journal of Rehabilitation Medicine</i> , 2018, 50, 420-428.	1.1	20
28	A comprehensive person-centered approach to adult spastic paresis: a consensus-based framework. <i>European Journal of Physical and Rehabilitation Medicine</i> , 2018, 54, 605-617.	2.2	38
29	Dimensionality and scaling properties of the Patient Categorisation Tool in patients with complex rehabilitation needs following acquired brain injury. <i>Journal of Rehabilitation Medicine</i> , 2018, 50, 435-443.	1.1	3
30	Conceptualization and Development of the Leg Activity Measure (LegA) for Patient and Carer Reported Assessment of Activity in the Paretic Leg. <i>Physiotherapy Research International</i> , 2017, 22, e1660.	1.5	5
31	A matter of life and death: controversy at the interface between clinical and legal decision-making in prolonged disorders of consciousness. <i>Journal of Medical Ethics</i> , 2017, 43, 469-475.	1.8	25
32	Poster 57: Time to Retreatment with Botulinum Toxin A in Upper Limb Spasticity Management: Initial Data from the Upper Limb International Spasticity (ULIS) Study. <i>PM and R</i> , 2017, 9, S155.	1.6	0
33	Staged residential post-acute rehabilitation for adults following acquired brain injury: A comparison of functional gains rated on the UK Functional Assessment Measure (UK FIM+FAM) and the Mayo-Portland Adaptability Inventory (MPAI-4). <i>Brain Injury</i> , 2017, 31, 1405-1413.	1.2	16
34	Unmet need for specialised rehabilitation following neurosurgery: can we maximise the potential cost-benefits?. <i>British Journal of Neurosurgery</i> , 2017, 31, 249-253.	0.8	4
35	Engaging patients with moderate to severe acquired brain injury in goal setting. <i>International Journal of Therapy and Rehabilitation</i> , 2016, 23, 20-31.	0.3	19
36	Poster 373 – Integrated Upper Limb Spasticity Management Including Botulinum Toxin A (BoNTA) on Patient-Centered Goal Attainment: Methodology for ULIS and Initial Goal-Setting Data. <i>PM and R</i> , 2016, 8, S331.	1.6	0

#	ARTICLE	IF	CITATIONS
37	Poster 313 Relationship Between AbobotulinumtoxinA Injections into Shoulder Muscles and Patient Centred Primary Goal Selection and Achievement: Sub-Analyses from the Upper Limb International Spasticity (ULIS)-II Study. PM and R, 2016, 8, S262.	1.6	0
38	Cost-efficiency of specialist inpatient rehabilitation for working-aged adults with complex neurological disabilities: a multicentre cohort analysis of a national clinical data set. BMJ Open, 2016, 6, e010238.	1.9	58
39	Cost-efficiency of specialist hyperacute in-patient rehabilitation services for medically unstable patients with complex rehabilitation needs: a prospective cohort analysis. BMJ Open, 2016, 6, e012112.	1.9	23
40	Common goal areas in the treatment of upper limb spasticity: a multicentre analysis. Clinical Rehabilitation, 2016, 30, 617-622.	2.2	34
41	Impact of integrated upper limb spasticity management including botulinum toxin A on patient-centred goal attainment: rationale and protocol for an international prospective, longitudinal cohort study (ULIS-III). BMJ Open, 2016, 6, e011157.	1.9	27
42	The UK Functional Assessment Measure (UK FIM+FAM): Psychometric Evaluation in Patients Undergoing Specialist Rehabilitation following a Stroke from the National UK Clinical Dataset. PLoS ONE, 2016, 11, e0147288.	2.5	21
43	Patient engagement and satisfaction with goal planning: Impact on outcome from rehabilitation. International Journal of Therapy and Rehabilitation, 2015, 22, 210-216.	0.3	60
44	Multi-disciplinary rehabilitation for acquired brain injury in adults of working age. The Cochrane Library, 2015, 2015, CD004170.	2.8	187
45	Systematic review of patient-reported outcome measures for functional performance in the lower limb. Journal of Rehabilitation Medicine, 2015, 47, 9-17.	1.1	29
46	Serial measurement of Wessex Head Injury Matrix in the diagnosis of patients in vegetative and minimally conscious states: a cohort analysis. BMJ Open, 2015, 5, e006051-e006051.	1.9	19
47	The Northwick Park Therapy Dependency Assessment scale: a psychometric analysis from a large multicentre neurorehabilitation dataset. Disability and Rehabilitation, 2015, 37, 1976-1983.	1.8	3
48	Factors Influencing Goal Attainment in Patients with Post-Stroke Upper Limb Spasticity Following Treatment with Botulinum Toxin A in Real-Life Clinical Practice: Sub-Analyses from the Upper Limb International Spasticity (ULIS)-II Study. Toxins, 2015, 7, 1192-1205.	3.4	17
49	A systematic review protocol to evaluate the psychometric properties of measures of function within adult neuro-rehabilitation. Systematic Reviews, 2015, 4, 86.	5.3	5
50	Development of a new tool to evaluate work support needs and guide vocational rehabilitation: the Work-ability Support Scale (WSS). Disability and Rehabilitation, 2015, 37, 247-258.	1.8	8
51	Rehabilitation interventions for foot drop in neuromuscular disease. The Cochrane Library, 2015, , CD003908.	2.8	1
52	Goal setting, using goal attainment scaling, as a method to identify patient selected items for measuring arm function. Physiotherapy, 2015, 101, 88-94.	0.4	15
53	Comparison of Rehabilitation Outcomes for Long Term Neurological Conditions: A Cohort Analysis of the Australian Rehabilitation Outcomes Centre Dataset for Adults of Working Age. PLoS ONE, 2015, 10, e0132275.	2.5	17
54	Functional Outcomes and Efficiency of Rehabilitation in a National Cohort of Patients with Guillain - Barré Syndrome and Other Inflammatory Polyneuropathies. PLoS ONE, 2014, 9, e110532.	2.5	20

#	ARTICLE	IF	CITATIONS
55	Service Use and Costs for People with Long-Term Neurological Conditions in the First Year following Discharge from In-Patient Neuro-Rehabilitation: A Longitudinal Cohort Study. PLoS ONE, 2014, 9, e113056.	2.5	13
56	Arm Activity measure (ArmA) application for recording functional gain following focal spasticity treatment. International Journal of Therapy and Rehabilitation, 2014, 21, 10-17.	0.3	15
57	The Needs and Provision Complexity Scale: a first psychometric analysis using multicentre data. Clinical Rehabilitation, 2014, 28, 687-695.	2.2	10
58	The Neurological Impairment Scale: reliability and validity as a predictor of functional outcome in neurorehabilitation. Disability and Rehabilitation, 2014, 36, 23-31.	1.8	29
59	A longitudinal, multicentre, cohort study of community rehabilitation service delivery in long-term neurological conditions. BMJ Open, 2014, 4, e004231.	1.9	16
60	The Work-ability Support Scale: Evaluation of Scoring Accuracy and Rater Reliability. Journal of Occupational Rehabilitation, 2014, 24, 511-524.	2.2	9
61	Prolonged disorders of consciousness: new national clinical guidelines from the Royal College of Physicians, London. Clinical Medicine, 2014, 14, 4-5.	1.9	20
62	Multidisciplinary rehabilitation following botulinum toxin and other focal intramuscular treatment for post-stroke spasticity. The Cochrane Library, 2013, , CD009689.	2.8	62
63	Conceptualisation and development of the arm activity measure (ArmA) for assessment of activity in the hemiparetic arm. Disability and Rehabilitation, 2013, 35, 1513-1518.	1.8	38
64	A comprehensive psychometric evaluation of the UK FIM+â€‰FAM. Disability and Rehabilitation, 2013, 35, 1885-1895.	1.8	60
65	Upper limb international spasticity study: rationale and protocol for a large, international, multicentre prospective cohort study investigating management and goal attainment following treatment with botulinum toxin A in real-life clinical practice. BMJ Open, 2013, 3, e002230.	1.9	37
66	Results from the Upper Limb International Spasticity Study-II (ULIS-II): a large, international, prospective cohort study investigating practice and goal attainment following treatment with botulinum toxin A in real-life clinical management. BMJ Open, 2013, 3, e002771.	1.9	64
67	The Needs and Provision Complexity Scale: a multicentre prospective cohort analysis of met and unmet needs and their cost implications for patients with complex neurological disability. BMJ Open, 2013, 3, e002353.	1.9	23
68	Initial psychometric evaluation of the Arm Activity Measure (ArmA): a measure of activity in the hemiparetic arm. Clinical Rehabilitation, 2013, 27, 728-740.	2.2	47
69	Systematic Review of Upperâ€‰limb Function Measurement Methods in Botulinum Toxin Intervention for Focal Spasticity. Physiotherapy Research International, 2013, 18, 178-189.	1.5	24
70	Costs of caring for adults with long-term neurological conditions. Journal of Rehabilitation Medicine, 2013, 45, 653-661.	1.1	18
71	A cost analysis of specialist inpatient neurorehabilitation services in the UK. Clinical Rehabilitation, 2012, 26, 256-263.	2.2	13
72	Healthcare tariffs for specialist inpatient neurorehabilitation services: rationale and development of a UK casemix and costing methodology. Clinical Rehabilitation, 2012, 26, 264-279.	2.2	25

#	ARTICLE	IF	CITATIONS
73	International casemix and funding models: lessons for rehabilitation. <i>Clinical Rehabilitation</i> , 2012, 26, 195-208.	2.2	32
74	fMRI for vegetative and minimally conscious states. <i>BMJ, The</i> , 2012, 345, e8045-e8045.	6.0	8
75	Engaging the hearts and minds of clinicians in outcome measurement – the UK rehabilitation outcomes collaborative approach. <i>Disability and Rehabilitation</i> , 2012, 34, 1871-1879.	1.8	33
76	The Rehabilitation Complexity Scale – extended version: detection of patients with highly complex needs. <i>Disability and Rehabilitation</i> , 2012, 34, 715-720.	1.8	39
77	Using the UKROC dataset to make the case for resources to improve cost-efficiency in neurological rehabilitation. <i>Disability and Rehabilitation</i> , 2012, 34, 1900-1906.	1.8	9
78	Health Policy: United States and International Perspectives. , 2012, , .		0
79	Goal Attainment Scaling and its relationship with standardized outcome measures: A commentary. <i>Journal of Rehabilitation Medicine</i> , 2011, 43, 70-72.	1.1	26
80	Symptomatic Therapy and Rehabilitation in Primary Progressive Multiple Sclerosis. <i>Neurology Research International</i> , 2011, 2011, 1-22.	1.3	27
81	Investigating muscle selection for botulinum toxin-A injections in adults with post-stroke upper limb spasticity. <i>Journal of Rehabilitation Medicine</i> , 2011, 43, 1032-1037.	1.1	25
82	Complex regional pain syndrome in adults: concise guidance. <i>Clinical Medicine</i> , 2011, 11, 596-600.	1.9	99
83	Multidisciplinary rehabilitation programmes following joint replacement at the hip and knee in chronic arthropathy. <i>The Cochrane Library</i> , 2010, 2010, CD004957.	2.8	103
84	Factor analysis and Rasch analysis of the Zarit Burden Interview for acquired brain injury carer research. <i>Journal of Rehabilitation Medicine</i> , 2010, 42, 302-309.	1.1	61
85	Botulinum toxin assessment, intervention and after-care for upper limb hypertonicity in adults: international consensus statement. <i>European Journal of Neurology</i> , 2010, 17, 74-93.	3.3	174
86	Goal attainment scaling in the evaluation of treatment of upper limb spasticity with botulinum toxin: A secondary analysis from a double-blind placebo-controlled randomized clinical trial. <i>Journal of Rehabilitation Medicine</i> , 2010, 42, 81-89.	1.1	109
87	Psychometric evaluation of the Northwick Park Dependency Scale. <i>Journal of Rehabilitation Medicine</i> , 2010, 42, 936-943.	1.1	23
88	Goal attainment scaling: a direct comparison of alternative rating methods. <i>Clinical Rehabilitation</i> , 2010, 24, 66-73.	2.2	59
89	The Rehabilitation Complexity Scale version 2: a clinimetric evaluation in patients with severe complex neurodisability. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2010, 81, 146-153.	1.9	28
90	Clinical practice improvement approach in multiple sclerosis rehabilitation: a pilot study. <i>International Journal of Rehabilitation Research</i> , 2010, 33, 238-247.	1.3	21

#	ARTICLE	IF	CITATIONS
91	The profile of patients and current practice of treatment of upper limb muscle spasticity with botulinum toxin type A: an international survey. <i>International Journal of Rehabilitation Research</i> , 2010, 33, 199-204.	1.3	70
92	Rasch analysis of the Beck Depression Inventory-II in a neurological rehabilitation sample. <i>Disability and Rehabilitation</i> , 2010, 32, 8-17.	1.8	61
93	Can the Northwick Park Dependency Scale be translated to a Barthel Index?. <i>Clinical Rehabilitation</i> , 2010, 24, 1112-1120.	2.2	4
94	Deriving a Barthel Index from the Northwick Park Dependency Scale and the Functional Independence Measure: are they equivalent?. <i>Clinical Rehabilitation</i> , 2010, 24, 1121-1126.	2.2	14
95	Factors contributing to work-ability for injured workers: literature review and comparison with available measures. <i>Disability and Rehabilitation</i> , 2010, 32, 1173-1183.	1.8	63
96	Multidisciplinary care for Guillain-Barré syndrome. <i>The Cochrane Library</i> , 2010, , CD008505.	2.8	29
97	Goal attainment scaling: Does it provide added value as a person-centred measure for evaluation of outcome in neurorehabilitation following acquired brain injury?. <i>Journal of Rehabilitation Medicine</i> , 2009, 41, 528-535.	1.1	118
98	Acquired brain injury and dementia: A comparison of carer experiences. <i>Brain Injury</i> , 2009, 23, 433-444.	1.2	50
99	Botulinum toxin A for treatment of upper limb spasticity following stroke: A multi-centre randomized placebo-controlled study of the effects on quality of life and other person-centred outcomes. <i>Journal of Rehabilitation Medicine</i> , 2009, 41, 536-544.	1.1	138
100	Development and initial validation of the Northwick Park Therapy Dependency Assessment. <i>Clinical Rehabilitation</i> , 2009, 23, 922-937.	2.2	17
101	Multiple sclerosis rehabilitation outcomes: analysis of a national casemix data set from Australia. <i>Multiple Sclerosis Journal</i> , 2009, 15, 869-875.	3.0	21
102	Work sampling: a quantitative analysis of nursing activity in a neurorehabilitation setting. <i>Journal of Advanced Nursing</i> , 2009, 65, 2097-2107.	3.3	31
103	Rehabilitation interventions for foot drop in neuromuscular disease. , 2009, , CD003908.		44
104	The UK FIM+FAM Extended Activities of Daily Living module: evaluation of scoring accuracy and reliability. <i>Disability and Rehabilitation</i> , 2009, 31, 825-830.	1.8	25
105	Goal attainment scaling (GAS) in rehabilitation: a practical guide. <i>Clinical Rehabilitation</i> , 2009, 23, 362-370.	2.2	583
106	Management of shoulder and proximal upper limb spasticity using botulinum toxin and concurrent therapy interventions: A preliminary analysis of goals and outcomes. <i>Disability and Rehabilitation</i> , 2009, 31, 220-226.	1.8	38
107	An examination of the factor structure of the Beck Depression Inventory-II in a neurorehabilitation inpatient sample. <i>Journal of the International Neuropsychological Society</i> , 2009, 15, 142-147.	1.8	33
108	Effectiveness of vocational rehabilitation intervention on the return to work and employment of persons with multiple sclerosis. <i>The Cochrane Library</i> , 2009, , CD007256.	2.8	51

#	ARTICLE	IF	CITATIONS
109	Use of Goal Attainment Scaling in Inpatient Rehabilitation for Persons With Multiple Sclerosis. Archives of Physical Medicine and Rehabilitation, 2008, 89, 652-659.	0.9	73
110	Multidisciplinary rehabilitation for adults with multiple sclerosis. Postgraduate Medical Journal, 2008, 84, 385-385.	1.8	21
111	Evaluation of functional outcome measures for the hemiparetic upper limb: A systematic review. Journal of Rehabilitation Medicine, 2008, 40, 787-795.	1.1	103
112	Evidence for the effectiveness of multi-disciplinary rehabilitation following acquired brain injury: a synthesis of two systematic approaches. Journal of Rehabilitation Medicine, 2008, 40, 691-701.	1.1	133
113	Long-term neurological conditions: management at the interface between neurology, rehabilitation and palliative care. Clinical Medicine, 2008, 8, 186-191.	1.9	64
114	Chronic spinal cord injury: management of patients in acute hospital settings. Clinical Medicine, 2008, 8, 70-74.	1.9	19
115	Screening for pain in patients with cognitive and communication difficulties: evaluation of the SPIN-screen. Clinical Medicine, 2008, 8, 393-398.	1.9	8
116	From diagnosis to death: exploring the interface between neurology, rehabilitation and palliative care in managing people with long-term neurological conditions. Clinical Medicine, 2007, 7, 129-136.	1.9	100
117	The Rehabilitation Complexity Scale: a simple, practical tool to identify "complex specialised" services in neurological rehabilitation. Clinical Medicine, 2007, 7, 593-599.	1.9	43
118	Diagnosis and treatment of depression following acquired brain injury: a comparison of practice in the UK and the Netherlands. Clinical Rehabilitation, 2007, 21, 805-811.	2.2	5
119	Can the Northwick Park Care Needs Assessment be used to estimate nursing staff requirements in an inpatient rehabilitation setting?. Clinical Rehabilitation, 2007, 21, 535-544.	2.2	1
120	Rehabilitation interventions for foot drop in neuromuscular disease. , 2007, , CD003908.		17
121	Complex Specialized Rehabilitation Following Severe Brain Injury. Journal of Head Trauma Rehabilitation, 2007, 22, 239-247.	1.7	9
122	Validation of the Memory and Behavior Problems Checklist-1990R for use in acquired brain injury. Brain Injury, 2007, 21, 817-824.	1.2	6
123	Cost-efficiency of longer-stay rehabilitation programmes: Can they provide value for money?. Brain Injury, 2007, 21, 1015-1021.	1.2	50
124	Politics, policy and payment "facilitators or barriers to person-centred rehabilitation?. Disability and Rehabilitation, 2007, 29, 1575-1582.	1.8	26
125	Serial injection of botulinum toxin for muscle imbalance due to regional spasticity in the upper limb. Disability and Rehabilitation, 2007, 29, 1806-1812.	1.8	19
126	Multidisciplinary rehabilitation for adults with multiple sclerosis. The Cochrane Library, 2007, , CD006036.	2.8	188



#	ARTICLE	IF	CITATIONS
127	Northwick Park Care Needs Assessment: adaptation for inpatient neurological rehabilitation settings. <i>Journal of Advanced Nursing</i> , 2007, 59, 612-622.	3.3	8
128	Assessment of shoulder pain in hemiplegia: Sensitivity of the ShoulderQ. <i>Disability and Rehabilitation</i> , 2006, 28, 389-395.	1.8	31
129	A pictorial scale of pain intensity (SPIN) for patients with communication impairments. <i>International Journal of Therapy and Rehabilitation</i> , 2006, 13, 457-463.	0.3	3
130	Multiple sclerosis: disability profile and quality of life in an Australian community cohort. <i>International Journal of Rehabilitation Research</i> , 2006, 29, 87-96.	1.3	77
131	Goal attainment for spasticity management using botulinum toxin. <i>Physiotherapy Research International</i> , 2006, 11, 24-34.	1.5	104
132	Efficiency of specialist rehabilitation in reducing dependency and costs of continuing care for adults with complex acquired brain injuries. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2006, 77, 634-639.	1.9	73
133	The use of standardized outcome measures in rehabilitation centres in the UK. <i>Clinical Rehabilitation</i> , 2006, 20, 609-615.	2.2	55
134	Generating the evidence base for the National Service Framework for Long Term Conditions: a new research typology. <i>Clinical Medicine</i> , 2006, 6, 91-97.	1.9	31
135	Development of a pictorial scale of pain intensity for patients with communication impairments: initial validation in a general population. <i>Clinical Medicine</i> , 2006, 6, 580-585.	1.9	34
136	The National Service Framework for Long Term Conditions: the challenges ahead. <i>Clinical Medicine</i> , 2005, 5, 203-206.	1.9	7
137	The Depression Intensity Scale Circles (DISCs): a first evaluation of a simple assessment tool for depression in the context of brain injury. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2005, 76, 1273-1278.	1.9	81
138	The national service framework for long term conditions: a novel approach for a "new style" NSF. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2005, 76, 901-902.	1.9	10
139	Use of antidepressant medication following acquired brain injury: concise guidance. <i>Clinical Medicine</i> , 2005, 5, 268-274.	1.9	16
140	Multi-disciplinary rehabilitation for acquired brain injury in adults of working age. , 2005, , CD004170.		145
141	Rehabilitation following acquired brain injury: concise guidance. <i>Clinical Medicine</i> , 2004, 4, 61-65.	1.9	21
142	Does reducing spasticity translate into functional benefit? An exploratory meta-analysis. <i>Journal of Neurology, Neurosurgery and Psychiatry</i> , 2004, 75, 1547-1551.	1.9	132
143	Prophylaxis and treatment of infective endocarditis in adults: a concise guide. <i>Clinical Medicine</i> , 2004, 4, 545-550.	1.9	58
144	The evidence for the cost-effectiveness of rehabilitation following acquired brain injury. <i>Clinical Medicine</i> , 2004, 4, 10-12.	1.9	35

#	ARTICLE	IF	CITATIONS
145	Management of Neurogenic Heterotopic Ossification. <i>Physiotherapy</i> , 2003, 89, 471-477.	0.4	7
146	Screening for ability to complete a questionnaire: a preliminary evaluation of the AbilityQ and ShoulderQ for assessing shoulder pain in stroke patients. <i>Clinical Rehabilitation</i> , 2003, 17, 150-157.	2.2	35
147	Poststroke depression: getting the full picture. <i>Lancet, The</i> , 2003, 361, 1757-1758.	13.7	28
148	Outpatient cognitive behavioral pain management programs: a randomized comparison of a group-based multidisciplinary versus an individual therapy model <sup>11</sup> No commercial party having a direct financial interest in the results of the research supporting this article has or will confer a benefit upon the author(s) or upon any organization with which the author(s) is/are associated.. <i>Archives of Physical Medicine and Rehabilitation</i> , 2003, 84, 781-788.	0.9	61
149	An evaluation of aftercare following discharge from a specialist in-patient rehabilitation service. <i>Disability and Rehabilitation</i> , 2003, 25, 1281-1288.	1.8	14
150	The development of clinical governance in the UK: its implications for rehabilitation medicine. <i>Clinical Medicine</i> , 2003, 3, 135-141.	1.9	5
151	Use of an integrated care pathway: a third round audit of the management of shoulder pain in neurological conditions. <i>Journal of Rehabilitation Medicine</i> , 2003, 35, 265-270.	1.1	9
152	Depression after stroke: a review of the evidence base to inform the development of an integrated care pathway. Part 2: Treatment alternatives. <i>Clinical Rehabilitation</i> , 2002, 16, 248-260.	2.2	67
153	Depression after stroke: a review of the evidence base to inform the development of an integrated care pathway. Part 1: Diagnosis, frequency and impact. <i>Clinical Rehabilitation</i> , 2002, 16, 231-247.	2.2	137
154	Reflex sympathetic dystrophy--a complex regional pain syndrome. <i>Disability and Rehabilitation</i> , 2002, 24, 939-947.	1.8	35
155	A completed audit cycle and integrated care pathway for the management of depression following brain injury in a rehabilitation setting. <i>Clinical Rehabilitation</i> , 2002, 16, 534-540.	2.2	16
156	Managing depression in brain injury rehabilitation: the use of an integrated care pathway and preliminary report of response to sertraline. <i>Clinical Rehabilitation</i> , 2002, 16, 261-268.	2.2	52
157	The Associated Reaction Rating Scale: a clinical tool to measure associated reactions in the hemiplegic upper limb. <i>Clinical Rehabilitation</i> , 2002, 16, 726-735.	2.2	22
158	Botulinum toxin in the management of spasticity in adults. <i>Clinical Medicine</i> , 2002, 2, 128-130.	1.9	38
159	Development of an integrated care pathway for the management of hemiplegic shoulder pain. <i>Disability and Rehabilitation</i> , 2002, 24, 390-398.	1.8	26
160	Standardized outcome assessment in brain injury rehabilitation for younger adults. <i>Disability and Rehabilitation</i> , 2002, 24, 383-389.	1.8	22
161	Shoulder pain after stroke: a review of the evidence base to inform the development of an integrated care pathway. <i>Clinical Rehabilitation</i> , 2002, 16, 276-298.	2.2	238
162	Clinical governance in rehabilitation medicine. The state of the art in 2002. A British Society of Rehabilitation Medicine report. <i>Clinical Rehabilitation</i> , 2002, 16 Suppl 1, 1-60.	2.2	2

#	ARTICLE	IF	CITATIONS
163	Accuracy of prediction of walking for young stroke patients by use of the FIM. <i>Physiotherapy Research International</i> , 2001, 6, 1-14.	1.5	5
164	Can brain-injured patients participate in an aerobic exercise programme during early inpatient rehabilitation?. <i>Clinical Rehabilitation</i> , 2001, 15, 535-544.	2.2	17
165	Clinical standards for specialist community rehabilitation services in the UK. <i>Clinical Rehabilitation</i> , 2001, 15, 611-623.	2.2	19
166	Clinical standards for inpatient specialist rehabilitation services in the UK. <i>Clinical Rehabilitation</i> , 2000, 14, 468-480.	2.2	29
167	Can young severely disabled stroke patients regain the ability to walk independently more than three months post stroke?. <i>Clinical Rehabilitation</i> , 2000, 14, 538-547.	2.2	26
168	Introduction. <i>Clinical Rehabilitation</i> , 1999, 13, 3-6.	2.2	14
169	Outcome Measures for Inpatient Neurorehabilitation Settings. <i>Neuropsychological Rehabilitation</i> , 1999, 9, 329-343.	1.6	15
170	Effectiveness of brain injury rehabilitation. <i>Clinical Rehabilitation</i> , 1999, 13, 7-24.	2.2	59
171	The UK FIM+FAM: development and evaluation. <i>Clinical Rehabilitation</i> , 1999, 13, 277-287.	2.2	200
172	Can a Barthel score be derived from the FIM?. <i>Clinical Rehabilitation</i> , 1999, 13, 56-63.	2.2	37
173	Outcome measurement in brain injury rehabilitation - towards a common language. <i>Clinical Rehabilitation</i> , 1999, 13, 273-275.	2.2	14
174	The Northwick Park Care Needs Assessment (NPCNA): a measure of community care needs: sensitivity to change during rehabilitation. <i>Clinical Rehabilitation</i> , 1999, 13, 482-491.	2.2	17
175	The Northwick Park Care Needs Assessment (NPCNA): a directly costable outcome measure in rehabilitation. <i>Clinical Rehabilitation</i> , 1999, 13, 253-267.	2.2	61
176	Three-dimensional motion analysis of upper limb movement in the bowing arm of string-playing musicians. <i>Clinical Biomechanics</i> , 1999, 14, 426-433.	1.2	52
177	Can a Barthel score be derived from the FIM?. <i>Clinical Rehabilitation</i> , 1999, 13, 56-63.	2.2	41
178	The Northwick Park Dependency Score (NPDS): a measure of nursing dependency in rehabilitation. <i>Clinical Rehabilitation</i> , 1998, 12, 304-318.	2.2	99
179	The use of standardized outcome measures in rehabilitation centres in the UK. <i>Clinical Rehabilitation</i> , 1997, 11, 306-313.	2.2	100
180	MRI and arthroscopic surgery: a combined breakthrough in management of shoulder pain.. <i>Annals of the Rheumatic Diseases</i> , 1996, 55, 405-406.	0.9	1

#	ARTICLE	IF	CITATIONS
181	Secondary safety of car adaptations for disabled motorists. Disability and Rehabilitation, 1996, 18, 317-327.	1.8	4
182	Rheumatoid Arthritis, Impairment and Mobility. Rheumatology, 1994, 33, 408-408.	1.9	0
183	THE NORTHWICK PARK NECK PAIN QUESTIONNAIRE, DEVISED TO MEASURE NECK PAIN AND DISABILITY. Rheumatology, 1994, 33, 469-474.	1.9	250
184	Self-referential selective memory in pain patients. British Journal of Clinical Psychology, 1993, 32, 365-374.	3.5	48
185	The pain beliefs questionnaire: an investigation of beliefs in the causes and consequences of pain. Pain, 1992, 51, 267-272.	4.2	119
186	Measurement of haematological indices of chronic rheumatic disease with two newer generation automated systems, the H1 and H6000 (Technicon).. Annals of the Rheumatic Diseases, 1991, 50, 583-587.	0.9	12
187	Autoantibody and idiotype profile of lung involvement in autoimmune rheumatic disease.. Annals of the Rheumatic Diseases, 1990, 49, 160-162.	0.9	17