

Judith Lechner

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3983182/publications.pdf>

Version: 2024-02-01

10
papers

624
citations

1478505

6
h-index

1588992

8
g-index

10
all docs

10
docs citations

10
times ranked

932
citing authors

#	ARTICLE	IF	CITATIONS
1	The pathology associated with diabetic retinopathy. Vision Research, 2017, 139, 7-14.	1.4	319
2	Higher plasma levels of complement C3a, C4a and C5a increase the risk of subretinal fibrosis in neovascular age-related macular degeneration. Immunity and Ageing, 2016, 13, 4.	4.2	81
3	Topical Delivery of Anti-VEGF Drugs to the Ocular Posterior Segment Using Cell-Penetrating Peptides. , 2017, 58, 2578.		70
4	Peripheral blood mononuclear cells from neovascular age-related macular degeneration patients produce higher levels of chemokines CCL2 (MCP-1) and CXCL8 (IL-8). Journal of Neuroinflammation, 2017, 14, 42.	7.2	49
5	Whole-mitochondrial genome sequencing in primary open-angle glaucoma using massively parallel sequencing identifies novel and known pathogenic variants. Genetics in Medicine, 2015, 17, 279-284.	2.4	38
6	Alterations in Circulating Immune Cells in Neovascular Age-Related Macular Degeneration. Scientific Reports, 2015, 5, 16754.	3.3	34
7	VEGF-B Is an Autocrine Gliotrophic Factor for M μ ller Cells under Pathologic Conditions. , 2020, 61, 35.		13
8	miR-130a activates the VEGFR2/STAT3/HIF1 α axis to potentiate the vasoregenerative capacity of endothelial colony-forming cells in hypoxia. Molecular Therapy - Nucleic Acids, 2021, 23, 968-981.	5.1	9
9	Plasma level of lipocalin 2 is increased in neovascular age-related macular degeneration patients, particularly those with macular fibrosis. Immunity and Ageing, 2020, 17, 35.	4.2	8
10	Sustained intraocular vascular endothelial growth factor neutralisation does not affect retinal and choroidal vasculature in Ins2 ^{AKITA} diabetic mice. Diabetes and Vascular Disease Research, 2019, 16, 440-449.	2.0	3