## Judith Lechner

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3983182/publications.pdf

Version: 2024-02-01

1478505 1588992 10 624 6 8 citations h-index g-index papers 10 10 10 932 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	miR-130a activates the VEGFR2/STAT3/HIF1 $\hat{l}$ ± axis to potentiate the vasoregenerative capacity of endothelial colony-forming cells in hypoxia. Molecular Therapy - Nucleic Acids, 2021, 23, 968-981.	5.1	9
2	Plasma level of lipocalin 2 is increased in neovascular age-related macular degeneration patients, particularly those with macular fibrosis. Immunity and Ageing, 2020, 17, 35.	4.2	8
3	VEGF-B Is an Autocrine Gliotrophic Factor for Mýller Cells under Pathologic Conditions. , 2020, 61, 35.		13
4	Sustained intraocular vascular endothelial growth factor neutralisation does not affect retinal and choroidal vasculature in Ins2 <sup>Akita</sup> diabetic mice. Diabetes and Vascular Disease Research, 2019, 16, 440-449.	2.0	3
5	Peripheral blood mononuclear cells from neovascular age-related macular degeneration patients produce higher levels of chemokines CCL2 (MCP-1) and CXCL8 (IL-8). Journal of Neuroinflammation, 2017, 14, 42.	7.2	49
6	The pathology associated with diabetic retinopathy. Vision Research, 2017, 139, 7-14.	1.4	319
7	Topical Delivery of Anti-VEGF Drugs to the Ocular Posterior Segment Using Cell-Penetrating Peptides. , 2017, 58, 2578.		70
8	Higher plasma levels of complement C3a, C4a and C5a increase the risk of subretinal fibrosis in neovascular age-related macular degeneration. Immunity and Ageing, 2016, 13, 4.	4.2	81
9	Alterations in Circulating Immune Cells in Neovascular Age-Related Macular Degeneration. Scientific Reports, 2015, 5, 16754.	3.3	34
10	Whole-mitochondrial genome sequencing in primary open-angle glaucoma using massively parallel sequencing identifies novel and known pathogenic variants. Genetics in Medicine, 2015, 17, 279-284.	2.4	38