

# Marcella Mun-San Kwan

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3984129/publications.pdf>

Version: 2024-02-01

20  
papers

1,684  
citations

516215

16  
h-index

752256

20  
g-index

20  
all docs

20  
docs citations

20  
times ranked

2544  
citing authors

#	ARTICLE	IF	CITATIONS
1	Factors associated with appointment non-attendance at a medical imaging department in regional Australia: a retrospective cohort analysis. <i>Journal of Medical Radiation Sciences</i> , 2018, 65, 192-199.	0.8	16
2	Effect of Exercise and Cognitive Training on Falls and Fall-Related Factors in Older Adults With Mild Cognitive Impairment: A Systematic Review. <i>Archives of Physical Medicine and Rehabilitation</i> , 2017, 98, 2079-2096.	0.5	61
3	The rural pipeline to longer-term rural practice: General practitioners and specialists. <i>PLoS ONE</i> , 2017, 12, e0180394.	1.1	55
4	Fall risk in Chinese community-dwelling older adults: A physiological profile assessment study. <i>Geriatrics and Gerontology International</i> , 2016, 16, 259-265.	0.7	11
5	Comparison of faecal microbiota in Blastocystis-positive and Blastocystis-negative irritable bowel syndrome patients. <i>Microbiome</i> , 2016, 4, 47.	4.9	77
6	Blastocystis specific serum immunoglobulin in patients with irritable bowel syndrome (IBS) versus healthy controls. <i>Parasites and Vectors</i> , 2015, 8, 453.	1.0	15
7	Effectiveness of Exergaming Training in Reducing Risk and Incidence of Falls in Frail Older Adults With a History of Falls. <i>Archives of Physical Medicine and Rehabilitation</i> , 2015, 96, 2096-2102.	0.5	78
8	Development and validation of a Chinese version of the Falls Efficacy Scale International. <i>Archives of Gerontology and Geriatrics</i> , 2013, 56, 169-174.	1.4	55
9	Increased Concern Is Protective for Falls in Chinese Older People: The Chopstix Fall Risk Study. <i>Journals of Gerontology - Series A Biological Sciences and Medical Sciences</i> , 2013, 68, 946-953.	1.7	38
10	Depressive symptoms in addition to visual impairment, reduced strength and poor balance predict falls in older Taiwanese people. <i>Age and Ageing</i> , 2012, 41, 606-612.	0.7	31
11	Sensorimotor function, balance abilities and pain influence Timed Up and Go performance in older community-living people. <i>Aging Clinical and Experimental Research</i> , 2011, 23, 196-201.	1.4	59
12	Minimal Chair Height Standing Ability Is Independently Associated With Falls in Taiwanese Older People. <i>Archives of Physical Medicine and Rehabilitation</i> , 2011, 92, 1080-1085.	0.5	11
13	Falls Incidence, Risk Factors, and Consequences in Chinese Older People: A Systematic Review. <i>Journal of the American Geriatrics Society</i> , 2011, 59, 536-543.	1.3	196
14	Effect on falls of providing single lens distance vision glasses to multifocal glasses wearers: VISIBLE randomised controlled trial. <i>BMJ: British Medical Journal</i> , 2010, 340, c2265-c2265.	2.4	104
15	Preventing falls in older multifocal glasses wearers by providing single-lens distance glasses: the protocol for the VISIBLE randomised controlled trial. <i>BMC Geriatrics</i> , 2009, 9, 10.	1.1	18
16	Group exercise can improve participants' mobility in an outpatient rehabilitation setting: a randomized controlled trial. <i>Clinical Rehabilitation</i> , 2008, 22, 493-502.	1.0	39
17	A novel weight-bearing strengthening program during rehabilitation of older people is feasible and improves standing up more than a non-weight-bearing strengthening program: a randomised trial. <i>Australian Journal of Physiotherapy</i> , 2007, 53, 147-153.	0.9	37
18	Foot pain in community-dwelling older people: an evaluation of the Manchester Foot Pain and Disability Index. <i>Rheumatology</i> , 2006, 45, 863-867.	0.9	126

#	ARTICLE	IF	CITATIONS
19	Reliability of the GAITRite® walkway system for the quantification of temporo-spatial parameters of gait in young and older people. <i>Gait and Posture</i> , 2004, 20, 20-25.	0.6	577
20	Reliability of Clinical Tests of Foot and Ankle Characteristics in Older People. <i>Journal of the American Podiatric Medical Association</i> , 2003, 93, 380-387.	0.2	80