

Atsuko Abiko

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/39889/publications.pdf>

Version: 2024-02-01

13
papers

246
citations

1307594

7
h-index

1372567

10
g-index

16
all docs

16
docs citations

16
times ranked

491
citing authors

#	ARTICLE	IF	CITATIONS
1	Long-term safety and efficacy of alogliptin, a DPP-4 inhibitor, in patients with type 2 diabetes: a 3-year prospective, controlled, observational study (J-BRAND Registry). <i>BMJ Open Diabetes Research and Care</i> , 2021, 9, e001787.	2.8	15
2	The work style and living condition survey of diabetologists and the expectations for the Japan Diabetes Society: results of questionnaires about the current state and the future prospect of their carrier in 2017. <i>Diabetology International</i> , 2020, 11, 299-308.	1.4	0
3	Increment of plasma glucose by exogenous glucagon is associated with present and future renal function in type 2 diabetes: a retrospective study from glucagon stimulation test. <i>BMC Endocrine Disorders</i> , 2019, 19, 99.	2.2	2
4	Safety and efficacy of once-a-weekly semaglutide vs additional oral antidiabetic drugs in Japanese people with inadequately controlled type 2 diabetes: A randomized trial. <i>Diabetes, Obesity and Metabolism</i> , 2018, 20, 1202-1212.	4.4	81
5	Impacts of Diabetes and an SGLT2 Inhibitor on the Glomerular Number and Volume in db/db Mice, as Estimated by Synchrotron Radiation Micro-CT at SPring-8. <i>EBioMedicine</i> , 2018, 36, 329-346.	6.1	25
6	Dipeptidyl peptidase-4 inhibitor treatment induces a greater increase in plasma levels of bioactive GIP than GLP-1 in non-diabetic subjects. <i>Molecular Metabolism</i> , 2017, 6, 226-231.	6.5	31
7	Prediabetes Exhibits Decreased Disposition Index Correlated with Deterioration of Glycemic Parameters in Nonobese Japanese Subjects: A Cross-Sectional Study from Medical Examination. <i>Metabolic Syndrome and Related Disorders</i> , 2017, 15, 296-303.	1.3	3
8	Expression of transcription factors in MEN1-associated pancreatic neuroendocrine tumors. <i>Endocrinology, Diabetes and Metabolism Case Reports</i> , 2017, 2017, .	0.5	1
9	Sodium-Glucose Cotransporter 2 Inhibitor and a Low Carbohydrate Diet Affect Gluconeogenesis and Glycogen Content Differently in the Kidney and the Liver of Non-Diabetic Mice. <i>PLoS ONE</i> , 2016, 11, e0157672.	2.5	20
10	High glucose induces platelet-derived growth factor-C via carbohydrate response element-binding protein in glomerular mesangial cells. <i>Physiological Reports</i> , 2016, 4, e12730.	1.7	7
11	Pancreatic glucose-dependent insulinotropic polypeptide (GIP) (1-30) expression is upregulated in diabetes and PEGylated GIP(1-30) can suppress the progression of low-dose-STZ-induced hyperglycaemia in mice. <i>Diabetologia</i> , 2016, 59, 533-541.	6.3	21
12	What is the truth about renin angiotensin blockers for diabetic patients?. <i>Diabetology International</i> , 2016, 7, 334-337.	1.4	0
13	Imaging of the Corneal Subbasal Whorl-like Nerve Plexus: More Accurate Depiction of the Extent of Corneal Nerve Damage in Patients With Diabetes. , 2015, 56, 5417.		40