Tiziana Genovese

List of Publications by Citations

Source: https://exaly.com/author-pdf/3992912/tiziana-genovese-publications-by-citations.pdf

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

162 papers

5,663 citations

44 h-index 61 g-index

168 ext. papers

6,278 ext. citations

5.4 avg, IF

4.96 L-index

#	Paper	IF	Citations
162	Rosiglitazone, a ligand of the peroxisome proliferator-activated receptor-gamma, reduces acute inflammation. <i>European Journal of Pharmacology</i> , 2004 , 483, 79-93	5.3	167
161	Reduction in the evolution of murine type II collagen-induced arthritis by treatment with rosiglitazone, a ligand of the peroxisome proliferator-activated receptor gamma. <i>Arthritis and Rheumatism</i> , 2003 , 48, 3544-56		128
160	Immunomodulatory effects of etanercept in an experimental model of spinal cord injury. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2006 , 316, 1006-16	4.7	127
159	Protective effects of anthocyanins from blackberry in a rat model of acute lung inflammation. <i>Free Radical Research</i> , 2003 , 37, 891-900	4	126
158	Uric acid protects against secondary damage after spinal cord injury. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2005 , 102, 3483-8	11.5	109
157	Attenuation in the evolution of experimental spinal cord trauma by treatment with melatonin. Journal of Pineal Research, 2005 , 38, 198-208	10.4	94
156	Lipopolysaccharide-induced tumor necrosis factor alpha production and not monocyte human leukocyte antigen-DR expression is correlated with survival in septic trauma patients. <i>Shock</i> , 2006 , 25, 129-34	3.4	92
155	Effects of palmitoylethanolamide on signaling pathways implicated in the development of spinal cord injury. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2008 , 326, 12-23	4.7	90
154	The P2Y-like receptor GPR17 as a sensor of damage and a new potential target in spinal cord injury. <i>Brain</i> , 2009 , 132, 2206-18	11.2	89
153	Glycyrrhizin attenuates the development of carrageenan-induced lung injury in mice. <i>Pharmacological Research</i> , 2008 , 58, 22-31	10.2	89
152	Absence of TLR4 reduces neurovascular unit and secondary inflammatory process after traumatic brain injury in mice. <i>PLoS ONE</i> , 2013 , 8, e57208	3.7	88
151	Role of endogenous and exogenous ligands for the peroxisome proliferators activated receptors alpha (PPAR-alpha) in the development of inflammatory bowel disease in mice. <i>Laboratory Investigation</i> , 2004 , 84, 1643-54	5.9	88
150	Role of glucocorticoid-induced TNF receptor family gene (GITR) in collagen-induced arthritis. <i>FASEB Journal</i> , 2005 , 19, 1253-65	0.9	88
149	Rosiglitazone and 15-deoxy-Delta12,14-prostaglandin J2, ligands of the peroxisome proliferator-activated receptor-gamma (PPAR-gamma), reduce ischaemia/reperfusion injury of the gut. <i>British Journal of Pharmacology</i> , 2003 , 140, 366-76	8.6	81
148	Effects of palmitoylethanolamide on release of mast cell peptidases and neurotrophic factors after spinal cord injury. <i>Brain, Behavior, and Immunity</i> , 2011 , 25, 1099-112	16.6	80
147	The role of the peroxisome proliferator-activated receptor-alpha (PPAR-alpha) in the regulation of acute inflammation. <i>Journal of Leukocyte Biology</i> , 2006 , 79, 999-1010	6.5	79
146	TNF-alpha blockage in a mouse model of SCI: evidence for improved outcome. <i>Shock</i> , 2008 , 29, 32-41	3.4	75

145	Increased GILZ expression in transgenic mice up-regulates Th-2 lymphokines. <i>Blood</i> , 2006 , 107, 1039-47	2.2	72
144	Myrtucommulone from Myrtus communis exhibits potent anti-inflammatory effectiveness in vivo. Journal of Pharmacology and Experimental Therapeutics, 2009 , 329, 76-86	4.7	64
143	Inducible nitric oxide synthase mediates bone loss in ovariectomized mice. <i>Endocrinology</i> , 2003 , 144, 1098-107	4.8	64
142	Green tea polyphenol extract attenuates colon injury induced by experimental colitis. <i>Free Radical Research</i> , 2005 , 39, 1017-25	4	60
141	Inhibition of the nuclear factor-kappaB activation with pyrrolidine dithiocarbamate attenuating inflammation and oxidative stress after experimental spinal cord trauma in rats. <i>Journal of Neurosurgery: Spine</i> , 2004 , 1, 311-21	2.8	60
140	Absence of endogenous interleukin-10 enhances secondary inflammatory process after spinal cord compression injury in mice. <i>Journal of Neurochemistry</i> , 2009 , 108, 1360-72	6	59
139	Glycogen synthase kinase-3 beta inhibition reduces secondary damage in experimental spinal cord trauma. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2006 , 318, 79-89	4.7	59
138	Inhibitors of poly(ADP-ribose) polymerase modulate signal transduction pathways and secondary damage in experimental spinal cord trauma. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2005 , 312, 449-57	4.7	58
137	Effects of combination M40403 and dexamethasone therapy on joint disease in a rat model of collagen-induced arthritis. <i>Arthritis and Rheumatism</i> , 2005 , 52, 1929-40		56
136	Rosiglitazone, a ligand of the peroxisome proliferator-activated receptor-gamma, reduces acute pancreatitis induced by cerulein. <i>Intensive Care Medicine</i> , 2004 , 30, 951-6	14.5	54
135	Inhibition or knock out of inducible nitric oxide synthase result in resistance to bleomycin-induced lung injury. <i>Respiratory Research</i> , 2005 , 6, 58	7.3	53
134	Effects of Tempol, a membrane-permeable radical scavenger, in a rodent model periodontitis. Journal of Clinical Periodontology, 2005 , 32, 1062-8	7.7	52
133	Melatonin reduces stress-activated/mitogen-activated protein kinases in spinal cord injury. <i>Journal of Pineal Research</i> , 2009 , 46, 79-86	10.4	50
132	Erythropoietin reduces the development of experimental inflammatory bowel disease. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2004 , 311, 1272-80	4.7	50
131	Treatment with a novel poly(ADP-ribose) glycohydrolase inhibitor reduces development of septic shock-like syndrome induced by zymosan in mice. <i>Critical Care Medicine</i> , 2004 , 32, 1365-74	1.4	49
130	Reduction of ischemic brain injury by administration of palmitoylethanolamide after transient middle cerebral artery occlusion in rats. <i>Brain Research</i> , 2012 , 1477, 45-58	3.7	48
129	Modulation of NADPH oxidase activation in cerebral ischemia/reperfusion injury in rats. <i>Brain Research</i> , 2011 , 1372, 92-102	3.7	48
128	Protective effect of orally administered carnosine on bleomycin-induced lung injury. <i>American Journal of Physiology - Lung Cellular and Molecular Physiology</i> , 2007 , 292, L1095-104	5.8	48

127	Erythropoietin reduces the degree of arthritis caused by type II collagen in the mouse. <i>Arthritis and Rheumatism</i> , 2005 , 52, 940-50		47
126	Effect of 17beta-estradiol on signal transduction pathways and secondary damage in experimental spinal cord trauma. <i>Shock</i> , 2008 , 29, 362-71	3.4	47
125	Natural almond skin reduced oxidative stress and inflammation in an experimental model of inflammatory bowel disease. <i>International Immunopharmacology</i> , 2011 , 11, 915-24	5.8	46
124	Melatonin regulates matrix metalloproteinases after traumatic experimental spinal cord injury. Journal of Pineal Research, 2008, 45, 149-56	10.4	46
123	Pyrrolidine dithiocarbamate reduces the severity of cerulein-induced murine acute pancreatitis. <i>Shock</i> , 2003 , 20, 544-50	3.4	46
122	Rosiglitazone, a ligand of the peroxisome proliferator-activated receptor-gamma, reduces the development of nonseptic shock induced by zymosan in mice. <i>Critical Care Medicine</i> , 2004 , 32, 457-66	1.4	45
121	PARG activity mediates intestinal injury induced by splanchnic artery occlusion and reperfusion. <i>FASEB Journal</i> , 2005 , 19, 558-66	0.9	45
120	The Antioxidant and Anti-Inflammatory Properties of L. Cashew Nuts in a Mouse Model of Colitis. <i>Nutrients</i> , 2020 , 12,	6.7	44
119	Modulation of nitric oxide homeostasis in a mouse model of spinal cord injury. <i>Journal of Neurosurgery: Spine</i> , 2006 , 4, 145-53	2.8	44
118	5-Lipoxygenase modulates colitis through the regulation of adhesion molecule expression and neutrophil migration. <i>Laboratory Investigation</i> , 2005 , 85, 808-22	5.9	44
117	Post-ischaemic thyroid hormone treatment in a rat model of acute stroke. <i>Brain Research</i> , 2013 , 1513, 92-102	3.7	43
116	The renal injury and inflammation caused by ischemia-reperfusion are reduced by genetic inhibition of TNF-R1: a comparison with infliximab treatment. <i>European Journal of Pharmacology</i> , 2013 , 700, 134-4	4€ ^{.3}	42
115	Role of endogenous ligands for the peroxisome proliferators activated receptors alpha in the secondary damage in experimental spinal cord trauma. <i>Experimental Neurology</i> , 2005 , 194, 267-78	5.7	42
114	Green tea polyphenol extract attenuates lung injury in experimental model of carrageenan-induced pleurisy in mice. <i>Respiratory Research</i> , 2005 , 6, 66	7.3	41
113	The selective adenosine A2A receptor agonist CGS 21680 reduces JNK MAPK activation in oligodendrocytes in injured spinal cord. <i>Shock</i> , 2009 , 32, 578-85	3.4	40
112	Effects of combination of melatonin and dexamethasone on secondary injury in an experimental mice model of spinal cord trauma. <i>Journal of Pineal Research</i> , 2007 , 43, 140-53	10.4	40
111	Effects of GW274150, a novel and selective inhibitor of iNOS activity, in acute lung inflammation. <i>British Journal of Pharmacology</i> , 2004 , 141, 979-87	8.6	38
110	Reduction in the development of cerulein-induced acute pancreatitis by treatment with M40401, a new selective superoxide dismutase mimetic. <i>Shock</i> , 2004 , 22, 254-61	3.4	38

(2007-2007)

109	Genetic and pharmacological inhibition of GITR-GITRL interaction reduces chronic lung injury induced by bleomycin instillation. <i>FASEB Journal</i> , 2007 , 21, 117-29	0.9	37
108	Erythropoietin reduces the development of nonseptic shock induced by zymosan in mice. <i>Critical Care Medicine</i> , 2006 , 34, 1168-77	1.4	37
107	Evidence for the role of mitogen-activated protein kinase signaling pathways in the development of spinal cord injury. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2008 , 325, 100-14	4.7	36
106	Pharmacological inhibition of leukotrienes in an animal model of bleomycin-induced acute lung injury. <i>Respiratory Research</i> , 2006 , 7, 137	7.3	36
105	GPI 6150, a PARP inhibitor, reduces the colon injury caused by dinitrobenzene sulfonic acid in the rat. <i>Biochemical Pharmacology</i> , 2002 , 64, 327-37	6	36
104	Hypericum perforatum attenuates the development of cerulein-induced acute pancreatitis in mice. <i>Shock</i> , 2006 , 25, 161-7	3.4	35
103	The Role of Cashew (L.) Nuts on an Experimental Model of Painful Degenerative Joint Disease. <i>Antioxidants</i> , 2020 , 9,	7.1	34
102	PPAR-alpha modulate the anti-inflammatory effect of glucocorticoids in the secondary damage in experimental spinal cord trauma. <i>Pharmacological Research</i> , 2009 , 59, 338-50	10.2	34
101	Absence of peroxisome proliferators-activated receptors (PPAR)alpha enhanced the multiple organ failure induced by zymosan. <i>Shock</i> , 2006 , 26, 477-84	3.4	34
100	Calpain I inhibitor ameliorates the indices of disease severity in a murine model of cerulein-induced acute pancreatitis. <i>Intensive Care Medicine</i> , 2004 , 30, 1645-51	14.5	34
99	Liver X receptor agonist treatment regulates inflammatory response after spinal cord trauma. Journal of Neurochemistry, 2010 , 112, 611-24	6	33
98	Glucocorticoid-induced TNF receptor family gene (GITR) knockout mice exhibit a resistance to splanchnic artery occlusion (SAO) shock. <i>Journal of Leukocyte Biology</i> , 2004 , 76, 933-40	6.5	32
97	Inhibitors of poly(ADP-ribose) polymerase modulate signal transduction pathways and the development of bleomycin-induced lung injury. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2005 , 313, 529-38	4.7	32
96	Glycyrrhizin reduces secondary inflammatory process after spinal cord compression injury in mice. <i>Shock</i> , 2009 , 31, 367-75	3.4	31
95	WY 14643, a potent exogenous PPAR-alpha ligand, reduces intestinal injury associated with splanchnic artery occlusion shock. <i>Shock</i> , 2004 , 22, 340-6	3.4	31
94	Effects of thalidomide in a mouse model of cerulein-induced acute pancreatitis. <i>Shock</i> , 2008 , 29, 89-97	3.4	31
93	16,16-Dimethyl prostaglandin E2 efficacy on prevention and protection from bleomycin-induced lung injury and fibrosis. <i>American Journal of Respiratory Cell and Molecular Biology</i> , 2009 , 41, 50-8	5.7	30
92	Beneficial effects of FeTSPP, a peroxynitrite decomposition catalyst, in a mouse model of spinal cord injury. <i>Free Radical Biology and Medicine</i> , 2007 , 43, 763-80	7.8	30

91	Effects of 3-aminobenzamide, an inhibitor of poly (ADP-ribose) polymerase, in a mouse model of acute pancreatitis induced by cerulein. <i>European Journal of Pharmacology</i> , 2006 , 549, 149-56	5.3	30
90	Etanercept attenuates the development of cerulein-induced acute pancreatitis in mice: a comparison with TNF-alpha genetic deletion. <i>Shock</i> , 2007 , 27, 542-51	3.4	30
89	Increased oxidative-related mechanisms in the spinal cord injury in old rats. <i>Neuroscience Letters</i> , 2006 , 393, 141-6	3.3	30
88	Pyrrolidine dithiocarbamate attenuates the development of organ failure induced by zymosan in mice. <i>Intensive Care Medicine</i> , 2003 , 29, 2016-25	14.5	30
87	Ultramicronized Palmitoylethanolamide and Paracetamol, a New Association to Relieve Hyperalgesia and Pain in a Sciatic Nerve Injury Model in Rat. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	29
86	Hypericum perforatum attenuates the development of carrageenan-induced lung injury in mice. Free Radical Biology and Medicine, 2006 , 40, 740-53	7.8	29
85	Poly(ADP-ribose) glycohydrolase activity mediates post-traumatic inflammatory reaction after experimental spinal cord trauma. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2006 , 319, 127-38	4.7	29
84	Effect of anthocyanins contained in a blackberry extract on the circulatory failure and multiple organ dysfunction caused by endotoxin in the rat. <i>Planta Medica</i> , 2004 , 70, 745-52	3.1	29
83	5-lipoxygenase knockout mice exhibit a resistance to acute pancreatitis induced by cerulein. <i>Immunology</i> , 2003 , 110, 120-30	7.8	29
82	ROLE OF ENDOGENOUS AND EXOGENOUS LIGANDS FOR THE PEROXISOME PROLIFERATOR-ACTIVATED RECEPTOR alpha IN THE DEVELOPMENT OF BLEOMYCIN-INDUCED LUNG INJURY. <i>Shock</i> , 2005 , 24, 547-55	3.4	29
81	Melatonin limits lung injury in bleomycin treated mice. Journal of Pineal Research, 2005, 39, 105-12	10.4	29
80	Cashew (L.) Nuts Counteract Oxidative Stress and Inflammation in an Acute Experimental Model of Carrageenan-Induced Paw Edema. <i>Antioxidants</i> , 2020 , 9,	7.1	29
79	Treatment with green tea extract attenuates secondary inflammatory response in an experimental model of spinal cord trauma. <i>Naunyn-Schmiedebergis Archives of Pharmacology</i> , 2009 , 380, 179-92	3.4	28
78	Glucocorticoid-induced tumor necrosis factor receptor-related (GITR)-Fc fusion protein inhibits GITR triggering and protects from the inflammatory response after spinal cord injury. <i>Molecular Pharmacology</i> , 2008 , 73, 1610-21	4.3	28
77	Role of poly(ADP-ribose) glycohydrolase in the development of inflammatory bowel disease in mice. <i>Free Radical Biology and Medicine</i> , 2007 , 42, 90-105	7.8	28
76	5-Lipoxygenase knockout mice exhibit a resistance to pleurisy and lung injury caused by carrageenan. <i>Journal of Leukocyte Biology</i> , 2003 , 73, 739-46	6.5	28
75	Increased carrageenan-induced acute lung inflammation in old rats. <i>Immunology</i> , 2005 , 115, 253-61	7.8	28
74	Green tea polyphenol extract attenuates ischemia/reperfusion injury of the gut. Naunyn-Schmiedebergrs Archives of Pharmacology, 2005, 371, 364-74	3.4	28

(2004-2009)

73	Effects of a metalloporphyrinic peroxynitrite decomposition catalyst, ww-85, in a mouse model of spinal cord injury. <i>Free Radical Research</i> , 2009 , 43, 631-45	4	27
72	GW0742, a high-affinity PPAR -beta/delta agonist, inhibits acute lung injury in mice. <i>Shock</i> , 2010 , 33, 42	6 ₃ 3,5	27
71	Glycogen synthase kinase 3beta inhibition reduces the development of nonseptic shock induced by zymosan in mice. <i>Shock</i> , 2007 , 27, 97-107	3.4	27
70	Effect of cyclopentanone prostaglandin 15-deoxy-delta12,14PGJ2 on early functional recovery from experimental spinal cord injury. <i>Shock</i> , 2008 , 30, 142-52	3.4	26
69	Biochemical Evaluation of the Antioxidant Effects of Hydroxytyrosol on Pancreatitis-Associated Gut Injury. <i>Antioxidants</i> , 2020 , 9,	7.1	26
68	Protective effects of glycyrrhizin in a gut hypoxia (ischemia)-reoxygenation (reperfusion) model. <i>Intensive Care Medicine</i> , 2009 , 35, 687-97	14.5	25
67	Protective effect of melatonin against the inflammatory response elicited by crude venom from isolated nematocysts of Pelagia noctiluca (Cnidaria, Scyphozoa). <i>Journal of Pineal Research</i> , 2009 , 47, 56-69	10.4	25
66	Anti-inflammatory and anti-apoptotic effects of fumonisin B1, an inhibitor of ceramide synthase, in a rodent model of splanchnic ischemia and reperfusion injury. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2008 , 327, 45-57	4.7	24
65	High-density lipoproteins reduce the intestinal damage associated with ischemia/reperfusion and colitis. <i>Shock</i> , 2004 , 21, 342-51	3.4	24
64	Beneficial effects of ethyl pyruvate in a mouse model of spinal cord injury. Shock, 2009, 32, 217-27	3.4	23
63	Green tea polyphenols ameliorate pancreatic injury in cerulein-induced murine acute pancreatitis. <i>Pancreas</i> , 2009 , 38, 954-67	2.6	23
62	Role of endogenous glutathione in the secondary damage in experimental spinal cord injury in mice. <i>Neuroscience Letters</i> , 2007 , 423, 41-6	3.3	23
61	Involvement of 5-lipoxygenase in spinal cord injury. <i>Journal of Neuroimmunology</i> , 2005 , 166, 55-64	3.5	23
60	PPAR-alpha contributes to the anti-inflammatory activity of 17beta-estradiol. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2009 , 331, 796-807	4.7	22
59	Protective effect of Hypericum perforatum in zymosan-induced multiple organ dysfunction syndrome: relationship to its inhibitory effect on nitric oxide production and its peroxynitrite scavenging activity. <i>Nitric Oxide - Biology and Chemistry</i> , 2007 , 16, 118-30	5	22
58	Neuroprotection and enhanced recovery with hypericum perforatum extract after experimental spinal cord injury in mice. <i>Shock</i> , 2006 , 25, 608-17	3.4	22
57	Ethyl pyruvate reduces the development of zymosan-induced generalized inflammation in mice. <i>Critical Care Medicine</i> , 2009 , 37, 270-82	1.4	21
56	Role of 5-lipoxygenase in the multiple organ failure induced by zymosan. <i>Intensive Care Medicine</i> , 2004 , 30, 1935-43	14.5	21

55	Treatment with PARP-1 inhibitors, GPI 15427 or GPI 16539, ameliorates intestinal damage in rat models of colitis and shock. <i>European Journal of Pharmacology</i> , 2005 , 527, 163-71	5.3	21
54	Consumption of . (Cashew Nuts) Inhibits Oxidative Stress through Modulation of the Nrf2/HO-1 and NF-kB Pathways. <i>Molecules</i> , 2020 , 25,	4.8	21
53	Thymosin A reduces IL-17-producing cells and IL-17 expression, and protects lungs from damage in bleomycin-treated mice. <i>Immunobiology</i> , 2014 , 219, 425-31	3.4	20
52	MK801 attenuates secondary injury in a mouse experimental compression model of spinal cord trauma. <i>BMC Neuroscience</i> , 2011 , 12, 31	3.2	20
51	Liver X receptor agonist treatment reduced splanchnic ischemia and reperfusion injury. <i>Journal of Leukocyte Biology</i> , 2010 , 87, 309-21	6.5	20
50	Effects of Hypericum perforatum extract in a rat model of ischemia and reperfusion injury. <i>Shock</i> , 2005 , 24, 255-63	3.4	20
49	Inhibition of tyrosine-kinase-mediated cellular signaling by tyrphostins AG 126 and AG556 modulates murine experimental acute pancreatitis. <i>Surgery</i> , 2005 , 138, 913-23	3.6	20
48	Synergistic interaction between methotrexate and a superoxide dismutase mimetic: pharmacologic and potential clinical significance. <i>Arthritis and Rheumatism</i> , 2005 , 52, 3755-60		20
47	Adelmidrol: A New Promising Antioxidant and Anti-Inflammatory Therapeutic Tool in Pulmonary Fibrosis. <i>Antioxidants</i> , 2020 , 9,	7.1	19
46	Absence of endogenous interleukin-10 enhances the evolution of acute lung injury. <i>European Cytokine Network</i> , 2002 , 13, 285-97	3.3	19
45	Inhibition of ceramide biosynthesis ameliorates pathological consequences of spinal cord injury. <i>Shock</i> , 2009 , 31, 634-44	3.4	18
44	Role of PPAR-delta in the development of zymosan-induced multiple organ failure: an experiment mice study. <i>Journal of Inflammation</i> , 2010 , 7, 12	6.7	18
43	Role of peroxisome proliferator-activated receptor-alpha in acute pancreatitis induced by cerulein. <i>Immunology</i> , 2006 , 118, 559-70	7.8	18
42	5-Aminoisoquinolinone reduces colon injury by experimental colitis. <i>Naunyn-Schmiedebergi</i> s <i>Archives of Pharmacology</i> , 2004 , 370, 464-73	3.4	18
41	Effects of glycogen synthase kinase-3beta inhibition on the development of cerulein-induced acute pancreatitis in mice. <i>Critical Care Medicine</i> , 2007 , 35, 2811-21	1.4	17
40	Reduced development of experimental periodontitis by treatment with M40403, a superoxide dismutase mimetic. <i>European Journal of Pharmacology</i> , 2005 , 516, 151-7	5.3	17
39	In vitro and in vivo properties of a fully human IgG1 monoclonal antibody that combats multidrug resistant Pseudomonas aeruginosa. <i>International Journal of Molecular Medicine</i> , 2012 , 30, 455-64	4.4	16
38	Thymosin A protects C57BL/6 mice from bleomycin-induced damage in the lung. <i>European Journal of Clinical Investigation</i> , 2013 , 43, 309-15	4.6	16

(2020-2020)

37	Cashew L.) Nuts Modulate the Nrf2 and NLRP3 Pathways in Pancreas and Lung after Induction of Acute Pancreatitis by Cerulein. <i>Antioxidants</i> , 2020 , 9,	7.1	16
36	Protective effects of thymosin A in a mouse model of lung fibrosis. <i>Annals of the New York Academy of Sciences</i> , 2012 , 1269, 69-73	6.5	15
35	5-lipoxygenase knockout mice exhibit a resistance to splanchnic artery occlusion shock. <i>Shock</i> , 2003 , 20, 230-6	3.4	14
34	Beneficial effects of 5-aminoisoquinolinone, a novel, potent, water-soluble, inhibitor of poly (ADP-ribose) polymerase, in a rat model of splanchnic artery occlusion and reperfusion. <i>European Journal of Pharmacology</i> , 2004 , 492, 203-10	5.3	14
33	Protective effects of M40401, a selective superoxide dismutase mimetic, on zymosan-induced nonseptic shock. <i>Critical Care Medicine</i> , 2004 , 32, 157-67	1.4	14
32	Absence of endogenous interleukin-10 enhanced organ dysfunction and mortality associated to zymosan-induced multiple organ dysfunction syndrome. <i>Cytokine</i> , 2008 , 41, 136-43	4	13
31	Effect of thalidomide on signal transduction pathways and secondary damage in experimental spinal cord trauma. <i>Shock</i> , 2008 , 30, 231-40	3.4	13
30	Fumonisin b1 reduces the development of multiple organ failure induced by zymosan in mice. <i>Shock</i> , 2009 , 31, 170-7	3.4	12
29	Effects of thymosin A and its N-terminal fragment Ac-SDKP on TGF-Etreated human lung fibroblasts and in the mouse model of bleomycin-induced lung fibrosis. <i>Expert Opinion on Biological Therapy</i> , 2015 , 15 Suppl 1, S211-21	5.4	11
28	Neuroprotective effects of olprinone after cerebral ischemia/reperfusion injury in rats. <i>Neuroscience Letters</i> , 2011 , 503, 93-9	3.3	11
27	PEA/Polydatin: Anti-Inflammatory and Antioxidant Approach to Counteract DNBS-Induced Colitis. <i>Antioxidants</i> , 2021 , 10,	7.1	11
26	AQX-1125, small molecule SHIP1 activator inhibits bleomycin-induced pulmonary fibrosis. <i>British Journal of Pharmacology</i> , 2017 , 174, 3045-3057	8.6	10
25	Beneficial effects of GW274150 treatment on the development of experimental colitis induced by dinitrobenzene sulfonic acid. <i>European Journal of Pharmacology</i> , 2005 , 507, 281-9	5.3	10
24	Inhibition of tyrosine kinase-mediated cellular signalling by Tyrphostins AG126 and AG556 modulates secondary damage in experimental spinal cord trauma. <i>Neuropharmacology</i> , 2007 , 52, 1454-	7 ^{§.5}	9
23	Prevention of carrageenan-induced pleurisy in mice by anti-CD30 ligand monoclonal antibody. <i>Clinical Immunology</i> , 2004 , 113, 64-73	9	9
22	The cyclopentenone prostaglandin 15-deoxydelta(12,14)-prostaglandin J2 attenuates the development of zymosan-induced shock. <i>Intensive Care Medicine</i> , 2005 , 31, 693-700	14.5	9
21	The Protective Effects of Pre- and Post-Administration of Micronized Palmitoylethanolamide Formulation on Postoperative Pain in Rats. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	9
20	Protective effects of Colomast, A New Formulation of Adelmidrol and Sodium Hyaluronate, in A Mouse Model of Acute Restraint Stress. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	8

19	Recombinant human activated protein C (Xigris) attenuates murine cerulein-induced acute pancreatitis via regulation of nuclear factor B and apoptotic pathways. <i>Pancreas</i> , 2012 , 41, 619-28	2.6	8
18	N-benzyloxycarbonyl-Val-Ala-Asp-fluoromethylketone reduces severity of experimental spinal cord injury. <i>Shock</i> , 2007 , 27, 258-65	3.4	8
17	ROLE of endogenous peroxisome proliferator-activated receptor-alpha (PPAR-alpha) ligands in the development of gut ischemia and reperfusion in mice. <i>Shock</i> , 2006 , 25, 17-22	3.4	8
16	Methylguanidine reduces the development of non septic shock induced by zymosan in mice. <i>Life Sciences</i> , 2004 , 75, 1417-33	6.8	8
15	Efficacy of treatment with verbascoside, biotechnologically produced by Syringa vulgaris plant cell cultures in an experimental mice model of spinal cord trauma. <i>Naunyn-Schmiedebergis Archives of Pharmacology</i> , 2010 , 382, 331-45	3.4	6
14	Mucosa-Associated Lymphoid Tissue Lymphoma Translocation 1 Inhibitor as a Novel Therapeutic Tool for Lung Injury. <i>International Journal of Molecular Sciences</i> , 2020 , 21,	6.3	5
13	The Methyl Ester of 2-Cyano-3,12-Dioxooleana-1,9-Dien-28-Oic Acid Reduces Endometrial Lesions Development by Modulating the NFkB and Nrf2 Pathways. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	5
12	Etanercept reduces acute tissue injury and mortality associated to zymosan-induced multiple organ dysfunction syndrome. <i>Shock</i> , 2008 , 29, 560-71	3.4	4
11	Novel Combination of COX-2 Inhibitor and Antioxidant Therapy for Modulating Oxidative Stress Associated with Intestinal Ischemic Reperfusion Injury and Endotoxemia. <i>Antioxidants</i> , 2020 , 9,	7.1	4
10	Inhibition of P2X7 Purinergic Receptor Ameliorates Fibromyalgia Syndrome by Suppressing NLRP3 Pathway. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	4
9	Palmitoylethanolamide/Baicalein Regulates the Androgen Receptor Signaling and NF- B /Nrf2 Pathways in Benign Prostatic Hyperplasia. <i>Antioxidants</i> , 2021 , 10,	7.1	4
8	Atrazine Inhalation Causes Neuroinflammation, Apoptosis and Accelerating Brain Aging. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	4
7	Epigallocatechin-3-Gallate Modulates Postoperative Pain by Regulating Biochemical and Molecular Pathways. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	3
6	Molecular and Biochemical Mechanism of Cannabidiol in the Management of the Inflammatory and Oxidative Processes Associated with Endometriosis. <i>International Journal of Molecular Sciences</i> , 2022 , 23, 5427	6.3	3
5	Autophagy and Mitophagy Promotion in a Rat Model of Endometriosis. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	2
4	Regulation of Inflammatory and Proliferative Pathways by Fotemustine and Dexamethasone in Endometriosis. <i>International Journal of Molecular Sciences</i> , 2021 , 22,	6.3	1
3	Role of Etanercept and Infliximab on Nociceptive Changes Induced by the Experimental Model of Fibromyalgia. <i>International Journal of Molecular Sciences</i> , 2022 , 23, 6139	6.3	1
2	Effects of glycogen synthase kinase-3[beta] inhibition on the development of cerulein-induced acute pancreatitis in mice *. <i>Critical Care Medicine</i> , 2007 , 35, 2811-2821	1.4	

Evidence for the role of PPAR-/ In the development of spinal cord injury. FASEB Journal, 2010, 24, lb461

0.9