

Hyoe Takata

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3994113/publications.pdf>

Version: 2024-02-01

29
papers

516
citations

516710

16
h-index

677142

22
g-index

32
all docs

32
docs citations

32
times ranked

437
citing authors

#	ARTICLE	IF	CITATIONS
1	Factors controlling dissolved ¹³⁷ Cs activities in coastal waters on the eastern and western sides of Honshu, Japan. <i>Science of the Total Environment</i> , 2022, 806, 151216.	8.0	2
2	A comparative study of riverine ¹³⁷ Cs dynamics during high-flow events at three contaminated river catchments in Fukushima. <i>Science of the Total Environment</i> , 2022, 821, 153408.	8.0	5
3	Temporal variability of ¹³⁷ Cs concentrations in coastal sediments off Fukushima. <i>Science of the Total Environment</i> , 2022, 831, 154670.	8.0	2
4	Environmental recovery from ¹³⁷ Cs contamination in Japanese coastal waters shown by comparison of temporal distributions with European seas. <i>Journal of Environmental Radioactivity</i> , 2022, 251-252, 106961.	1.7	2
5	Importance of desorption process from Abukuma River's suspended particles in increasing dissolved ¹³⁷ Cs in coastal water during river-flood caused by typhoons. <i>Chemosphere</i> , 2021, 281, 130751.	8.2	11
6	The contribution of ¹³⁷ Cs export flux from the Tone River Japan to the marine environment. <i>Science of the Total Environment</i> , 2020, 701, 134550.	8.0	16
7	Temporal trends of ¹³⁷ Cs concentration in seawaters and bottom sediments in coastal waters around Japan: implications for the Kd concept in the dynamic marine environment. <i>Journal of Radioanalytical and Nuclear Chemistry</i> , 2020, 323, 567-580.	1.5	20
8	Suspended Particleâ€“Water Interactions Increase Dissolved ¹³⁷ Cs Activities in the Nearshore Seawater during Typhoon Hagibis. <i>Environmental Science & Technology</i> , 2020, 54, 10678-10687.	10.0	27
9	A 30-year record reveals re-equilibration rates of ¹³⁷ Cs in marine biota after the Fukushima Dai-ichi nuclear power plant accident: Concentration ratios in pre- and post-event conditions. <i>Science of the Total Environment</i> , 2019, 675, 694-704.	8.0	20
10	Appearances of Fukushima Daiichi Nuclear Power Plant-Derived ¹³⁷ Cs in Coastal Waters around Japan: Results from Marine Monitoring off Nuclear Power Plants and Facilities, 1983â€“2016. <i>Environmental Science & Technology</i> , 2018, 52, 2629-2637.	10.0	31
11	Radiocesium in the swash zones off the coast of the Japan Sea. <i>Applied Radiation and Isotopes</i> , 2018, 141, 64-67.	1.5	0
12	Temporal variation of cesium isotope concentrations and atom ratios in zooplankton in the Pacific off the east coast of Japan. <i>Scientific Reports</i> , 2017, 7, 39874.	3.3	20
13	Decline in radiocesium in seafloor sediments off Fukushima and nearby prefectures. <i>Journal of Oceanography</i> , 2017, 73, 529-545.	1.7	37
14	The Contribution of Sources to the Sustained Elevated Inventory of ¹³⁷ Cs in Offshore Waters East of Japan after the Fukushima Dai-ichi Nuclear Power Station Accident. <i>Environmental Science & Technology</i> , 2016, 50, 6957-6963.	10.0	24
15	A new approach to evaluate factors controlling elemental sedimentâ€“seawater distribution coefficients (Kd) in coastal regions, Japan. <i>Science of the Total Environment</i> , 2016, 543, 315-325.	8.0	14
16	Remobilization of radiocesium on riverine particles in seawater: The contribution of desorption to the export flux to the marine environment. <i>Marine Chemistry</i> , 2015, 176, 51-63.	2.3	41
17	Bromine and iodine in Japanese soils determined with polarizing energy dispersive X-ray fluorescence spectrometry. <i>Soil Science and Plant Nutrition</i> , 2015, 61, 751-760.	1.9	22
18	Radiocesiums (¹³⁴ Cs, ¹³⁷ Cs) in zooplankton in the waters of Miyagi, Fukushima and Ibaraki Prefectures. <i>Journal of Radioanalytical and Nuclear Chemistry</i> , 2015, 303, 1265-1271.	1.5	8

#	ARTICLE	IF	CITATIONS
19	Distributions of Pu isotopes in seawater and bottom sediments in the coast of the Japanese archipelago before and soon after the Fukushima Dai-ichi Nuclear Power Station accident. <i>Journal of Environmental Radioactivity</i> , 2015, 142, 113-123.	1.7	19
20	Distribution coefficients (Kd) of strontium and significance of oxides and organic matter in controlling its partitioning in coastal regions of Japan. <i>Science of the Total Environment</i> , 2014, 490, 979-986.	8.0	18
21	A sensitive and simple analytical method for the determination of stable Cs in estuarine and coastal waters. <i>Analytical Methods</i> , 2013, 5, 2558.	2.7	10
22	Distribution coefficients (Kd) of stable iodine in estuarine and coastal regions, Japan, and their relationship to salinity and organic carbon in sediments. <i>Environmental Monitoring and Assessment</i> , 2013, 185, 3645-3658.	2.7	15
23	Long Term Temporal Changes of ^{90}Sr and ^{137}Cs in Seawater, Bottom Sediment and Marine Organism Samples - from the Chernobyl Accident to Immediately after the Fukushima Accident -. <i>Bunseki Kagaku</i> , 2013, 62, 455-474.	0.2	21
24	Distributions of trace metals Co, Cu and Cd in northern Sagami Bay, Japan and their relationship to estuarine variables. <i>Estuarine, Coastal and Shelf Science</i> , 2012, 111, 84-94.	2.1	10
25	Rapid determination of total iodine in Japanese coastal seawater using SF-ICP-MS. <i>Microchemical Journal</i> , 2012, 100, 42-47.	4.5	37
26	Determination of ^{232}Th in seawater by ICP-MS after preconcentration and separation using a chelating resin. <i>Talanta</i> , 2011, 85, 1772-1777.	5.5	26
27	Determination of naturally occurring uranium concentrations in seawater, sediment, and marine organisms in Japanese estuarine areas. <i>Journal of Radioanalytical and Nuclear Chemistry</i> , 2011, 287, 795-799.	1.5	23
28	Processes controlling cobalt distribution in two temperate estuaries, Sagami Bay and Wakasa Bay, Japan. <i>Estuarine, Coastal and Shelf Science</i> , 2010, 89, 294-305.	2.1	20
29	Sediment-Water Distribution Coefficients of Stable Elements in Four Estuarine Areas in Japan. <i>Journal of Nuclear Science and Technology</i> , 2010, 47, 111-122.	1.3	14