

Edinaldo Nelson dos Santos-Silva

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3995188/publications.pdf>

Version: 2024-02-01

21
papers

127
citations

1478505

6
h-index

1474206

9
g-index

22
all docs

22
docs citations

22
times ranked

125
citing authors

#	ARTICLE	IF	CITATIONS
1	Loss of a freshwater copepod species from El Junco Lake, Galápagos following the introduction and eradication of the Nile tilapia. <i>Aquatic Conservation: Marine and Freshwater Ecosystems</i> , 2021, 31, 3651-3656.	2.0	1
2	Taxonomic notes on the typology of four species of genus <i>Notodiaptomus</i> Kiefer, 1936 (Copepoda: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5	0.5	0
3	Class Copepoda. , 2020, , 663-796.		6
4	An annotated checklist of freshwater Copepoda (Crustacea, Hexanauplia) from continental Ecuador and the Galapagos Archipelago. <i>ZooKeys</i> , 2019, 871, 55-77.	1.1	5
5	<i>Eunotia amazonica</i> sp. nov. (Bacillariophyta), a common stalk-forming species from the Rio Negro basin (Brazilian Amazon). <i>European Journal of Phycology</i> , 2018, 53, 166-179.	2.0	5
6	Composition, abundance, and diversity of limnetic cladocerans (Crustacea: Anomopoda and) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 542	0.3	1
7	Distribution of European and African species of genus <i>Diaptomus</i> (Copepoda: Calanoida: Diaptomidae): a track analysis. <i>Nauplius</i> , 2018, 26, .	0.3	1
8	Revision of the <i>Remaneicaris argentina</i> -group (Copepoda, Harpacticoida, Parastenocarididae): supplementary description of species, and description of the first semi-terrestrial <i>Remaneicaris</i> from the tropical forest of Southeast Mexico. <i>Zootaxa</i> , 2017, 4238, 499.	0.5	7
9	A new species of <i>Notodiaptomus</i> from the Ecuadorian Andes (Copepoda, Calanoida, Diaptomidae). <i>ZooKeys</i> , 2017, 697, 59-71.	1.1	1
10	A new genus of Parastenocarididae (Copepoda, Harpacticoida) from the Tocantins River basin (GoiÃs,) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 542	1.1	7
11	The density and biomass of mesozooplankton and ichthyoplankton in the Negro and the Amazon Rivers during the rainy season: the ecological importance of the confluence boundary. <i>PeerJ</i> , 2017, 5, e3308.	2.0	4
12	New Diaptomidae records (Crustacea: Copepoda: Calanoida: Diaptomidae) in the Neotropical region. <i>Check List</i> , 2013, 9, 700.	0.4	11
13	Composition, species richness and patterns of nycthemeral vertical distribution of planktonic cladocereans in a black water Amazonian lake. <i>Nauplius</i> , 2011, 19, 109-122.	0.3	13
14	A new <i>Argyrodiaptomus</i> (Copepoda: Calanoida: Diaptomidae) from the southwestern Brazilian Amazon. <i>Zootaxa</i> , 2007, 1518, 1-29.	0.5	3
15	The Neotropical Genus <i>Notodiaptomus</i> Kiefer, 1936 (Calanoida: Diaptomidae): Redescription of the Type Species <i>Notodiaptomus deitersi</i> (Poppe, 1891) and Designation of a Neotype. <i>Studies on Neotropical Fauna and Environment</i> , 1999, 34, 114-128.	1.0	11
16	Redescription and distribution of <i>Mastigodiaptomus montezumae</i> (Copepoda, Calanoida, Diaptomidae) in Mexico. <i>Hydrobiologia</i> , 1996, 328, 207-213.	2.0	11
17	Some littoral anomopods (Crustacea) from Central Amazonia. <i>Hydrobiologia</i> , 1995, 315, 227-230.	2.0	12
18	<i>Keraladiaptomus rangareddyi</i> a new genus and new species of Diaptominae (Copepoda, Calanoida,) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 542	2.0	1

#	ARTICLE	IF	CITATIONS
19	A NEW SPECIES OF <i>Rhacodiaptomus</i> Kiefer, 1936, FROM THE BRAZILIAN AMAZON (CRUSTACEA:COPEPODA:) Tj ET Oq 1 1 0.784314 r g B 0.71 2	0.71	2
20	Revision of <i>Brasilibathynellocaris</i> Jakobi, 1972 (Copepoda: Harpacticoida: Parastenocarididae) with redefinition of the genus. <i>Zoological Journal of the Linnean Society</i> , 0, 159, 527-566.	2.3	18
21	A new genus of Parastenocarididae Chappuis, 1940 (Copepoda: Harpacticoida) from the Amazonian Region, Brazil, with close affinity to <i>Murunducaris</i> Reid, 1994. <i>Nauplius</i> , 0, 29, .	0.3	0