

Taro Nakamura

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3996774/publications.pdf>

Version: 2024-02-01

46
papers

1,130
citations

361045

20
h-index

414034

32
g-index

48
all docs

48
docs citations

48
times ranked

773
citing authors

#	ARTICLE	IF	CITATIONS
1	Quick-Freeze, Deep-Etch Electron Microscopy Reveals the Characteristic Architecture of the Fission Yeast Spore. <i>Journal of Fungi (Basel, Switzerland)</i> , 2021, 7, 7.	1.5	6
2	Meiosis-specific localization of the exocytic Rab Ypt2 in fission yeast. <i>Small GTPases</i> , 2020, 11, 146-154.	0.7	3
3	The Fission Yeast RNA-Binding Protein Meu5 Is Involved in Outer Forespore Membrane Breakdown during Spore Formation. <i>Journal of Fungi (Basel, Switzerland)</i> , 2020, 6, 284.	1.5	4
4	The asymmetric chemical structures of two mating pheromones reflect their differential roles in mating of fission yeast. <i>Journal of Cell Science</i> , 2019, 132, .	1.2	11
5	The fission yeast SPB component Dms1 is required to initiate forespore membrane formation and maintain meiotic SPB components. <i>PLoS ONE</i> , 2018, 13, e0197879.	1.1	1
6	The exocytic Rabs Ypt3 and Ypt2 regulate the early step of biogenesis of the spore plasma membrane in fission yeast. <i>Molecular Biology of the Cell</i> , 2016, 27, 3317-3328.	0.9	7
7	C3-P-05Spore surface of <i>Schizosaccharomyces pombe</i> visualized by Quick-Freeze and Deep-Etch (QFDE) replica electron microscopy. <i>Microscopy (Oxford, England)</i> , 2015, 64, i128.2-i128.	0.7	0
8	Molecular coevolution of a sex pheromone and its receptor triggers reproductive isolation in <i>Schizosaccharomyces pombe</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015, 112, 4405-4410.	3.3	28
9	The fission yeast spore is coated by a proteinaceous surface layer comprising mainly Isp3. <i>Molecular Biology of the Cell</i> , 2014, 25, 1549-1559.	0.9	16
10	The meiosis-specific nuclear passenger protein is required for proper assembly of forespore membrane in fission yeast. <i>Journal of Cell Science</i> , 2014, 127, 4429-42.	1.2	4
11	The Fission Yeast Synaptobrevin Ortholog Syb1 Plays an Important Role in Forespore Membrane Formation and Spore Maturation. <i>Eukaryotic Cell</i> , 2013, 12, 1162-1170.	3.4	4
12	Distal and Proximal Actions of Peptide Pheromone M-Factor Control Different Conjugation Steps in Fission Yeast. <i>PLoS ONE</i> , 2013, 8, e69491.	1.1	14
13	Remarkably Simple Sequence Requirement of the M-Factor Pheromone of <i>Schizosaccharomyces pombe</i> . <i>Genetics</i> , 2012, 191, 815-825.	1.2	18
14	The fission yeast pleckstrin homology domain protein Spo7 is essential for initiation of forespore membrane assembly and spore morphogenesis. <i>Molecular Biology of the Cell</i> , 2011, 22, 3442-3455.	0.9	9
15	Endocytosis is essential for dynamic translocation of a syntaxin 1 orthologue during fission yeast meiosis. <i>Molecular Biology of the Cell</i> , 2011, 22, 3658-3670.	0.9	16
16	Autophagy in the fission yeast <i>Schizosaccharomyces pombe</i> . <i>FEBS Letters</i> , 2010, 584, 1327-1334.	1.3	43
17	Role of Septins in the Orientation of Forespore Membrane Extension during Sporulation in Fission Yeast. <i>Molecular and Cellular Biology</i> , 2010, 30, 2057-2074.	1.1	38
18	<i>Schizosaccharomyces pombe</i> Calmodulin, Cam1, Plays a Crucial Role in Sporulation by Recruiting and Stabilizing the Spindle Pole Body Components Responsible for Assembly of the Forespore Membrane. <i>Eukaryotic Cell</i> , 2010, 9, 1925-1935.	3.4	12

#	ARTICLE	IF	CITATIONS
19	Mannosylinositol phosphorylceramide is a major sphingolipid component and is required for proper localization of plasma-membrane proteins in <i>Schizosaccharomyces pombe</i> . <i>Journal of Cell Science</i> , 2010, 123, 1578-1587.	1.2	47
20	NBRP databases: databases of biological resources in Japan. <i>Nucleic Acids Research</i> , 2010, 38, D26-D32.	6.5	44
21	The <i>Schizosaccharomyces pombe</i> Syntaxin 1 Homolog, Psy1, Is Essential in the Development of the Forespore Membrane. <i>Bioscience, Biotechnology and Biochemistry</i> , 2009, 73, 339-345.	0.6	24
22	Ectopic Overproduction of a Sporulation-Specific Transcription Factor Induces Assembly of Prespore-Like Membranous Compartments in Vegetative Cells of Fission Yeast. <i>Genetics</i> , 2009, 183, 1195-1199.	1.2	6
23	Autophagy-deficient <i>Schizosaccharomyces pombe</i> mutants undergo partial sporulation during nitrogen starvation. <i>Microbiology (United Kingdom)</i> , 2009, 155, 3816-3826.	0.7	63
24	Two Fission Yeast Rab7 Homologs, Ypt7 and Ypt71, Play Antagonistic Roles in the Regulation of Vacuolar Morphology. <i>Traffic</i> , 2009, 10, 912-924.	1.3	34
25	Live Observation of Forespore Membrane Formation in Fission Yeast. <i>Molecular Biology of the Cell</i> , 2008, 19, 3544-3553.	0.9	39
26	Meiotic Spindle Pole Bodies Acquire the Ability to Assemble the Spore Plasma Membrane by Sequential Recruitment of Sporulation-specific Components in Fission Yeast. <i>Molecular Biology of the Cell</i> , 2008, 19, 2476-2487.	0.9	29
27	Geranylgeranyl Diphosphate Synthase in Fission Yeast Is a Heteromer of Farnesyl Diphosphate Synthase (FPS), Fps1, and an FPS-like Protein, Spo9, Essential for Sporulation. <i>Molecular Biology of the Cell</i> , 2007, 18, 3568-3581.	0.9	34
28	Localization of Type I Myosin and F-actin to the Leading Edge Region of the Forespore Membrane in <i>Schizosaccharomyces pombe</i> . <i>Cell Structure and Function</i> , 2006, 31, 181-195.	0.5	21
29	The cation-transporting P-type ATPase Cta4 is required for assembly of the forespore membrane in fission yeast. <i>Genes and Genetic Systems</i> , 2005, 80, 317-324.	0.2	7
30	A Fission Yeast SNAP-25 Homologue, SpSec9, Is Essential for Cytokinesis and Sporulation. <i>Cell Structure and Function</i> , 2005, 30, 15-24.	0.5	29
31	A Role for Fission Yeast Rab GTPase Ypt7p in Sporulation. <i>Cell Structure and Function</i> , 2005, 30, 43-49.	0.5	21
32	Control of Late Meiosis and Ascospore Formation. , 2004, , 311-327.		23
33	Sorting nexin homologues are targets of phosphatidylinositol 3-phosphate in sporulation of <i>Schizosaccharomyces pombe</i> . <i>Genes To Cells</i> , 2004, 9, 561-574.	0.5	20
34	The Sec14 family glycerophospholipid-transfer protein is required for structural integrity of the spindle pole body during meiosis in fission yeast. <i>Genes To Cells</i> , 2004, 9, 1275-1286.	0.5	15
35	Role of phosphatidylinositol 3-phosphate in formation of forespore membrane in <i>Schizosaccharomyces pombe</i> . <i>Yeast</i> , 2003, 20, 193-206.	0.8	23
36	Ribosomal proteins S0 and S21 are involved in the stability of 18S rRNA in fission yeast, <i>Schizosaccharomyces pombe</i> . <i>Biochemical and Biophysical Research Communications</i> , 2003, 311, 942-947.	1.0	11

#	ARTICLE	IF	CITATIONS
37	The Fission Yeast <i>spo14+</i> Gene Encoding a Functional Homologue of Budding Yeast Sec12 Is Required for the Development of Forespore Membranes. <i>Molecular Biology of the Cell</i> , 2003, 14, 1109-1124.	0.9	47
38	The <i>Schizosaccharomyces pombe cdt2+</i> Gene, a Target of G1-S Phase-Specific Transcription Factor Complex DSC1, Is Required for Mitotic and Premeiotic DNA Replication. <i>Genetics</i> , 2003, 164, 881-893.	1.2	27
39	Novel Fission Yeast Cdc7-Dbf4-Like Kinase Complex Required for the Initiation and Progression of Meiotic Second Division. <i>Molecular and Cellular Biology</i> , 2002, 22, 309-320.	1.1	61
40	Irreversible deacylation of plasma membrane phospholipids by the combined action of Mg ²⁺ and a long-chain acyl-CoA synthetase inhibitor in <i>Saccharomyces cerevisiae</i> . <i>Journal of Bioscience and Bioengineering</i> , 2002, 94, 258-63.	1.1	2
41	The <i>Schizosaccharomyces pombe spo20+</i> Gene Encoding a Homologue of <i>Saccharomyces cerevisiae</i> Sec14 Plays an Important Role in Forespore Membrane Formation. <i>Molecular Biology of the Cell</i> , 2001, 12, 901-917.	0.9	74
42	The <i>Schizosaccharomyces pombe spo3+</i> Gene Is Required for Assembly of the Forespore Membrane and Genetically Interacts with <i>psy1+</i> -encoding Syntaxin-like Protein. <i>Molecular Biology of the Cell</i> , 2001, 12, 3955-3972.	0.9	131
43	Genetic evidence for the functional redundancy of the calcineurin- and Mpk1-mediated pathways in the regulation of cellular events important for growth in <i>S. pombe</i> . <i>Molecular Genetics and Genomics</i> , 1996, 251, 211.	2.4	6
44	Cloning and characterization of the <i>Saccharomyces cerevisiae</i> SVS1 gene which encodes a serine- and threonine-rich protein required for vanadate resistance. <i>Gene</i> , 1995, 165, 25-29.	1.0	19
45	Cloning and Molecular Analysis of cDNA Encoding a Carboxymethylcellulase of the Yeast <i>Cryptococcus flavus</i> . <i>Bioscience, Biotechnology and Biochemistry</i> , 1992, 56, 1230-1235.	0.6	14
46	Ca ²⁺ /calmodulin-activated protein phosphatase. <i>FEBS Letters</i> , 1992, 309, 103-106.	1.3	25