

Francisco J Acosta

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/399747/publications.pdf>

Version: 2024-02-01

20
papers

419
citations

933447

10
h-index

794594

19
g-index

21
all docs

21
docs citations

21
times ranked

669
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Nonadherence after hospital discharge in patients with schizophrenia or schizoaffective disorder: A six-month naturalistic follow-up study. <i>Comprehensive Psychiatry</i> , 2021, 108, 152240. | 3.1 | 15 |
| 2 | Prospective study of variables associated with nonadherence to psychotherapy. <i>Journal of Mental Health</i> , 2020, 29, 581-589. | 1.9 | 0 |
| 3 | Suicide Coverage in the Digital Press Media: Adherence to World Health Organization Guidelines and Effectiveness of Different Interventions Aimed at Media Professionals. <i>Health Communication</i> , 2020, 35, 1623-1632. | 3.1 | 12 |
| 4 | Painful insight vs. usable insight in schizophrenia. Do they have different influences on suicidal behavior?. <i>Schizophrenia Research</i> , 2020, 220, 147-154. | 2.0 | 4 |
| 5 | Testing the hypothesis of subtypes of nonadherence in schizophrenia and schizoaffective disorder: A prospective study. <i>World Journal of Psychiatry</i> , 2020, 10, 260-271. | 2.7 | 3 |
| 6 | Effectiveness and cost-effectiveness of a multicomponent intervention to improve medication adherence in people with depressive disorders - MAPDep: a study protocol for a cluster randomized controlled trial. <i>Patient Preference and Adherence</i> , 2019, Volume 13, 309-319. | 1.8 | 6 |
| 7 | Suicidal risk, hopelessness and depression in patients with schizophrenia and internalized stigma. <i>Actas Espanolas De Psiquiatria</i> , 2018, 46, 33-41. | 0.1 | 7 |
| 8 | Should full adherence be a necessary goal in schizophrenia? Full versus non-full adherence to antipsychotic treatment. <i>Comprehensive Psychiatry</i> , 2014, 55, 33-39. | 3.1 | 12 |
| 9 | Influence of antipsychotic treatment type and regimen on the functionality of patients with schizophrenia. <i>Nordic Journal of Psychiatry</i> , 2014, 68, 180-188. | 1.3 | 4 |
| 10 | Antipsychotic treatment dosing profile in patients with schizophrenia evaluated with electronic monitoring (MEMSÂ®). <i>Schizophrenia Research</i> , 2013, 146, 196-200. | 2.0 | 17 |
| 11 | Creencias sobre la depresi3n y sus tratamientos: variables asociadas e influencia de las creencias en la adherencia. <i>Revista De Psiquiatr3a Y Salud Mental</i> , 2013, 6, 86-92. | 1.8 | 15 |
| 12 | Beliefs about illness and their relationship with hopelessness, depression, insight and suicide attempts in schizophrenia. <i>Psychiatria Danubina</i> , 2013, 25, 49-54. | 0.4 | 16 |
| 13 | Hopelessness and suicidal risk in bipolar disorder. A study in clinically nonsyndromal patients. <i>Comprehensive Psychiatry</i> , 2012, 53, 1103-1109. | 3.1 | 33 |
| 14 | Qualitative content of auditory hallucinations and suicidal behavior in schizophrenia. <i>Schizophrenia Research</i> , 2012, 134, 298-299. | 2.0 | 4 |
| 15 | Medication adherence in schizophrenia. <i>World Journal of Psychiatry</i> , 2012, 2, 74. | 2.7 | 136 |
| 16 | Suicidal behavior in schizophrenia and its relationship to the quality of psychotic symptoms and insight - a case report. <i>Psychiatria Danubina</i> , 2012, 24, 97-9. | 0.4 | 2 |
| 17 | Evaluation of noncompliance in schizophrenia patients using electronic monitoring (MEMSÂ®) and its relationship to sociodemographic, clinical and psychopathological variables. <i>Schizophrenia Research</i> , 2009, 107, 213-217. | 2.0 | 99 |
| 18 | Evaluation of compliance with mems in schizophrenic patients and its relationship with clinical and psychopathological variables. Preliminary data. <i>European Psychiatry</i> , 2007, 22, S101. | 0.2 | 0 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Are there subtypes of suicidal schizophrenia? A prospective study. <i>Schizophrenia Research</i> , 2006, 86, 215-220. | 2.0 | 19 |
| 20 | A psychopathological study of a group of schizophrenic patients after attempting suicide. Are there two different clinical subtypes?. <i>European Psychiatry</i> , 2003, 18, 190-192. | 0.2 | 14 |