

Su Chi Lim

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4005387/publications.pdf>

Version: 2024-02-01

135
papers

3,839
citations

159585

30
h-index

155660

55
g-index

136
all docs

136
docs citations

136
times ranked

6211
citing authors

#	ARTICLE	IF	CITATIONS
1	Risk of Incident Heart Failure in Individuals With Early-Onset Type 2 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2022, 107, e178-e187.	3.6	7
2	MODY5 Hepatocyte Nuclear Factor 1A (HNF1A)-Associated Nephropathy: experience from a regional monogenic diabetes referral centre in Singapore. <i>Journal of Investigative Medicine High Impact Case Reports</i> , 2022, 10, 232470962110656.	0.6	2
3	A real-world study on SGLT2 inhibitors and diabetic kidney disease progression. <i>CKJ: Clinical Kidney Journal</i> , 2022, 15, 1403-1414.	2.9	7
4	Insulin Allergy to Detemir Followed by Rapid Onset of Diabetic Ketoacidosis: A Case Report and Literature Review. <i>Frontiers in Endocrinology</i> , 2022, 13, 844040.	3.5	3
5	Amino acid profile of skeletal muscle loss in type 2 diabetes: Results from a 7-year longitudinal study in asians. <i>Diabetes Research and Clinical Practice</i> , 2022, 186, 109803.	2.8	8
6	Genetic Risk Score for Plasma Uric Acid Levels Is Associated With Early Rapid Kidney Function Decline in Type 2 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2022, 107, e2792-e2800.	3.6	3
7	Ethnic-Specific Type 2 Diabetes Risk Factor PAX4 R192H Is Associated with Attention-Specific Cognitive Impairment in Chinese with Type 2 Diabetes. <i>Journal of Alzheimer's Disease</i> , 2022, , 1-9.	2.6	0
8	Roux-en-Y gastric bypass versus best medical treatment for type 2 diabetes mellitus in adults with body mass index between 27 and 32 kg/m ² : A 5-year randomized control trial. <i>Diabetes Research and Clinical Practice</i> , 2022, , 109900.	2.8	3
9	Association of early-onset Type 2 diabetes with cognitive impairment is partially mediated by increased pulse pressure. <i>Journal of Diabetes and Its Complications</i> , 2022, 36, 108209.	2.3	1
10	A head-to-head comparison between Guardian Connect and FreeStyle Libre systems and an evaluation of user acceptability of sensors in patients with type 1 diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2022, 38, .	4.0	9
11	Clinical variable-based cluster analysis identifies novel subgroups with a distinct genetic signature, lipidomic pattern and cardio-renal risks in Asian patients with recent-onset type 2 diabetes. <i>Diabetologia</i> , 2022, 65, 2146-2156.	6.3	7
12	Decline in skeletal muscle mass is associated with cognitive decline in type 2 diabetes mellitus. <i>Journal of Diabetes and Its Complications</i> , 2022, , 108258.	2.3	3
13	Higher ratio of extracellular water to total body water was associated with reduced cognitive function in type 2 diabetes. <i>Journal of Diabetes</i> , 2021, 13, 222-231.	1.8	10
14	Letter to the Editor: "Prevention of Adrenal Crisis: Cortisol Responses to Major Stress Compared to Stress Dose Hydrocortisone Delivery". <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021, 106, e395-e396.	3.6	1
15	Association of leukocyte telomere length with obesity-related traits in Asian children with early-onset obesity. <i>Pediatric Obesity</i> , 2021, 16, e12771.	2.8	9
16	Association between depressive symptoms and pulse wave velocity is mediated by increased adiposity in older adults with type 2 diabetes. <i>Journal of Psychiatry and Neuroscience</i> , 2021, 46, E176-E183.	2.4	1
17	Association of leukocyte telomere length with chronic kidney disease in East Asians with type 2 diabetes: a Mendelian randomization study. <i>CKJ: Clinical Kidney Journal</i> , 2021, 14, 2371-2376.	2.9	12
18	Low muscle mass is associated with progression of chronic kidney disease and albuminuria " An 8-year longitudinal study in Asians with Type 2 Diabetes. <i>Diabetes Research and Clinical Practice</i> , 2021, 174, 108777.	2.8	8

#	ARTICLE	IF	CITATIONS
19	Association of Genetic Variants for Plasma LRG1 With Rapid Decline in Kidney Function in Patients With Type 2 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2021, 106, 2384-2394.	3.6	9
20	Association of overhydration and serum pigment epithelium-derived factor with CKD progression in diabetic kidney disease: A prospective cohort study. <i>Diabetes Research and Clinical Practice</i> , 2021, 174, 108754.	2.8	3
21	Impact of COVID-19 and partial lockdown on access to care, self-management and psychological well-being among people with diabetes: A cross-sectional study. <i>International Journal of Clinical Practice</i> , 2021, 75, e14319.	1.7	14
22	Low frequency variants associated with leukocyte telomere length in the Singapore Chinese population. <i>Communications Biology</i> , 2021, 4, 519.	4.4	15
23	Decreased GLUT2 and glucose uptake contribute to insulin secretion defects in MODY3/HNF1A hiPSC-derived mutant β^2 cells. <i>Nature Communications</i> , 2021, 12, 3133.	12.8	36
24	Aging-induced isoDGR-modified fibronectin activates monocytic and endothelial cells to promote atherosclerosis. <i>Atherosclerosis</i> , 2021, 324, 58-68.	0.8	10
25	Glucokinase activating mutation causing hypoglycaemia diagnosed late in adult who fasts for Ramadhan. <i>Endocrinology, Diabetes and Metabolism Case Reports</i> , 2021, 2021, .	0.5	0
26	Higher extracellular water to total body water ratio was associated with chronic kidney disease progression in type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2021, 35, 107930.	2.3	11
27	Transformation of Electronic Health Records and Questionnaire Data to OMOP CDM: A Feasibility Study Using SG_T2DM Dataset. <i>Applied Clinical Informatics</i> , 2021, 12, 757-767.	1.7	13
28	Association of Plasma Leucine-Rich Alpha-2 Glycoprotein 1 (LRG1) with All-Cause and Cause-Specific Mortality in Individuals with Type 2 Diabetes. <i>Clinical Chemistry</i> , 2021, 67, 1640-1649.	3.2	4
29	Associations of young onset age and genetic risk of beta cell dysfunction with glycaemic progression in individuals with type 2 diabetes. <i>Diabetes and Metabolism</i> , 2021, 47, 101238.	2.9	6
30	Association of Plasma Leucine-Rich α -2 Glycoprotein 1, a Modulator of Transforming Growth Factor- β^2 Signaling Pathway, With Incident Heart Failure in Individuals With Type 2 Diabetes. <i>Diabetes Care</i> , 2021, 44, 571-577.	8.6	11
31	Glucose Awareness to Motivate and Enable Solutions (GAMES) in diabetes mellitus using flash glucose monitoring: A clinical programme. <i>Diabetic Medicine</i> , 2021, , e14733.	2.3	3
32	The Kynurenine Pathway in Acute Kidney Injury and Chronic Kidney Disease. <i>American Journal of Nephrology</i> , 2021, 52, 771-787.	3.1	27
33	Risk of progressive chronic kidney disease in individuals with early-onset type 2 diabetes: a prospective cohort study. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, 115-121.	0.7	15
34	Association of Urine Haptoglobin With Risk of All-Cause and Cause-Specific Mortality in Individuals With Type 2 Diabetes: A Transethnic Collaborative Work. <i>Diabetes Care</i> , 2020, 43, 625-633.	8.6	6
35	Response to multiple glucose-lowering agents in a sib-pair with a novel HNF1 α (MODY3) variant. <i>European Journal of Human Genetics</i> , 2020, 28, 518-520.	2.8	5
36	The prevalence of sarcopenic obesity and its association with cognitive performance in type 2 diabetes in Singapore. <i>Clinical Nutrition</i> , 2020, 39, 2274-2281.	5.0	35

#	ARTICLE	IF	CITATIONS
37	Clinical experience from a regional monogenic diabetes referral centre in Singapore. <i>Diabetes Research and Clinical Practice</i> , 2020, 168, 108390.	2.8	3
38	Correlation of Telomere Length in Adipose Tissue and Leukocytes and its Association with Postsurgical Weight Loss. <i>Obesity</i> , 2020, 28, 2424-2430.	3.0	4
39	Aortic pulse wave velocity, central pulse pressure, augmentation index and chronic kidney disease progression in individuals with type 2 diabetes: a 3-year prospective study. <i>BMC Nephrology</i> , 2020, 21, 359.	1.8	12
40	Long-Term Observation of a Man With Severe Obesity and Undiagnosed Monogenic Diabetes Serendipitously Treated With Metabolic Surgery. <i>Journal of Investigative Medicine High Impact Case Reports</i> , 2020, 8, 232470962097487.	0.6	0
41	Association between Lower Extremity Skeletal Muscle Mass and Impaired Cognitive Function in Type 2 Diabetes. <i>Scientific Reports</i> , 2020, 10, 2956.	3.3	22
42	Plasma osteoprotegerin as a biomarker of poor glycaemic control that predicts progression of albuminuria in type 2 diabetes mellitus: A 3-year longitudinal cohort study. <i>Diabetes Research and Clinical Practice</i> , 2020, 161, 107992.	2.8	5
43	Role of endothelium-independent vasodilation in the relationship between visceral adiposity and reduced cognitive performance in older adults with type 2 diabetes. <i>Microcirculation</i> , 2020, 27, e12609.	1.8	1
44	Zinc- α -2-glycoprotein is associated with non-albuminuric chronic kidney disease progression in type 2 diabetes: a retrospective study with 4-year follow-up. <i>Diabetic Medicine</i> , 2020, 37, 1919-1926.	2.3	3
45	Metabolic Surgery Diabetes Remission (MDR) Score: a New Preoperative Scoring System for Predicting Type 2 Diabetes Remission at 1 Year After Metabolic Surgery in the Singapore Multi-ethnic Asian Setting. <i>Obesity Surgery</i> , 2020, 30, 3387-3393.	2.1	8
46	Association of traditional and novel measures of central obesity with cognitive performance in older multi-ethnic Asians with type 2 diabetes. <i>Clinical Obesity</i> , 2020, 10, e12352.	2.0	9
47	Endothelin-1 predicts incident diabetic peripheral neuropathy in Type 2 Diabetes: a cohort study. <i>European Journal of Endocrinology</i> , 2020, 182, 429-438.	3.7	9
48	Medical Costs Associated with Severity of Chronic Kidney Disease in Type 2 Diabetes Mellitus in Singapore. <i>Annals of the Academy of Medicine, Singapore</i> , 2020, 49, 731-741.	0.4	10
49	Association between vascular measures and cognitive function in type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2020, 34, 107724.	2.3	1
50	Medical costs associated with chronic kidney disease progression in an Asian population with type 2 diabetes mellitus. <i>Nephrology</i> , 2019, 24, 534-541.	1.6	15
51	Brown Adipose Tissue: Multimodality Evaluation by PET, MRI, Infrared Thermography, and Whole-Body Calorimetry (TACTICAL). <i>Obesity</i> , 2019, 27, 1434-1442.	3.0	40
52	Leucine-rich α -2-glycoprotein predicts proliferative diabetic retinopathy in type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2019, 33, 651-656.	2.3	13
53	Large-Scale Whole-Genome Sequencing of Three Diverse Asian Populations in Singapore. <i>Cell</i> , 2019, 179, 736-749.e15.	28.9	126
54	Relationship Between Fasting Plasma Glucagon Level and Renal Function—A Cross-Sectional Study in Individuals With Type 2 Diabetes. <i>Journal of the Endocrine Society</i> , 2019, 3, 273-283.	0.2	8

#	ARTICLE	IF	CITATIONS
55	Arterial Stiffness Modulates the Association of Resting Heart Rate With Rapid Renal Function Decline in Individuals With Type 2 Diabetes Mellitus. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2019, 39, 2437-2444.	2.4	6
56	Association between gain in adiposity and diabetic kidney disease worsening in type 2 diabetes is mediated by deteriorating glycaemic control: A 3-year follow-up analysis. <i>Diabetes Research and Clinical Practice</i> , 2019, 157, 107812.	2.8	2
57	Loci for human leukocyte telomere length in the Singaporean Chinese population and trans-ethnic genetic studies. <i>Nature Communications</i> , 2019, 10, 2491.	12.8	64
58	Association of haptoglobin phenotype with incident acute myocardial infarction in Chinese patients with type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2019, 18, 65.	6.8	14
59	Cause-Specific Mortality in Multiethnic South East Asians With Type 2 Diabetes Mellitus. <i>Asia-Pacific Journal of Public Health</i> , 2019, 31, 306-314.	1.0	5
60	Association of central arterial stiffness with the presence and severity of diabetic retinopathy in Asians with type 2 diabetes. <i>Diabetes and Vascular Disease Research</i> , 2019, 16, 498-505.	2.0	15
61	Precision medicine for a man presented with diabetes at 2-month old. <i>European Journal of Human Genetics</i> , 2019, 27, 989-993.	2.8	2
62	Gain in adiposity over 3 years is associated with progressive renal decline in multi-ethnic South-East Asians with type 2 diabetes. <i>Journal of Diabetes</i> , 2019, 11, 316-325.	1.8	15
63	PAX4 R192H is associated with younger onset of Type 2 diabetes in East Asians in Singapore. <i>Journal of Diabetes and Its Complications</i> , 2019, 33, 53-58.	2.3	7
64	Impact of haemoglobin A1c trajectories on chronic kidney disease progression in type 2 diabetes. <i>Nephrology</i> , 2019, 24, 1026-1032.	1.6	15
65	Ethnic disparities in relationships of obesity indices with telomere length in Asians with type 2 diabetes. <i>Journal of Diabetes</i> , 2019, 11, 386-393.	1.8	13
66	Genetic Polymorphisms and Cytokine Profile of Different Ethnicities in Septic Shock Patients, and their Association with Mortality. <i>Indian Journal of Critical Care Medicine</i> , 2019, 23, 135-138.	0.9	4
67	Discovery and validation of serum creatinine variability as novel biomarker for predicting onset of albuminuria in Type 2 diabetes mellitus. <i>Diabetes Research and Clinical Practice</i> , 2018, 138, 8-15.	2.8	9
68	Microvascular endothelial function is an independent predictor for albuminuria progression among Asians with type 2 diabetes—A prospective cohort study. <i>Microcirculation</i> , 2018, 25, e12453.	1.8	0
69	Excess visceral adiposity is associated with diabetic retinopathy in a multiethnic Asian cohort with longstanding type 2 diabetes. <i>Endocrine Research</i> , 2018, 43, 186-194.	1.2	30
70	Sex modulates the association of fibroblast growth factor 21 with end-stage renal disease in Asian people with Type 2 diabetes: a 6.3-year prospective cohort study. <i>Diabetic Medicine</i> , 2018, 35, 880-886.	2.3	8
71	Capsinoids activate brown adipose tissue (BAT) with increased energy expenditure associated with subthreshold 18-fluorine fluorodeoxyglucose uptake in BAT-positive humans confirmed by positron emission tomography scan. <i>American Journal of Clinical Nutrition</i> , 2018, 107, 62-70.	4.7	42
72	Spectrum of mutations in index patients with familial hypercholesterolemia in Singapore: Single center study. <i>Atherosclerosis</i> , 2018, 269, 106-116.	0.8	22

#	ARTICLE	IF	CITATIONS
73	Short Leukocyte Telomere Length Predicts Albuminuria Progression in Individuals With Type 2 Diabetes. <i>Kidney International Reports</i> , 2018, 3, 592-601.	0.8	21
74	A Genome-Wide Association Study of Diabetic Kidney Disease in Subjects With Type 2 Diabetes. <i>Diabetes</i> , 2018, 67, 1414-1427.	0.6	136
75	Genetic markers for urine haptoglobin is associated with decline in renal function in type 2 diabetes in East Asians. <i>Scientific Reports</i> , 2018, 8, 5109.	3.3	15
76	Efficacy of self-monitoring of blood glucose versus retrospective continuous glucose monitoring in improving glycaemic control in diabetic kidney disease patients. <i>Nephrology</i> , 2018, 23, 264-268.	1.6	16
77	Long-term outcomes of patients with type 2 diabetes attending a multidisciplinary diabetes kidney disease clinic. <i>Journal of Diabetes</i> , 2018, 10, 572-580.	1.8	9
78	Urine Tricarboxylic Acid Cycle Metabolites Predict Progressive Chronic Kidney Disease in Type 2 Diabetes. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018, 103, 4357-4364.	3.6	48
79	Long-term prospective observation suggests that glomerular hyperfiltration is associated with rapid decline in renal filtration function: A multiethnic study. <i>Diabetes and Vascular Disease Research</i> , 2018, 15, 417-423.	2.0	20
80	The role of triglyceride glucose index in development of Type 2 diabetes mellitus. <i>Diabetes Research and Clinical Practice</i> , 2018, 143, 43-49.	2.8	91
81	Low-Dose Levothyroxine Reduces Intrahepatic Lipid Content in Patients With Type 2 Diabetes Mellitus and NAFLD. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2018, 103, 2698-2706.	3.6	70
82	The association of serum creatinine and estimated glomerular filtration rate variability with diabetic retinopathy in Asians with type 2 diabetes: A nested case-control study. <i>Diabetes and Vascular Disease Research</i> , 2018, 15, 548-558.	2.0	16
83	SP428IMPACT OF HBA1C TRAJECTORIES ON CHRONIC KIDNEY DISEASE PROGRESSION IN TYPE 2 DIABETES. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, i492-i493.	0.7	0
84	Association of circulating proinflammatory marker, leucine-rich alpha ₂ -glycoprotein (LRG1), following metabolic/bariatric surgery. <i>Diabetes/Metabolism Research and Reviews</i> , 2018, 34, e3029.	4.0	25
85	Association of plasma osteopontin with diabetic retinopathy in Asians with type 2 diabetes. <i>Molecular Vision</i> , 2018, 24, 165-173.	1.1	9
86	Genetic variants in the receptor for advanced glycation end products (<i>RAGE</i>) gene were associated with circulating soluble RAGE level but not with renal function among Asians with type 2 diabetes: a genome-wide association study. <i>Nephrology Dialysis Transplantation</i> , 2017, 32, gfw263.	0.7	18
87	Association of the anti-angiogenic factor secreted protein and rich in cysteine (SPARC) with vascular complications among Chinese type 2 diabetic patients in Singapore. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 1222-1227.	2.3	2
88	Profiling of Plasma Metabolites Suggests Altered Mitochondrial Fuel Usage and Remodeling of Sphingolipid Metabolism in Individuals With Type 2 Diabetes and Kidney Disease. <i>Kidney International Reports</i> , 2017, 2, 470-480.	0.8	68
89	HbA1c variability in type 2 diabetes is associated with the occurrence of new-onset albuminuria within three years. <i>Diabetes Research and Clinical Practice</i> , 2017, 128, 32-39.	2.8	14
90	Baseline predictors of aortic stiffness progression among multi-ethnic Asians with type 2 diabetes. <i>Atherosclerosis</i> , 2017, 260, 102-109.	0.8	11

#	ARTICLE	IF	CITATIONS
91	Arterial stiffness is an independent predictor for albuminuria progression among Asians with type 2 diabetes—A prospective cohort study. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 933-938.	2.3	20
92	Association of apolipoprotein-CIII (apoC-III), endothelium-dependent vasodilation and peripheral neuropathy in a multi-ethnic population with type 2 diabetes. <i>Metabolism: Clinical and Experimental</i> , 2017, 72, 75-82.	3.4	9
93	Glutamic acid decarboxylase and islet antigen 2 antibody profiles in people with adult-onset diabetes mellitus: a comparison between mixed ethnic populations in Singapore and Germany. <i>Diabetic Medicine</i> , 2017, 34, 1145-1153.	2.3	11
94	Traffic-derived particulate matter exposure and histone H3 modification: A repeated measures study. <i>Environmental Research</i> , 2017, 153, 112-119.	7.5	52
95	Effect of long-term glycemic variability on estimated glomerular filtration rate decline among patients with type 2 diabetes mellitus: insights from the diabetic nephropathy cohort in Singapore. <i>Journal of Diabetes</i> , 2017, 9, 908-919.	1.8	21
96	Plasma Leucine-Rich Î±-2-Glycoprotein 1 Predicts Rapid eGFR Decline and Albuminuria Progression in Type 2 Diabetes Mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2017, 102, 3683-3691.	3.6	43
97	Development and validation of a predictive model for Chronic Kidney Disease progression in Type 2 Diabetes Mellitus based on a 13-year study in Singapore. <i>Diabetes Research and Clinical Practice</i> , 2017, 123, 49-54.	2.8	36
98	Direct medical cost associated with diabetic retinopathy severity in type 2 diabetes in Singapore. <i>PLoS ONE</i> , 2017, 12, e0180949.	2.5	20
99	Ethnic disparities in risk of cardiovascular disease, end-stage renal disease and all-cause mortality: a prospective study among Asian people with Type 2 diabetes. <i>Diabetic Medicine</i> , 2016, 33, 332-339.	2.3	35
100	Profile of the Paraoxonase 1 (PON1) Gene 192Q/R Polymorphism and Clinical Associations among Older Singaporean Chinese with Alzheimer's and Mixed Dementia. <i>Dementia and Geriatric Cognitive Disorders Extra</i> , 2016, 6, 43-54.	1.3	6
101	Urinary Haptoglobin Predicts Rapid Renal Function Decline in Asians With Type 2 Diabetes and Early Kidney Disease. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2016, 101, 3794-3802.	3.6	19
102	A preliminary study to evaluate the strategy of combining clinical criteria and next generation sequencing (NGS) for the identification of monogenic diabetes among multi-ethnic Asians. <i>Diabetes Research and Clinical Practice</i> , 2016, 119, 13-22.	2.8	37
103	Long-term diabetes outcomes in multi-ethnic Asians living in Singapore. <i>Diabetes Research and Clinical Practice</i> , 2016, 111, 83-92.	2.8	11
104	Onset and progression of kidney disease in type 2 diabetes among multi-ethnic Asian population. <i>Journal of Diabetes and Its Complications</i> , 2016, 30, 1248-1254.	2.3	12
105	Central arterial stiffness is associated with systemic inflammation among Asians with type 2 diabetes. <i>Diabetes and Vascular Disease Research</i> , 2016, 13, 303-306.	2.0	3
106	Circulating and visceral adipose miR-100 is down-regulated in patients with obesity and Type 2 diabetes. <i>Molecular and Cellular Endocrinology</i> , 2016, 427, 112-123.	3.2	47
107	Loss of Fas apoptosis inhibitory molecule leads to spontaneous obesity and hepatosteatosis. <i>Cell Death and Disease</i> , 2016, 7, e2091-e2091.	6.3	22
108	MicroRNAs as biomarkers of hepatotoxicity in a randomized placebo-controlled study of simvastatin and ubiquinol supplementation. <i>Experimental Biology and Medicine</i> , 2016, 241, 317-330.	2.4	20

#	ARTICLE	IF	CITATIONS
109	Health Promotion Board "Ministry of Health Clinical Practice Guidelines: Obesity. Singapore Medical Journal, 2015, 57, 292-300.	0.6	49
110	Vascular cell adhesion molecule-1, but not intercellular adhesion molecule-1, is associated with diabetic kidney disease in Asians with type 2 diabetes. Journal of Diabetes and Its Complications, 2015, 29, 707-712.	2.3	38
111	Evaluation of body adiposity index as a predictor of aortic stiffness in multi-ethnic Asian population with type 2 diabetes. Diabetes and Vascular Disease Research, 2015, 12, 111-118.	2.0	16
112	Ethnic disparity in central arterial stiffness and its determinants among Asians with type 2 diabetes. Atherosclerosis, 2015, 242, 22-28.	0.8	18
113	The role of fibroblast growth factor 21 in diabetes and its complications: A review from clinical perspective. Diabetes Research and Clinical Practice, 2015, 108, 382-389.	2.8	45
114	Elevation of a Novel Angiogenic Factor, Leucine-Rich- α 2-Glycoprotein (LRG1), Is Associated With Arterial Stiffness, Endothelial Dysfunction, and Peripheral Arterial Disease in Patients With Type 2 Diabetes. Journal of Clinical Endocrinology and Metabolism, 2015, 100, 1586-1593.	3.6	70
115	Prednisone for community-acquired pneumonia: not yet time. Lancet, The, 2015, 386, 431.	13.7	5
116	Chronic kidney disease, cardiovascular disease and mortality: A prospective cohort study in a multi-ethnic Asian population. European Journal of Preventive Cardiology, 2015, 22, 1018-1026.	1.8	56
117	Elevated Serum Leptin, Adiponectin and Leptin to Adiponectin Ratio Is Associated with Chronic Kidney Disease in Asian Adults. PLoS ONE, 2015, 10, e0122009.	2.5	48
118	Prevalence of Chronic Kidney Disease in Adults with Type 2 Diabetes Mellitus. Annals of the Academy of Medicine, Singapore, 2015, 44, 164-71.	0.4	19
119	Hepatotoxicity of anti-tuberculosis chemotherapy in patients with liver cirrhosis. International Journal of Tuberculosis and Lung Disease, 2014, 18, 347-351.	1.2	13
120	Elevated circulating alpha-klotho by angiotensin II receptor blocker losartan is associated with reduction of albuminuria in type 2 diabetic patients. JRAAS - Journal of the Renin-Angiotensin-Aldosterone System, 2014, 15, 487-490.	1.7	34
121	High normal albuminuria is independently associated with aortic stiffness in patients with Type 2 diabetes. Diabetic Medicine, 2014, 31, 1199-1204.	2.3	34
122	Association of plasma soluble α -klotho with pro-endothelin-1 in patients with type 2 diabetes. Atherosclerosis, 2014, 233, 415-418.	0.8	26
123	Relationship between circulating irisin, renal function and body composition in type 2 diabetes. Journal of Diabetes and Its Complications, 2014, 28, 208-213.	2.3	85
124	Obesity is a determinant of arterial stiffness independent of traditional risk factors in Asians with young-onset type 2 diabetes. Atherosclerosis, 2014, 236, 286-291.	0.8	19
125	Adipocytokine zinc α 2 glycoprotein (ZAG) as a novel urinary biomarker for normoalbuminuric diabetic nephropathy. Diabetic Medicine, 2012, 29, 945-949.	2.3	25
126	Microarray analysis of multiple candidate genes and associated plasma proteins for nephropathy secondary to type 2 diabetes among Chinese individuals. Diabetologia, 2009, 52, 1343-1351.	6.3	20

#	ARTICLE	IF	CITATIONS
127	Prevalence and Risk Factors for Diabetic Retinopathy. <i>Ophthalmology</i> , 2008, 115, 1869-1875.	5.2	354
128	The effect of coenzyme Q10 on microcirculatory endothelial function of subjects with type 2 diabetes mellitus. <i>Atherosclerosis</i> , 2008, 196, 966-969.	0.8	39
129	The Relationship Between Adrenomedullin, Metabolic Factors, and Vascular Function in Individuals With Type 2 Diabetes. <i>Diabetes Care</i> , 2007, 30, 1513-1519.	8.6	41
130	Angiotensin receptor antagonist vs. angiotensin-converting enzyme inhibitor in Asian subjects with type 2 diabetes and albuminuria ? a randomized crossover study. <i>Diabetes, Obesity and Metabolism</i> , 2007, 9, 477-482.	4.4	7
131	Relationship between common functional polymorphisms of the p22phox gene (âˆˆ930A > G and +242C >) Tj ETQq1 1 0.784314 rgBT (C 1037-1041.	2.3	22
132	Oxidative burden in prediabetic and diabetic individuals: evidence from plasma coenzyme Q10. <i>Diabetic Medicine</i> , 2006, 23, 1344-1349.	2.3	56
133	The Effect of Hormonal Replacement Therapy on the Vascular Reactivity and Endothelial Function of Healthy Individuals and Individuals with Type 2 Diabetes1. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1999, 84, 4159-4164.	3.6	51
134	Soluble intercellular adhesion molecule, vascular cell adhesion molecule, and impaired microvascular reactivity are early markers of vasculopathy in type 2 diabetic individuals without microalbuminuria.. <i>Diabetes Care</i> , 1999, 22, 1865-1870.	8.6	122
135	Microvascular and macrovascular reactivity is reduced in subjects at risk for type 2 diabetes.. <i>Diabetes</i> , 1999, 48, 1856-1862.	0.6	635