

# Bishnu Bashyal

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4013101/publications.pdf>

Version: 2024-02-01

10  
papers

133  
citations

1937685

4  
h-index

1474206

9  
g-index

10  
all docs

10  
docs citations

10  
times ranked

226  
citing authors

#	ARTICLE	IF	CITATIONS
1	Improved methodology for the isolation of false smut pathogen <i>Ustilaginoidea virens</i> of rice. <i>Indian Phytopathology</i> , 2021, 74, 249-252.	1.2	5
2	Identification and expression analysis of pathogenicity-related genes of <i>Rhizoctonia solani</i> anastomosis groups infecting rice. <i>3 Biotech</i> , 2021, 11, 394.	2.2	2
3	Identification of rice seed infection routes of <i>Fusarium fujikuroi</i> inciting bakanae disease of rice. <i>Journal of Plant Pathology</i> , 2020, 102, 113-121.	1.2	19
4	Molecular detection of <i>Fusarium graminearum</i> causing head blight of wheat by loop mediated isothermal amplification (LAMP) assay. <i>Indian Phytopathology</i> , 2020, 73, 667-672.	1.2	4
5	Understanding the secondary metabolite production of <i>Gibberella fujikuroi</i> species complex in genomic era. <i>Indian Phytopathology</i> , 2019, 72, 607-617.	1.2	4
6	Histopathological Studies of <i>Rhizoctonia solani</i> Infection Process in Different Cultivars of Mungbean [ <i>Vigna radiata</i> (L.) Wilczek]. <i>The National Academy of Sciences, India</i> , 2018, 41, 269-273.	1.3	3
7	Occurrence, identification and pathogenicity of <i>Fusarium</i> species associated with bakanae disease of basmati rice in India. <i>European Journal of Plant Pathology</i> , 2016, 144, 457-466.	1.7	37
8	Emergence of Aggressive Population in the <i>Bipolaris sorokiniana</i> of Barley ( <i>Hordeum vulgare</i> L.) Through Anastomosis. <i>Proceedings of the National Academy of Sciences India Section B - Biological Sciences</i> , 2015, 85, 935-941.	1.0	2
9	Pathogenicity, Ecology and Genetic Diversity of the <i>Fusarium</i> spp. Associated with an Emerging Bakanae Disease of Rice ( <i>Oryza sativa</i> L.) in India. , 2014, , 307-314.		17
10	Association of melanin content with conidiogenesis in <i>Bipolaris Sorokiniana</i> of barley ( <i>Hordeum</i> ) Tj ETQq0 0 0 rgBT/Overlock 10 Tf 50 3	3.6	40