

Hyung Jik Kim

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4013674/publications.pdf>

Version: 2024-02-01

42
papers

678
citations

759233

12
h-index

580821

25
g-index

42
all docs

42
docs citations

42
times ranked

1148
citing authors

#	ARTICLE	IF	CITATIONS
1	Prevalence of and factors associated with sarcopenia in elderly patients with end-stage renal disease. <i>Clinical Nutrition</i> , 2014, 33, 64-68.	5.0	212
2	Impact of sarcopenia on long-term mortality and cardiovascular events in patients undergoing hemodialysis. <i>Korean Journal of Internal Medicine</i> , 2019, 34, 599-607.	1.7	79
3	Could Solutions Low in Glucose Degradation Products Preserve Residual Renal Function in Incident Peritoneal Dialysis Patients? A 1-Year Multicenter Prospective Randomized Controlled Trial (Balnet) <i>Tj ETQq1 1 0.7843 14 rgB44Overlo</i>	1.4	14
4	Increased Neutrophil Extracellular Trap Formation in Uremia Is Associated with Chronic Inflammation and Prevalent Coronary Artery Disease. <i>Journal of Immunology Research</i> , 2017, 2017, 1-10.	2.2	28
5	Acute kidney injury can predict in-hospital and long-term mortality in elderly patients undergoing hip fracture surgery. <i>PLoS ONE</i> , 2017, 12, e0176259.	2.5	28
6	Obesity-related decrease in intraoperative blood flow is associated with maturation failure of radiocephalic arteriovenous fistula. <i>Journal of Vascular Surgery</i> , 2015, 62, 1010-1017.e1.	1.1	23
7	Fat-to-Lean Mass Ratio Can Predict Cardiac Events and All-Cause Mortality in Patients Undergoing Hemodialysis. <i>Annals of Nutrition and Metabolism</i> , 2018, 73, 241-249.	1.9	21
8	Prognostic role of circulating neutrophil extracellular traps levels for long-term mortality in new end-stage renal disease patients. <i>Clinical Immunology</i> , 2020, 210, 108263.	3.2	21
9	Excessive Weight Gain during the First Year of Peritoneal Dialysis Is Associated with Inflammation, Diabetes Mellitus, and a Rapid Decrease in Residual Renal Function. <i>PLoS ONE</i> , 2015, 10, e0139033.	2.5	16
10	The relationship between autophagy, increased neutrophil extracellular traps formation and endothelial dysfunction in chronic kidney disease. <i>Clinical Immunology</i> , 2018, 197, 189-197.	3.2	16
11	Late stage 3 chronic kidney disease is an independent risk factor for sarcopenia, but not proteinuria. <i>Scientific Reports</i> , 2021, 11, 18472.	3.3	16
12	Incidence of malignancy and related mortality after kidney transplantation: a nationwide, population-based cohort study in Korea. <i>Scientific Reports</i> , 2020, 10, 21398.	3.3	14
13	Effects of Excessive Body Fat Accumulation on Long-Term Outcomes during Peritoneal Dialysis. <i>Peritoneal Dialysis International</i> , 2019, 39, 268-275.	2.3	13
14	Outcomes of vascular access in hemodialysis patients: Analysis based on the Korean National Health Insurance Database from 2008 to 2016. <i>Kidney Research and Clinical Practice</i> , 2019, 38, 391-398.	2.2	13
15	A Low Baseline Glomerular Filtration Rate Predicts Poor Clinical Outcome at 3 Months after Acute		

#	ARTICLE	IF	CITATIONS
19	Four-Year Changes in Visceral Fat Mass and the Risk of Developing Proteinuria in the General Population. <i>PLoS ONE</i> , 2015, 10, e0131119.	2.5	10
20	Serum cystatin C to creatinine ratio is associated with sarcopenia in non-dialysis-dependent chronic kidney disease. <i>Kidney Research and Clinical Practice</i> , 2022, 41, 580-590.	2.2	10
21	Increased body fat rather than body weight has harmful effects on 4-year changes of renal function in the general elderly population with a normal or mildly impaired renal function. <i>Clinical Interventions in Aging</i> , 2014, 9, 1277.	2.9	9
22	Impact of rapid ultrafiltration rate on changes in the echocardiographic left atrial volume index in patients undergoing haemodialysis: a longitudinal observational study. <i>BMJ Open</i> , 2017, 7, e013990.	1.9	8
23	Proton-pump inhibitors for prevention of upper gastrointestinal bleeding in patients undergoing dialysis. <i>World Journal of Gastroenterology</i> , 2015, 21, 4919.	3.3	8
24	B-type natriuretic peptide predicts an ischemic etiology of acute heart failure in patients with stage 4-5 chronic kidney disease. <i>Clinical Biochemistry</i> , 2014, 47, 344-348.	1.9	7
25	Vitamin D: a possible modifying factor linking obesity to vascular calcification in hemodialysis patients. <i>Nutrition and Metabolism</i> , 2017, 14, 27.	3.0	7
26	Comparison of tolvaptan treatment between patients with the SIADH and congestive heart failure: a single-center experience. <i>Korean Journal of Internal Medicine</i> , 2018, 33, 561-567.	1.7	7
27	Anatomical variants of upper arm veins on preoperative mapping venography for hemodialysis access in Korean adults. <i>Journal of Vascular Access</i> , 2019, 20, 270-275.	0.9	7
28	The relationship between plasma leptin and nutritional status in chronic hemodialysis patients. <i>Journal of Korean Medical Science</i> , 1999, 14, 546.	2.5	6
29	Comparison of dynamic arterial and venous pressure between metal needles and plastic cannulas in incident hemodialysis patients with arteriovenous graft. <i>Journal of Vascular Access</i> , 2021, 22, 42-47.	0.9	4
30	Long-term prognosis of end-stage renal disease patients with normal myocardial perfusion as determined by single photon emission computed tomography. <i>Korean Journal of Internal Medicine</i> , 2018, 33, 148-156.	1.7	3
31	Serum alkaline phosphatase and \hat{I}^3 -glutamyl transferase in acute pyelonephritis. <i>Kidney Research and Clinical Practice</i> , 2019, 38, 205-211.	2.2	2
32	Custom-made Doppler ultrasound flow simulator for dialysis access using continuous renal replacement therapy machine. <i>Journal of Vascular Access</i> , 2019, 20, 701-705.	0.9	1
33	Neutrophil extracellular traps and heparin-induced antibodies contribute to vascular access thrombosis in hemodialysis patients. <i>Kidney Research and Clinical Practice</i> , 2021, 40, 712-723.	2.2	1
34	SP483 EXCESS WEIGHT GAIN DURING THE FIRST YEAR OF PERITONEAL DIALYSIS IS A RISK FACTOR FOR A RAPID DECREASE IN RESIDUAL RENAL FUNCTION. <i>Nephrology Dialysis Transplantation</i> , 2015, 30, iii539-iii539.	0.7	0
35	SP475 RAPID ULTRAFILTRATION RATE IS ASSOCIATED WITH THE INCREASE OF ECHOCARDIOGRAPHIC LEFT ATRIAL VOLUME INDEX AMONG HEMODIALYSIS PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2016, 31, i251-i252.	0.7	0
36	SP622 LOW VITAMIN D AS A MODIFYING FACTOR IN THE RELATIONSHIP BETWEEN OBESITY AND VASCULAR CALCIFICATION IN HEMODIALYSIS PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2016, 31, i302-i303.	0.7	0

#	ARTICLE	IF	CITATIONS
37	FP505 THE SIGNIFICANCE OF REPEATED FLUID STATUS MEASUREMENTS BY BIOIMPEDANCE ANALYSIS FOR PREDICTING PROGNOSIS OF PERITONEAL DIALYSIS PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2018, 33, i207-i207.	0.7	0
38	SP407 Serum protein carbonyl levels are associated with sarcopenia and long-term outcomes in hemodialysis patients. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.7	0
39	FP728 LOW LEAN TISSUE AND HIGH FAT TISSUE INDEX ARE ASSOCIATED WITH DECREASED QUALITY OF LIFE IN DIALYSIS PATIENTS. <i>Nephrology Dialysis Transplantation</i> , 2019, 34, .	0.7	0
40	P1580 LONGITUDINAL CHANGES IN BODY COMPOSITION ARE ASSOCIATED WITH ALL-CAUSE MORTALITY IN PATIENTS ON PERITONEAL DIALYSIS. <i>Nephrology Dialysis Transplantation</i> , 2020, 35, .	0.7	0
41	Role of peribrachial fat as a key determinant of brachial artery dilatation for successful arteriovenous fistula maturation in hemodialysis patients. <i>Scientific Reports</i> , 2020, 10, 3841.	3.3	0
42	A Case of Hemocholecystitis in a Patient with Microscopic Polyangiitis. <i>Korean Journal of Medicine</i> , 2012, 83, 127.	0.3	0