

Mark A Lemmon

List of Publications by Citations

Source: <https://exaly.com/author-pdf/4013850/mark-a-lemmon-publications-by-citations.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

156
papers

22,607
citations

76
h-index

150
g-index

163
ext. papers

24,672
ext. citations

14.5
avg, IF

7.21
L-index

#	Paper	IF	Citations
156	Cell signaling by receptor tyrosine kinases. <i>Cell</i> , 2010 , 141, 1117-34	56.2	2994
155	Membrane recognition by phospholipid-binding domains. <i>Nature Reviews Molecular Cell Biology</i> , 2008 , 9, 99-111	48.7	1130
154	An open-and-shut case? Recent insights into the activation of EGF/ErbB receptors. <i>Molecular Cell</i> , 2003 , 12, 541-52	17.6	693
153	Heparin-induced oligomerization of FGF molecules is responsible for FGF receptor dimerization, activation, and cell proliferation. <i>Cell</i> , 1994 , 79, 1015-24	56.2	605
152	Signal-dependent membrane targeting by pleckstrin homology (PH) domains. <i>Biochemical Journal</i> , 2000 , 350, 1-18	3.8	588
151	EGF activates its receptor by removing interactions that autoinhibit ectodomain dimerization. <i>Molecular Cell</i> , 2003 , 11, 507-17	17.6	565
150	Structure of the high affinity complex of inositol trisphosphate with a phospholipase C pleckstrin homology domain. <i>Cell</i> , 1995 , 83, 1037-46	56.2	557
149	Specific and high-affinity binding of inositol phosphates to an isolated pleckstrin homology domain. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995 , 92, 10472-6	11.5	492
148	Sequence specificity in the dimerization of transmembrane alpha-helices. <i>Biochemistry</i> , 1992 , 31, 12719-25	3.5	482
147	PH domains: diverse sequences with a common fold recruit signaling molecules to the cell surface. <i>Cell</i> , 1996 , 85, 621-4	56.2	459
146	Phosphoinositide recognition domains. <i>Traffic</i> , 2003 , 4, 201-13	5.7	457
145	Activation of phospholipase C gamma by PI 3-kinase-induced PH domain-mediated membrane targeting. <i>EMBO Journal</i> , 1998 , 17, 414-22	13	448
144	Regulation of growth factor activation by proteoglycans: what is the role of the low affinity receptors?. <i>Cell</i> , 1995 , 83, 357-60	56.2	442
143	Regulation of signal transduction and signal diversity by receptor oligomerization. <i>Trends in Biochemical Sciences</i> , 1994 , 19, 459-63	10.3	397
142	Specificity and promiscuity in phosphoinositide binding by pleckstrin homology domains. <i>Journal of Biological Chemistry</i> , 1998 , 273, 30497-508	5.4	362
141	ErbB3/HER3 intracellular domain is competent to bind ATP and catalyze autophosphorylation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2010 , 107, 7692-7	11.5	343
140	Structural basis for discrimination of 3-phosphoinositides by pleckstrin homology domains. <i>Molecular Cell</i> , 2000 , 6, 373-84	17.6	310

139	Identification and analysis of PH domain-containing targets of phosphatidylinositol 3-kinase using a novel in vivo assay in yeast. <i>EMBO Journal</i> , 1998 , 17, 5374-87	13	307
138	Genome-wide analysis of membrane targeting by <i>S. cerevisiae</i> pleckstrin homology domains. <i>Molecular Cell</i> , 2004 , 13, 677-88	17.6	280
137	A dimerization motif for transmembrane alpha-helices. <i>Nature Structural Biology</i> , 1994 , 1, 157-63		278
136	Two EGF molecules contribute additively to stabilization of the EGFR dimer. <i>EMBO Journal</i> , 1997 , 16, 281-94	13	271
135	Svp1p defines a family of phosphatidylinositol 3,5-bisphosphate effectors. <i>EMBO Journal</i> , 2004 , 23, 1922-33	13	269
134	Crystal structure at 2.2 Å resolution of the pleckstrin homology domain from human dynamin. <i>Cell</i> , 1994 , 79, 199-209	56.2	260
133	The single transmembrane domains of ErbB receptors self-associate in cell membranes. <i>Journal of Biological Chemistry</i> , 2002 , 277, 4704-12	5.4	253
132	Measurement of the binding of tyrosyl phosphopeptides to SH2 domains: a reappraisal. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995 , 92, 3199-203	11.5	251
131	The EGFR family: not so prototypical receptor tyrosine kinases. <i>Cold Spring Harbor Perspectives in Biology</i> , 2014 , 6, a020768	10.2	246
130	ALK mutations confer differential oncogenic activation and sensitivity to ALK inhibition therapy in neuroblastoma. <i>Cancer Cell</i> , 2014 , 26, 682-94	24.3	236
129	Identification of the binding site for acidic phospholipids on the pH domain of dynamin: implications for stimulation of GTPase activity. <i>Journal of Molecular Biology</i> , 1996 , 255, 14-21	6.5	231
128	The juxtamembrane region of the EGF receptor functions as an activation domain. <i>Molecular Cell</i> , 2009 , 34, 641-51	17.6	229
127	Signal-dependent membrane targeting by pleckstrin homology (PH) domains. <i>Biochemical Journal</i> , 2000 , 350, 1	3.8	224
126	Phosphatidylinositol-4,5-bisphosphate is required for endocytic coated vesicle formation. <i>Current Biology</i> , 1998 , 8, 1399-402	6.3	218
125	Epidermal growth factor receptor dimerization and activation require ligand-induced conformational changes in the dimer interface. <i>Molecular and Cellular Biology</i> , 2005 , 25, 7734-42	4.8	216
124	Pleckstrin homology domains and the cytoskeleton. <i>FEBS Letters</i> , 2002 , 513, 71-6	3.8	196
123	Phosphatidylinositol 3,5-bisphosphate: metabolism and cellular functions. <i>Trends in Biochemical Sciences</i> , 2006 , 31, 52-63	10.3	191
122	SH2 and PTB domains in tyrosine kinase signaling. <i>Science Signaling</i> , 2003 , 2003, RE12	8.8	189

121	Pleckstrin homology (PH) domains and phosphoinositides. <i>Biochemical Society Symposia</i> , 2007 , 74, 81-93		176
120	Differential inhibitor sensitivity of anaplastic lymphoma kinase variants found in neuroblastoma. <i>Science Translational Medicine</i> , 2011 , 3, 108ra114	17.5	175
119	Functional selectivity of EGF family peptide growth factors: implications for cancer. <i>Pharmacology & Therapeutics</i> , 2009 , 122, 1-8	13.9	171
118	Specificity and promiscuity in membrane helix interactions. <i>Quarterly Reviews of Biophysics</i> , 1994 , 27, 157-218	7	168
117	The glycoporphin A transmembrane domain dimer: sequence-specific propensity for a right-handed supercoil of helices. <i>Biochemistry</i> , 1992 , 31, 12726-32	3.2	166
116	Pleckstrin homology (PH) domains and phosphoinositides. <i>Biochemical Society Symposia</i> , 2007 , 81-93		163
115	The pleckstrin homology domains of dynamin isoforms require oligomerization for high affinity phosphoinositide binding. <i>Journal of Biological Chemistry</i> , 1998 , 273, 27725-33	5.4	161
114	Identification of the Rac-GEF P-Rex1 as an essential mediator of ErbB signaling in breast cancer. <i>Molecular Cell</i> , 2010 , 40, 877-92	17.6	160
113	Ligand-induced ErbB receptor dimerization. <i>Experimental Cell Research</i> , 2009 , 315, 638-48	4.2	160
112	All phox homology (PX) domains from <i>Saccharomyces cerevisiae</i> specifically recognize phosphatidylinositol 3-phosphate. <i>Journal of Biological Chemistry</i> , 2001 , 276, 44179-84	5.4	158
111	EGFR Ligands Differentially Stabilize Receptor Dimers to Specify Signaling Kinetics. <i>Cell</i> , 2017 , 171, 683-695	65.1	147
110	Structural basis for negative cooperativity in growth factor binding to an EGF receptor. <i>Cell</i> , 2010 , 142, 568-79	56.2	139
109	Erlotinib binds both inactive and active conformations of the EGFR tyrosine kinase domain. <i>Biochemical Journal</i> , 2012 , 448, 417-23	3.8	137
108	Conditional peripheral membrane proteins: facing up to limited specificity. <i>Structure</i> , 2012 , 20, 15-27	5.2	132
107	Mechanism of activation and inhibition of the HER4/ErbB4 kinase. <i>Structure</i> , 2008 , 16, 460-7	5.2	132
106	Kinase associated-1 domains drive MARK/PAR1 kinases to membrane targets by binding acidic phospholipids. <i>Cell</i> , 2010 , 143, 966-77	56.2	126
105	Structures of Eklkotho reveal a Qip codeQlike mechanism for endocrine FGF signalling. <i>Nature</i> , 2018 , 553, 501-505	50.4	118
104	Argos inhibits epidermal growth factor receptor signalling by ligand sequestration. <i>Nature</i> , 2004 , 430, 1040-4	50.4	114

103	ErbB3/HER3 does not homodimerize upon neuregulin binding at the cell surface. <i>FEBS Letters</i> , 2004 , 569, 332-6	3.8	112
102	The ALK/ROS1 Inhibitor PF-06463922 Overcomes Primary Resistance to Crizotinib in ALK-Driven Neuroblastoma. <i>Cancer Discovery</i> , 2016 , 6, 96-107	24.4	111
101	Determining selectivity of phosphoinositide-binding domains. <i>Methods</i> , 2006 , 39, 122-33	4.6	109
100	Dominant-negative inhibition of receptor-mediated endocytosis by a dynamin-1 mutant with a defective pleckstrin homology domain. <i>Current Biology</i> , 1999 , 9, 261-4	6.3	106
99	Extracellular domains drive homo- but not hetero-dimerization of erbB receptors. <i>EMBO Journal</i> , 2000 , 19, 4632-43	13	103
98	Palmitoylation of the EGFR ligand Spitz by Rasp increases Spitz activity by restricting its diffusion. <i>Developmental Cell</i> , 2006 , 10, 167-76	10.2	99
97	Normalization of nomenclature for peptide motifs as ligands of modular protein domains. <i>FEBS Letters</i> , 2002 , 513, 141-4	3.8	99
96	A structure-based model for ligand binding and dimerization of EGF receptors. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004 , 101, 929-34	11.5	98
95	Inhibition of nuclear import and cell-cycle progression by mutated forms of the dynamin-like GTPase MxB. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004 , 101, 8957-62	11.5	91
94	Molecular determinants in pleckstrin homology domains that allow specific recognition of phosphoinositides. <i>Biochemical Society Transactions</i> , 2001 , 29, 377-84	5.1	91
93	Mechanism for activation of mutated epidermal growth factor receptors in lung cancer. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, E3595-604	11.5	89
92	Dynamin GTPase regulation is altered by PH domain mutations found in centronuclear myopathy patients. <i>EMBO Journal</i> , 2010 , 29, 3054-67	13	88
91	Solution structure of pleckstrin homology domain of dynamin by heteronuclear NMR spectroscopy. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1995 , 92, 816-20	11.5	88
90	The tethered configuration of the EGF receptor extracellular domain exerts only a limited control of receptor function. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004 , 101, 923-8	11.5	86
89	Live cell imaging with protein domains capable of recognizing phosphatidylinositol 4,5-bisphosphate; a comparative study. <i>BMC Cell Biology</i> , 2009 , 10, 67		84
88	Ligand-induced structural transitions in ErbB receptor extracellular domains. <i>Structure</i> , 2007 , 15, 942-54	5.2	83
87	Kit receptor dimerization is driven by bivalent binding of stem cell factor. <i>Journal of Biological Chemistry</i> , 1997 , 272, 6311-7	5.4	82
86	N-terminal domains elicit formation of functional Pmel17 amyloid fibrils. <i>Journal of Biological Chemistry</i> , 2009 , 284, 35543-55	5.4	79

85	On the nature of low- and high-affinity EGF receptors on living cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006 , 103, 5735-40	11.5	78
84	Quantitative analysis of the effect of phosphoinositide interactions on the function of Dbl family proteins. <i>Journal of Biological Chemistry</i> , 2001 , 276, 45868-75	5.4	78
83	Role of Inn1 and its interactions with Hof1 and Cyk3 in promoting cleavage furrow and septum formation in <i>S. cerevisiae</i> . <i>Journal of Cell Biology</i> , 2009 , 185, 995-1012	7.3	77
82	Essential role for Rac in heregulin beta1 mitogenic signaling: a mechanism that involves epidermal growth factor receptor and is independent of ErbB4. <i>Molecular and Cellular Biology</i> , 2006 , 26, 831-42	4.8	77
81	Nuclear signaling by receptor tyrosine kinases: the first robin of spring. <i>Cell</i> , 2006 , 127, 45-8	56.2	77
80	The Dark Side of Cell Signaling: Positive Roles for Negative Regulators. <i>Cell</i> , 2016 , 164, 1172-1184	56.2	72
79	High-affinity binding of a FYVE domain to phosphatidylinositol 3-phosphate requires intact phospholipid but not FYVE domain oligomerization. <i>Biochemistry</i> , 2001 , 40, 8581-7	3.2	72
78	EGF-independent activation of cell-surface EGF receptors harboring mutations found in gefitinib-sensitive lung cancer. <i>Oncogene</i> , 2007 , 26, 1567-76	9.2	71
77	Dimerization of the p185neu transmembrane domain is necessary but not sufficient for transformation. <i>Oncogene</i> , 1997 , 14, 687-96	9.2	70
76	Specific role for the PH domain of dynamin-1 in the regulation of rapid endocytosis in adrenal chromaffin cells. <i>EMBO Journal</i> , 1997 , 16, 1565-74	13	70
75	TIPE3 is the transfer protein of lipid second messengers that promote cancer. <i>Cancer Cell</i> , 2014 , 26, 465-473	24.3	69
74	Ala-insertion scanning mutagenesis of the glycophorin A transmembrane helix: a rapid way to map helix-helix interactions in integral membrane proteins. <i>Protein Science</i> , 1996 , 5, 1339-41	6.3	68
73	ErbB2 resembles an autoinhibited invertebrate epidermal growth factor receptor. <i>Nature</i> , 2009 , 461, 287-91	50.4	67
72	Assessing the range of kinase autoinhibition mechanisms in the insulin receptor family. <i>Biochemical Journal</i> , 2012 , 448, 213-20	3.8	65
71	Thermodynamic studies of tyrosyl-phosphopeptide binding to the SH2 domain of p56lck. <i>Biochemistry</i> , 1994 , 33, 5070-6	3.2	64
70	The role of the pleckstrin homology domain in membrane targeting and activation of phospholipase Cbeta(1). <i>Journal of Biological Chemistry</i> , 2000 , 275, 14873-81	5.4	56
69	Receptor tyrosine kinases with intracellular pseudokinase domains. <i>Biochemical Society Transactions</i> , 2013 , 41, 1029-36	5.1	54
68	Complex relationship between ligand binding and dimerization in the epidermal growth factor receptor. <i>Cell Reports</i> , 2014 , 9, 1306-17	10.6	53

67	Loss of phosphatidylinositol 3-phosphate binding by the C-terminal Tiam-1 pleckstrin homology domain prevents in vivo Rac1 activation without affecting membrane targeting. <i>Journal of Biological Chemistry</i> , 2003 , 278, 11457-64	5.4	53
66	Crystal structure of fibroblast growth factor 9 reveals regions implicated in dimerization and autoinhibition. <i>Journal of Biological Chemistry</i> , 2001 , 276, 4322-9	5.4	53
65	A possible effector role for the pleckstrin homology (PH) domain of dynamin. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2009 , 106, 13359-64	11.5	51
64	The p21-activated protein kinase-related kinase Cla4 is a coincidence detector of signaling by Cdc42 and phosphatidylinositol 4-phosphate. <i>Journal of Biological Chemistry</i> , 2004 , 279, 17101-10	5.4	50
63	Helix-helix interactions inside lipid bilayers. <i>Current Opinion in Structural Biology</i> , 1992 , 2, 511-518	8.1	50
62	Antibody targeting of anaplastic lymphoma kinase induces cytotoxicity of human neuroblastoma. <i>Oncogene</i> , 2012 , 31, 4859-67	9.2	47
61	Membrane activity of the phospholipase C-delta1 pleckstrin homology (PH) domain. <i>Biochemical Journal</i> , 2005 , 389, 435-41	3.8	47
60	Specificity and promiscuity in membrane helix interactions. <i>FEBS Letters</i> , 1994 , 346, 17-20	3.8	47
59	Mutations in or near the transmembrane domain alter PMEL amyloid formation from functional to pathogenic. <i>PLoS Genetics</i> , 2011 , 7, e1002286	6	41
58	Structural basis for EGFR ligand sequestration by Argos. <i>Nature</i> , 2008 , 453, 1271-5	50.4	41
57	Loss of pleckstrin defines a novel pathway for PKC-mediated exocytosis. <i>Blood</i> , 2009 , 113, 3577-84	2.2	40
56	Phosphoinositide binding by the pleckstrin homology domains of Ipl and Tih1. <i>Journal of Biological Chemistry</i> , 2002 , 277, 49935-44	5.4	39
55	Putting together structures of epidermal growth factor receptors. <i>Current Opinion in Structural Biology</i> , 2014 , 29, 95-101	8.1	37
54	The Dbs PH domain contributes independently to membrane targeting and regulation of guanine nucleotide-exchange activity. <i>Biochemical Journal</i> , 2006 , 400, 563-72	3.8	37
53	Computational analysis of EGFR inhibition by Argos. <i>Developmental Biology</i> , 2005 , 284, 523-35	3.1	33
52	Comparison of <i>Saccharomyces cerevisiae</i> F-BAR domain structures reveals a conserved inositol phosphate binding site. <i>Structure</i> , 2015 , 23, 352-63	5.2	32
51	Specificity of the myotubularin family of phosphatidylinositol-3-phosphatase is determined by the PH/GRAM domain. <i>Journal of Biological Chemistry</i> , 2006 , 281, 31762-9	5.4	32
50	Pleckstrin homology domains: two halves make a hole?. <i>Cell</i> , 2005 , 120, 574-6	56.2	32

49	Structural Insights into Pseudokinase Domains of Receptor Tyrosine Kinases. <i>Molecular Cell</i> , 2020 , 79, 390-405.e7	17.6	30
48	Thermodynamic studies of SHC phosphotyrosine interaction domain recognition of the NPXpY motif. <i>Journal of Biological Chemistry</i> , 1996 , 271, 4770-5	5.4	28
47	Drug Sensitivity and Allele Specificity of First-Line Osimertinib Resistance Mutations. <i>Cancer Research</i> , 2020 , 80, 2017-2030	10.1	27
46	Ligand regulation of a constitutively dimeric EGF receptor. <i>Nature Communications</i> , 2015 , 6, 7380	17.4	26
45	Molecular dynamics analysis of conserved hydrophobic and hydrophilic bond-interaction networks in ErbB family kinases. <i>Biochemical Journal</i> , 2011 , 436, 241-51	3.8	26
44	The EGFR Exon 19 Mutant L747-A750>P Exhibits Distinct Sensitivity to Tyrosine Kinase Inhibitors in Lung Adenocarcinoma. <i>Clinical Cancer Research</i> , 2019 , 25, 6382-6391	12.9	21
43	Overcoming resistance to HER2 inhibitors through state-specific kinase binding. <i>Nature Chemical Biology</i> , 2016 , 12, 923-930	11.7	21
42	Dimerization of Tie2 mediated by its membrane-proximal FNIII domains. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017 , 114, 4382-4387	11.5	20
41	Insulin and epidermal growth factor receptor family members share parallel activation mechanisms. <i>Protein Science</i> , 2020 , 29, 1331-1344	6.3	18
40	Molecular determinants of KA1 domain-mediated autoinhibition and phospholipid activation of MARK1 kinase. <i>Biochemical Journal</i> , 2017 , 474, 385-398	3.8	16
39	Non-acylated Wnts Can Promote Signaling. <i>Cell Reports</i> , 2019 , 26, 875-883.e5	10.6	16
38	The EGF receptor family as therapeutic targets in breast cancer. <i>Breast Disease</i> , 2003 , 18, 33-43	1.6	15
37	Genome-wide analysis of signaling domain function. <i>Current Opinion in Chemical Biology</i> , 2003 , 7, 103-9	9.7	14
36	Kinetics of receptor tyrosine kinase activation define ERK signaling dynamics. <i>Science Signaling</i> , 2020 , 13,	8.8	14
35	A new twist in the transmembrane signaling tool-kit. <i>Cell</i> , 2007 , 130, 213-5	56.2	11
34	Analysis of phosphoinositide binding by pleckstrin homology domain from dynamin. <i>Methods in Enzymology</i> , 2001 , 329, 457-68	1.7	9
33	Structural Basis for MARK1 Kinase Autoinhibition by Its KA1 Domain. <i>Structure</i> , 2018 , 26, 1137-1143.e3	5.2	9
32	EGFR mutations cause a lethal syndrome of epithelial dysfunction with progeroid features. <i>Molecular Genetics & Genomic Medicine</i> , 2015 , 3, 452-8	2.3	8

31	Biochemistry. KSR plays CRAF-ty. <i>Science</i> , 2011 , 332, 1043-4	33.3	8
30	Computational algorithms for in silico profiling of activating mutations in cancer. <i>Cellular and Molecular Life Sciences</i> , 2019 , 76, 2663-2679	10.3	7
29	Protein kinase C regulation: C1 meets C-tail. <i>Structure</i> , 2011 , 19, 144-6	5.2	7
28	Specificity of the Myotubularin Family of Phosphatidylinositol-3-phosphatase Is Determined by the PH/GRAM Domain. <i>Journal of Biological Chemistry</i> , 2006 , 281, 31762-31769	5.4	7
27	Drugging the "Undruggable" MYCN Oncogenic Transcription Factor: Overcoming Previous Obstacles to Impact Childhood Cancers. <i>Cancer Research</i> , 2021 , 81, 1627-1632	10.1	7
26	Occupy EGFR. <i>Cancer Discovery</i> , 2012 , 2, 398-400	24.4	6
25	ROR and RYK extracellular region structures suggest that receptor tyrosine kinases have distinct WNT-recognition modes. <i>Cell Reports</i> , 2021 , 37, 109834	10.6	6
24	Neuregulin Signaling Is a Mechanism of Therapeutic Resistance in Head and Neck Squamous Cell Carcinoma. <i>Molecular Cancer Therapeutics</i> , 2019 , 18, 2124-2134	6.1	5
23	Structural basis for ligand reception by anaplastic lymphoma kinase. <i>Nature</i> , 2021 , 600, 148-152	50.4	5
22	Regulation of Kinase Activity in the <i>Caenorhabditis elegans</i> EGF Receptor, LET-23. <i>Structure</i> , 2018 , 26, 270-281.e4	5.2	4
21	Argos mutants define an affinity threshold for spitz inhibition in vivo. <i>Journal of Biological Chemistry</i> , 2006 , 281, 28993-9001	5.4	4
20	Glioblastoma mutations alter EGFR dimer structure to prevent ligand bias.. <i>Nature</i> , 2022 ,	50.4	4
19	Smoothing out the patches. <i>Science</i> , 2018 , 362, 26-27	33.3	3
18	Flipping ATP to AMPlify Kinase Functions. <i>Cell</i> , 2018 , 175, 641-642	56.2	3
17	Phosphatidylserine binding directly regulates TIM-3 function. <i>Biochemical Journal</i> , 2021 , 478, 3331-3349	3.8	3
16	Pleckstrin Homology (PH) Domains 2010 , 1093-1101		2
15	PH Domains 2005 , 337-363		2
14	Deletion Mutations Keep Kinase Inhibitors in the Loop. <i>Cancer Cell</i> , 2016 , 29, 423-425	24.3	2

13	Comparison of tyrosine kinase domain properties for the neurotrophin receptors TrkA and TrkB. <i>Biochemical Journal</i> , 2020 , 477, 4053-4070	3.8	1
12	Computational studies of anaplastic lymphoma kinase mutations reveal common mechanisms of oncogenic activation. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021 , 118,	11.5	1
11	Looking lively: emerging principles of pseudokinase signaling.. <i>Trends in Biochemical Sciences</i> , 2022 ,	10.3	1
10	Dynamics of protein kinases and pseudokinases by HDX-MS.. <i>Methods in Enzymology</i> , 2022 , 667, 303-338	1.7	0
9	Structural bases for specific phosphoinositide binding by PH domains. <i>Biochemical Society Transactions</i> , 1999 , 27, A73-A73	5.1	
8	Pleckstrin Homology (PH) Domains 2003 , 161-169		
7	Drug Sensitivity and Allele-specificity of First-line Osimertinib Resistance EGFR Mutations. <i>FASEB Journal</i> , 2020 , 34, 1-1	0.9	
6	Auto-inhibition of dynamin GTPase activity is regulated by PH domain interactions. <i>FASEB Journal</i> , 2009 , 23, 697.3	0.9	
5	Regulation of the epidermal growth factor receptor intracellular domain. <i>FASEB Journal</i> , 2009 , 23, 883.2	0.9	
4	Structural basis for EGFR ligand sequestration by Argos. <i>FASEB Journal</i> , 2009 , 23, 883.7	0.9	
3	ErbB2/HER2/Neu resembles an autoinhibited invertebrate EGF receptor. <i>FASEB Journal</i> , 2009 , 23, 884.3	0.9	
2	Phosphoinositide-mimicking peptide sequences are binding targets for PH domains. <i>FASEB Journal</i> , 2009 , 23, 873.7	0.9	
1	Characterization of Novel PtdIns(4,5)P2 Effector Domains. <i>FASEB Journal</i> , 2009 , 23, 873.6	0.9	