

Alfredo Falcone

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/402071/publications.pdf>

Version: 2024-02-01

395
papers

27,623
citations

10979

71
h-index

6831

155
g-index

409
all docs

409
docs citations

409
times ranked

25612
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 1 | ESMO consensus guidelines for the management of patients with metastatic colorectal cancer. <i>Annals of Oncology</i> , 2016, 27, 1386-1422. | 0.6 | 2,545 |
| 2 | Regorafenib monotherapy for previously treated metastatic colorectal cancer (CORRECT): an international, multicentre, randomised, placebo-controlled, phase 3 trial. <i>Lancet</i> , The, 2013, 381, 303-312. | 6.3 | 2,276 |
| 3 | Phase III Trial of Infusional Fluorouracil, Leucovorin, Oxaliplatin, and Irinotecan (FOLFOXIRI) Compared With Infusional Fluorouracil, Leucovorin, and Irinotecan (FOLFIRI) As First-Line Treatment for Metastatic Colorectal Cancer: The Gruppo Oncologico Nord Ovest. <i>Journal of Clinical Oncology</i> , 2007, 25, 1670-1676. | 0.8 | 1,083 |
| 4 | Randomized Trial of TAS-102 for Refractory Metastatic Colorectal Cancer. <i>New England Journal of Medicine</i> , 2015, 372, 1909-1919. | 13.9 | 1,027 |
| 5 | Pembrolizumab versus paclitaxel for previously treated, advanced gastric or gastro-oesophageal junction cancer (KEYNOTE-061): a randomised, open-label, controlled, phase 3 trial. <i>Lancet</i> , The, 2018, 392, 123-133. | 6.3 | 984 |
| 6 | Initial Therapy with FOLFOXIRI and Bevacizumab for Metastatic Colorectal Cancer. <i>New England Journal of Medicine</i> , 2014, 371, 1609-1618. | 13.9 | 845 |
| 7 | FOLFOXIRI plus bevacizumab versus FOLFIRI plus bevacizumab as first-line treatment of patients with metastatic colorectal cancer: updated overall survival and molecular subgroup analyses of the open-label, phase 3 TRIBE study. <i>Lancet Oncology</i> , The, 2015, 16, 1306-1315. | 5.1 | 835 |
| 8 | Clonal evolution and resistance to EGFR blockade in the blood of colorectal cancer patients. <i>Nature Medicine</i> , 2015, 21, 795-801. | 15.2 | 809 |
| 9 | KRAS codon 61, 146 and BRAF mutations predict resistance to cetuximab plus irinotecan in KRAS codon 12 and 13 wild-type metastatic colorectal cancer. <i>British Journal of Cancer</i> , 2009, 101, 715-721. | 2.9 | 509 |
| 10 | Neratinib after trastuzumab-based adjuvant therapy in HER2-positive breast cancer (ExteNET): 5-year analysis of a randomised, double-blind, placebo-controlled, phase 3 trial. <i>Lancet Oncology</i> , The, 2017, 18, 1688-1700. | 5.1 | 451 |
| 11 | Prognosis of patients with peritoneal metastatic colorectal cancer given systemic therapy: an analysis of individual patient data from prospective randomised trials from the Analysis and Research in Cancers of the Digestive System (ARCAD) database. <i>Lancet Oncology</i> , The, 2016, 17, 1709-1719. | 5.1 | 442 |
| 12 | MicroRNA-21 in Pancreatic Cancer: Correlation with Clinical Outcome and Pharmacologic Aspects Underlying Its Role in the Modulation of Gemcitabine Activity. <i>Cancer Research</i> , 2010, 70, 4528-4538. | 0.4 | 409 |
| 13 | PTEN Expression and KRAS Mutations on Primary Tumors and Metastases in the Prediction of Benefit From Cetuximab Plus Irinotecan for Patients With Metastatic Colorectal Cancer. <i>Journal of Clinical Oncology</i> , 2009, 27, 2622-2629. | 0.8 | 402 |
| 14 | Primary Tumor Location as a Prognostic Factor in Metastatic Colorectal Cancer. <i>Journal of the National Cancer Institute</i> , 2015, 107, . | 3.0 | 385 |
| 15 | Atezolizumab with or without cobimetinib versus regorafenib in previously treated metastatic colorectal cancer (IMblaze370): a multicentre, open-label, phase 3, randomised, controlled trial. <i>Lancet Oncology</i> , The, 2019, 20, 849-861. | 5.1 | 368 |
| 16 | Quantitative evidence for early metastatic seeding in colorectal cancer. <i>Nature Genetics</i> , 2019, 51, 1113-1122. | 9.4 | 315 |
| 17 | Analysis of circulating DNA and protein biomarkers to predict the clinical activity of regorafenib and assess prognosis in patients with metastatic colorectal cancer: a retrospective, exploratory analysis of the CORRECT trial. <i>Lancet Oncology</i> , The, 2015, 16, 937-948. | 5.1 | 286 |
| 18 | Rechallenge for Patients With KRAS and BRAF Wild-Type Metastatic Colorectal Cancer With Acquired Resistance to First-line Cetuximab and Irinotecan. <i>JAMA Oncology</i> , 2019, 5, 343. | 3.4 | 280 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 19 | Pharmacogenetic Profiling in Patients With Advanced Colorectal Cancer Treated With First-Line FOLFOX-4 Chemotherapy. <i>Journal of Clinical Oncology</i> , 2007, 25, 1247-1254. | 0.8 | 250 |
| 20 | Bevacizumab with FOLFOXIRI (irinotecan, oxaliplatin, fluorouracil, and folinate) as first-line treatment for metastatic colorectal cancer: a phase 2 trial. <i>Lancet Oncology</i> , The, 2010, 11, 845-852. | 5.1 | 234 |
| 21 | High Concordance of <i>KRAS</i> Status Between Primary Colorectal Tumors and Related Metastatic Sites: Implications for Clinical Practice. <i>Oncologist</i> , 2008, 13, 1270-1275. | 1.9 | 218 |
| 22 | Long-Term Outcome of Initially Unresectable Metastatic Colorectal Cancer Patients Treated with 5-Fluorouracil/Leucovorin, Oxaliplatin, and Irinotecan (FOLFOXIRI) Followed by Radical Surgery of Metastases. <i>Annals of Surgery</i> , 2009, 249, 420-425. | 2.1 | 213 |
| 23 | The Detection of Androgen Receptor Splice Variant 7 in Plasma-derived Exosomal RNA Strongly Predicts Resistance to Hormonal Therapy in Metastatic Prostate Cancer Patients. <i>European Urology</i> , 2017, 71, 680-687. | 0.9 | 213 |
| 24 | A Systematic Review of the Burden of Pancreatic Cancer in Europe: Real-World Impact on Survival, Quality of Life and Costs. <i>Journal of Gastrointestinal Cancer</i> , 2015, 46, 201-211. | 0.6 | 199 |
| 25 | Upfront FOLFOXIRI plus bevacizumab and reintroduction after progression versus mFOLFOX6 plus bevacizumab followed by FOLFIRI plus bevacizumab in the treatment of patients with metastatic colorectal cancer (TRIBE2): a multicentre, open-label, phase 3, randomised, controlled trial. <i>Lancet Oncology</i> , The, 2020, 21, 497-507. | 5.1 | 196 |
| 26 | Correlation of <i>CDA, ERCC1</i> , and <i>XPD</i> Polymorphisms with Response and Survival in Gemcitabine/Cisplatin-Treated Advanced Non-Small Cell Lung Cancer Patients. <i>Clinical Cancer Research</i> , 2008, 14, 1797-1803. | 3.2 | 193 |
| 27 | Ramucirumab with cisplatin and fluoropyrimidine as first-line therapy in patients with metastatic gastric or junctional adenocarcinoma (RAINFALL): a double-blind, randomised, placebo-controlled, phase 3 trial. <i>Lancet Oncology</i> , The, 2019, 20, 420-435. | 5.1 | 191 |
| 28 | PD-L1 mRNA expression in plasma-derived exosomes is associated with response to anti-PD-1 antibodies in melanoma and NSCLC. <i>British Journal of Cancer</i> , 2018, 118, 820-824. | 2.9 | 190 |
| 29 | ALK, ROS1, and NTRK Rearrangements in Metastatic Colorectal Cancer. <i>Journal of the National Cancer Institute</i> , 2017, 109, . | 3.0 | 183 |
| 30 | Cetuximab rechallenge in metastatic colorectal cancer patients: how to come away from acquired resistance?. <i>Annals of Oncology</i> , 2012, 23, 2313-2318. | 0.6 | 170 |
| 31 | Mucinous histology predicts for poor response rate and overall survival of patients with colorectal cancer and treated with first-line oxaliplatin- and/or irinotecan-based chemotherapy. <i>British Journal of Cancer</i> , 2009, 100, 881-887. | 2.9 | 164 |
| 32 | Randomized Trial of Two Induction Chemotherapy Regimens in Metastatic Colorectal Cancer: An Updated Analysis. <i>Journal of the National Cancer Institute</i> , 2011, 103, 21-30. | 3.0 | 160 |
| 33 | A randomised clinical trial of two docetaxel regimens (weekly vs 3 week) in the second-line treatment of non-small-cell lung cancer. The DISTAL 01 study. <i>British Journal of Cancer</i> , 2004, 91, 1996-2004. | 2.9 | 158 |
| 34 | Treatment with 5-Fluorouracil/Folinic Acid, Oxaliplatin, and Irinotecan Enables Surgical Resection of Metastases in Patients With Initially Unresectable Metastatic Colorectal Cancer. <i>Annals of Surgical Oncology</i> , 2006, 13, 58-65. | 0.7 | 156 |
| 35 | Early tumor shrinkage and depth of response predict long-term outcome in metastatic colorectal cancer patients treated with first-line chemotherapy plus bevacizumab: results from phase III TRIBE trial by the Gruppo Oncologico del Nord Ovest. <i>Annals of Oncology</i> , 2015, 26, 1188-1194. | 0.6 | 153 |
| 36 | Biweekly Chemotherapy With Oxaliplatin, Irinotecan, Infusional Fluorouracil, and Leucovorin: A Pilot Study in Patients With Metastatic Colorectal Cancer. <i>Journal of Clinical Oncology</i> , 2002, 20, 4006-4014. | 0.8 | 148 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 37 | EZH2 inhibition: targeting the crossroad of tumor invasion and angiogenesis. <i>Cancer and Metastasis Reviews</i> , 2012, 31, 753-761. | 2.7 | 148 |
| 38 | BRAF and RAS mutations as prognostic factors in metastatic colorectal cancer patients undergoing liver resection. <i>British Journal of Cancer</i> , 2015, 112, 1921-1928. | 2.9 | 146 |
| 39 | Individual Patient Data Meta-Analysis of FOLFOXIRI Plus Bevacizumab Versus Doublets Plus Bevacizumab as Initial Therapy of Unresectable Metastatic Colorectal Cancer. <i>Journal of Clinical Oncology</i> , 2020, 38, 3314-3324. | 0.8 | 139 |
| 40 | First-line chemotherapy for mCRC—a review and evidence-based algorithm. <i>Nature Reviews Clinical Oncology</i> , 2015, 12, 607-619. | 12.5 | 138 |
| 41 | BRAF codons 594 and 596 mutations identify a new molecular subtype of metastatic colorectal cancer at favorable prognosis. <i>Annals of Oncology</i> , 2015, 26, 2092-2097. | 0.6 | 137 |
| 42 | Cetuximab plus gemcitabine and cisplatin compared with gemcitabine and cisplatin alone in patients with advanced pancreatic cancer: a randomised, multicentre, phase II trial. <i>Lancet Oncology</i> , The, 2008, 9, 39-44. | 5.1 | 130 |
| 43 | Phase II study of cetuximab in combination with cisplatin and docetaxel in patients with untreated advanced gastric or gastro-oesophageal junction adenocarcinoma (DOCETUX study). <i>British Journal of Cancer</i> , 2009, 101, 1261-1268. | 2.9 | 130 |
| 44 | Role of <i>NRAS</i> mutations as prognostic and predictive markers in metastatic colorectal cancer. <i>International Journal of Cancer</i> , 2015, 136, 83-90. | 2.3 | 126 |
| 45 | Pharmacogenetic Profiling for Cetuximab Plus Irinotecan Therapy in Patients With Refractory Advanced Colorectal Cancer. <i>Journal of Clinical Oncology</i> , 2008, 26, 1427-1434. | 0.8 | 124 |
| 46 | Nuclear Factor- κ B Tumor Expression Predicts Response and Survival in Irinotecan-Refractory Metastatic Colorectal Cancer Treated With Cetuximab-Irinotecan Therapy. <i>Journal of Clinical Oncology</i> , 2007, 25, 3930-3935. | 0.8 | 121 |
| 47 | Body Mass Index Is Prognostic in Metastatic Colorectal Cancer: Pooled Analysis of Patients From First-Line Clinical Trials in the ARCAD Database. <i>Journal of Clinical Oncology</i> , 2016, 34, 144-150. | 0.8 | 116 |
| 48 | Hypersensitivity reactions related to oxaliplatin (OHP). <i>British Journal of Cancer</i> , 2003, 89, 477-481. | 2.9 | 113 |
| 49 | Genetic modulation of the Let-7 microRNA binding to KRAS 3' untranslated region and survival of metastatic colorectal cancer patients treated with salvage cetuximab+irinotecan. <i>Pharmacogenomics Journal</i> , 2010, 10, 458-464. | 0.9 | 109 |
| 50 | Natural history of bone metastasis in colorectal cancer: final results of a large Italian bone metastases study. <i>Annals of Oncology</i> , 2012, 23, 2072-2077. | 0.6 | 108 |
| 51 | Neutrophil-to-Lymphocyte Ratio (NLR), Platelet-to-Lymphocyte Ratio (PLR), and Outcomes with Nivolumab in Pretreated Non-Small Cell Lung Cancer (NSCLC): A Large Retrospective Multicenter Study. <i>Advances in Therapy</i> , 2020, 37, 1145-1155. | 1.3 | 102 |
| 52 | First-line treatment of metastatic colorectal cancer with irinotecan, oxaliplatin and 5-fluorouracil/leucovorin (FOLFOXIRI): results of a phase II study with a simplified biweekly schedule. <i>Annals of Oncology</i> , 2004, 15, 1766-1772. | 0.6 | 99 |
| 53 | Pharmacogenetic profiling in patients with advanced colorectal cancer treated with first-line FOLFIRI chemotherapy. <i>Pharmacogenomics Journal</i> , 2008, 8, 278-288. | 0.9 | 97 |
| 54 | Relationship between 5-fluorouracil disposition, toxicity and dihydropyrimidine dehydrogenase activity in cancer patients. <i>Annals of Oncology</i> , 2001, 12, 1301-1306. | 0.6 | 94 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Circulating endothelial cells and endothelial progenitors as predictive markers of clinical response to bevacizumab-based first-line treatment in advanced colorectal cancer patients. <i>Annals of Oncology</i> , 2010, 21, 2382-2389. | 0.6 | 94 |
| 56 | Location of Primary Tumor and Benefit From Anti-Epidermal Growth Factor Receptor Monoclonal Antibodies in Patients With <i>RAS</i> and <i>BRAF</i> Wild-Type Metastatic Colorectal Cancer. <i>Oncologist</i> , 2016, 21, 988-994. | 1.9 | 94 |
| 57 | Cyclophosphamide-methotrexate $\hat{=}$ metronomic $\hat{=}$ chemotherapy for the palliative treatment of metastatic breast cancer. A comparative pharmacoeconomic evaluation. <i>Annals of Oncology</i> , 2005, 16, 1243-1252. | 0.6 | 91 |
| 58 | Personalizing Survival Predictions in Advanced Colorectal Cancer: The ARCAD Nomogram Project. <i>Journal of the National Cancer Institute</i> , 2018, 110, 638-648. | 3.0 | 90 |
| 59 | Regorafenib for Patients with Metastatic Colorectal Cancer Who Progressed After Standard Therapy: Results of the Large, Single-Arm, Open-Label Phase IIIb CONSIGN Study. <i>Oncologist</i> , 2019, 24, 185-192. | 1.9 | 89 |
| 60 | Individual Patient Data Analysis of Progression-Free Survival Versus Overall Survival As a First-Line End Point for Metastatic Colorectal Cancer in Modern Randomized Trials: Findings From the Analysis and Research in Cancers of the Digestive System Database. <i>Journal of Clinical Oncology</i> , 2015, 33, 22-28. | 0.8 | 87 |
| 61 | Activity and Safety of Cetuximab Plus Modified FOLFOXIRI Followed by Maintenance With Cetuximab or Bevacizumab for <i>RAS</i> and <i>BRAF</i> Wild-type Metastatic Colorectal Cancer. <i>JAMA Oncology</i> , 2018, 4, 529. | 3.4 | 87 |
| 62 | FOLFOXIRI in combination with panitumumab as first-line treatment in quadruple wild-type (<i>KRAS</i> ,) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 Nord Ovest (GONO). <i>Annals of Oncology</i> , 2013, 24, 2062-2067. | 0.6 | 86 |
| 63 | Antiangiogenic and anticolorrectal cancer effects of metronomic irinotecan chemotherapy alone and in combination with semaxinib. <i>British Journal of Cancer</i> , 2008, 98, 1619-1629. | 2.9 | 85 |
| 64 | Clinical and Pharmacodynamic Evaluation of Metronomic Cyclophosphamide, Celecoxib, and Dexamethasone in Advanced Hormone-refractory Prostate Cancer. <i>Clinical Cancer Research</i> , 2009, 15, 4954-4962. | 3.2 | 85 |
| 65 | Pharmacodynamic and pharmacogenetic angiogenesis-related markers of first-line FOLFOXIRI plus bevacizumab schedule in metastatic colorectal cancer. <i>British Journal of Cancer</i> , 2011, 104, 1262-1269. | 2.9 | 85 |
| 66 | Randomized trial on adjuvant treatment with FOLFIRI followed by docetaxel and cisplatin versus 5-fluorouracil and folinic acid for radically resected gastric cancer. <i>Annals of Oncology</i> , 2014, 25, 1373-1378. | 0.6 | 84 |
| 67 | The good, the bad and the ugly: a tale of miR-101, miR-21 and miR-155 in pancreatic intraductal papillary mucinous neoplasms. <i>Annals of Oncology</i> , 2013, 24, 734-741. | 0.6 | 83 |
| 68 | Contribution of <i>KRAS</i> mutations and c.2369C > T (p.T790M) <i>EGFR</i> to acquired resistance to EGFR-TKIs in <i>EGFR</i> mutant NSCLC: a study on circulating tumor DNA. <i>Oncotarget</i> , 2017, 8, 13611-13619. | 0.8 | 81 |
| 69 | Cetuximab Rechallenge Plus Avelumab in Pretreated Patients With <i>RAS</i> Wild-type Metastatic Colorectal Cancer. <i>JAMA Oncology</i> , 2021, 7, 1529. | 3.4 | 80 |
| 70 | Prognosis of mucinous histology for patients with radically resected stage II and III colon cancer. <i>Annals of Oncology</i> , 2012, 23, 135-141. | 0.6 | 79 |
| 71 | Glycolysis gene expression analysis and selective metabolic advantage in the clinical progression of colorectal cancer. <i>Pharmacogenomics Journal</i> , 2017, 17, 258-264. | 0.9 | 79 |
| 72 | High Let-7a MicroRNA Levels in <i>KRAS</i> -Mutated Colorectal Carcinomas May Rescue Anti-EGFR Therapy Effects in Patients with Chemotherapy-Refractory Metastatic Disease. <i>Oncologist</i> , 2012, 17, 823-829. | 1.9 | 74 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Anti-HER agents in gastric cancer: from bench to bedside. <i>Nature Reviews Gastroenterology and Hepatology</i> , 2011, 8, 369-383. | 8.2 | 73 |
| 74 | Current status and perspectives in immunotherapy for metastatic melanoma. <i>Oncotarget</i> , 2018, 9, 12452-12470. | 0.8 | 73 |
| 75 | Negative hyper-selection of metastatic colorectal cancer patients for anti-EGFR monoclonal antibodies: the PRESSING case-control study. <i>Annals of Oncology</i> , 2017, 28, 3009-3014. | 0.6 | 72 |
| 76 | Overexpression of TK1 and CDK9 in plasma-derived exosomes is associated with clinical resistance to CDK4/6 inhibitors in metastatic breast cancer patients. <i>Breast Cancer Research and Treatment</i> , 2019, 178, 57-62. | 1.1 | 71 |
| 77 | Maintenance Therapy With Panitumumab Alone vs Panitumumab Plus Fluorouracil-Leucovorin in Patients With <i>RAS</i> Wild-Type Metastatic Colorectal Cancer. <i>JAMA Oncology</i> , 2019, 5, 1268. | 3.4 | 70 |
| 78 | Retrospective exploratory analysis of VEGF polymorphisms in the prediction of benefit from first-line FOLFIRI plus bevacizumab in metastatic colorectal cancer. <i>BMC Cancer</i> , 2011, 11, 247. | 1.1 | 69 |
| 79 | Multivariate prognostic factors analysis for second-line chemotherapy in advanced biliary tract cancer. <i>British Journal of Cancer</i> , 2014, 110, 2165-2169. | 2.9 | 69 |
| 80 | FOLFIRINOX and translational studies: Towards personalized therapy in pancreatic cancer. <i>World Journal of Gastroenterology</i> , 2016, 22, 6987. | 1.4 | 68 |
| 81 | Insulin-like growth factor 1 expression correlates with clinical outcome in <i>KRAS</i> wild type colorectal cancer patients treated with cetuximab and irinotecan. <i>International Journal of Cancer</i> , 2010, 127, 1941-1947. | 2.3 | 67 |
| 82 | Association of Polymorphisms in <i>AKT1</i> and <i>EGFR</i> with Clinical Outcome and Toxicity in Non-Small Cell Lung Cancer Patients Treated with Gefitinib. <i>Molecular Cancer Therapeutics</i> , 2010, 9, 581-593. | 1.9 | 67 |
| 83 | High-Throughput MicroRNA (miRNAs) Arrays Unravel the Prognostic Role of MiR-211 in Pancreatic Cancer. <i>PLoS ONE</i> , 2012, 7, e49145. | 1.1 | 67 |
| 84 | Epidermal Growth Factor Receptor (EGFR) gene copy number (GCN) correlates with clinical activity of irinotecan-cetuximab in <i>K-RAS</i> wild-type colorectal cancer: a fluorescence in situ (FISH) and chromogenic in situ hybridization (CISH) analysis. <i>BMC Cancer</i> , 2009, 9, 303. | 1.1 | 66 |
| 85 | Early changes in plasma DNA levels of mutant <i>KRAS</i> as a sensitive marker of response to chemotherapy in pancreatic cancer. <i>Scientific Reports</i> , 2017, 7, 7931. | 1.6 | 66 |
| 86 | Epidermal growth factor receptor (EGFR) gene promoter methylation and cetuximab treatment in colorectal cancer patients. <i>British Journal of Cancer</i> , 2011, 104, 1786-1790. | 2.9 | 65 |
| 87 | Prospective Validation of Candidate SNPs of VEGF/VEGFR Pathway in Metastatic Colorectal Cancer Patients Treated with First-Line FOLFIRI Plus Bevacizumab. <i>PLoS ONE</i> , 2013, 8, e66774. | 1.1 | 64 |
| 88 | A pharmacokinetic-based test to prevent severe 5-fluorouracil toxicity. <i>Clinical Pharmacology and Therapeutics</i> , 2006, 80, 384-395. | 2.3 | 63 |
| 89 | A pharmacokinetic and pharmacodynamic study on metronomic irinotecan in metastatic colorectal cancer patients. <i>British Journal of Cancer</i> , 2008, 98, 1312-1319. | 2.9 | 63 |
| 90 | Impact of <i>ABCG2</i> polymorphisms on the clinical outcome and toxicity of gefitinib in non-small-cell lung cancer patients. <i>Pharmacogenomics</i> , 2011, 12, 159-170. | 0.6 | 63 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | Clinical, pharmacokinetic and pharmacodynamic evaluations of metronomic UFT and cyclophosphamide plus celecoxib in patients with advanced refractory gastrointestinal cancers. <i>Angiogenesis</i> , 2012, 15, 275-286. | 3.7 | 61 |
| 92 | First-line anti-EGFR monoclonal antibodies in panRAS wild-type metastatic colorectal cancer: A systematic review and meta-analysis. <i>Critical Reviews in Oncology/Hematology</i> , 2015, 96, 156-166. | 2.0 | 61 |
| 93 | Interprofessional spiritual care in oncology: a literature review. <i>ESMO Open</i> , 2019, 4, e000465. | 2.0 | 61 |
| 94 | Vascular Endothelial Growth Factor Levels in Immunodepleted Plasma of Cancer Patients As a Possible Pharmacodynamic Marker for Bevacizumab Activity. <i>Journal of Clinical Oncology</i> , 2007, 25, 1816-1818. | 0.8 | 56 |
| 95 | Clinico-pathological nomogram for predicting BRAF mutational status of metastatic colorectal cancer. <i>British Journal of Cancer</i> , 2016, 114, 30-36. | 2.9 | 56 |
| 96 | Early magnesium modifications as a surrogate marker of efficacy of cetuximab-based anticancer treatment in KRAS wild-type advanced colorectal cancer patients. <i>Annals of Oncology</i> , 2011, 22, 1141-1146. | 0.6 | 54 |
| 97 | Second-line chemotherapy in advanced biliary cancer progressed to first-line platinum-gemcitabine combination: a multicenter survey and pooled analysis with published data. <i>Journal of Experimental and Clinical Cancer Research</i> , 2015, 34, 156. | 3.5 | 54 |
| 98 | Efficacy of FOLFOXIRI plus bevacizumab in liver-limited metastatic colorectal cancer: A pooled analysis of clinical studies by Gruppo Oncologico del Nord Ovest. <i>European Journal of Cancer</i> , 2017, 73, 74-84. | 1.3 | 54 |
| 99 | AtezoTRIBE: a randomised phase II study of FOLFOXIRI plus bevacizumab alone or in combination with atezolizumab as initial therapy for patients with unresectable metastatic colorectal cancer. <i>BMC Cancer</i> , 2020, 20, 683. | 1.1 | 53 |
| 100 | Sorafenib plus daily low-dose temozolomide for relapsed glioblastoma: a phase II study. <i>Anticancer Research</i> , 2013, 33, 3487-94. | 0.5 | 53 |
| 101 | 5-Fluorouracil Pharmacokinetics Predicts Disease-free Survival in Patients Administered Adjuvant Chemotherapy for Colorectal Cancer. <i>Clinical Cancer Research</i> , 2008, 14, 2749-2755. | 3.2 | 52 |
| 102 | Sequence Effect of Irinotecan and Fluorouracil Treatment on Pharmacokinetics and Toxicity in Chemotherapy-Naive Metastatic Colorectal Cancer Patients. <i>Journal of Clinical Oncology</i> , 2001, 19, 3456-3462. | 0.8 | 51 |
| 103 | Histopathologic evaluation of liver metastases from colorectal cancer in patients treated with FOLFOXIRI plus bevacizumab. <i>British Journal of Cancer</i> , 2013, 108, 2549-2556. | 2.9 | 51 |
| 104 | Baseline elevated leukocyte count in peripheral blood is associated with poor survival in patients with advanced non-small cell lung cancer: a prognostic model. <i>Journal of Cancer Research and Clinical Oncology</i> , 2008, 134, 1143-1149. | 1.2 | 49 |
| 105 | VEGF gene polymorphisms and susceptibility to colorectal cancer disease in Italian population. <i>International Journal of Colorectal Disease</i> , 2009, 24, 165-170. | 1.0 | 47 |
| 106 | Magnitude of benefit of the addition of bevacizumab to first-line chemotherapy for metastatic colorectal cancer: meta-analysis of randomized clinical trials. <i>Journal of Experimental and Clinical Cancer Research</i> , 2010, 29, 58. | 3.5 | 46 |
| 107 | FOLFOXIRI or FOLFOXIRI plus bevacizumab as first-line treatment of metastatic colorectal cancer: a propensity score-adjusted analysis from two randomized clinical trials. <i>Annals of Oncology</i> , 2016, 27, 843-849. | 0.6 | 46 |
| 108 | Trifluridine/Tipiracil (TAS-102) in Refractory Metastatic Colorectal Cancer: A Multicenter Register in the Frame of the Italian Compassionate Use Program. <i>Oncologist</i> , 2018, 23, 1178-1187. | 1.9 | 46 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|------|-----------|
| 109 | Safety and effectiveness of regorafenib in patients with metastatic colorectal cancer in routine clinical practice in the prospective, observational CORRELATE study. <i>European Journal of Cancer</i> , 2019, 123, 146-154. | 1.3 | 46 |
| 110 | Prognostic clinical factors in pretreated colorectal cancer patients receiving regorafenib: Implications for clinical management. <i>Oncotarget</i> , 2015, 6, 33982-33992. | 0.8 | 46 |
| 111 | Estimating 12-week death probability in patients with refractory metastatic colorectal cancer: the Colon Life nomogram. <i>Annals of Oncology</i> , 2017, 28, 555-561. | 0.6 | 43 |
| 112 | Human Equilibrative Nucleoside Transporter 1 (hENT1) Levels Predict Response to Gemcitabine in Patients With Biliary Tract Cancer (BTC). <i>Current Cancer Drug Targets</i> , 2011, 11, 123-129. | 0.8 | 42 |
| 113 | VEGF-A polymorphisms predict progression-free survival among advanced castration-resistant prostate cancer patients treated with metronomic cyclophosphamide. <i>British Journal of Cancer</i> , 2013, 109, 957-964. | 2.9 | 41 |
| 114 | An EZH2 polymorphism is associated with clinical outcome in metastatic colorectal cancer patients. <i>Annals of Oncology</i> , 2012, 23, 1207-1213. | 0.6 | 40 |
| 115 | Prospective validation of a lymphocyte infiltration prognostic test in stage III colon cancer patients treated with adjuvant FOLFOX. <i>European Journal of Cancer</i> , 2017, 82, 16-24. | 1.3 | 40 |
| 116 | Erlotinib after Failure of Gefitinib in Patients with Advanced Non-small Cell Lung Cancer Previously Responding to Gefitinib. <i>Journal of Thoracic Oncology</i> , 2008, 3, 912-914. | 0.5 | 39 |
| 117 | High concordance of BRAF status between primary colorectal tumours and related metastatic sites: implications for clinical practice. <i>Annals of Oncology</i> , 2010, 21, 1565. | 0.6 | 38 |
| 118 | First-line treatment with FOLFOXIRI for advanced pancreatic cancer in clinical practice: Patients' outcome and analysis of prognostic factors. <i>International Journal of Cancer</i> , 2016, 139, 938-945. | 2.3 | 38 |
| 119 | Safety and tolerability of subcutaneous trastuzumab for the adjuvant treatment of human epidermal growth factor receptor 2-positive early breast cancer: SafeHer phase III study's primary analysis of 2573 patients. <i>European Journal of Cancer</i> , 2017, 82, 237-246. | 1.3 | 38 |
| 120 | Metastatic BRAF K601E-mutated melanoma reaches complete response to MEK inhibitor trametinib administered for over 36 months. <i>Experimental Hematology and Oncology</i> , 2017, 6, 6. | 2.0 | 38 |
| 121 | Clinical Calculator for Early Mortality in Metastatic Colorectal Cancer: An Analysis of Patients From 28 Clinical Trials in the Aide et Recherche en Cancérologie Digestive Database. <i>Journal of Clinical Oncology</i> , 2017, 35, 1929-1937. | 0.8 | 37 |
| 122 | Immune Checkpoint Inhibitors in pMMR Metastatic Colorectal Cancer: A Tough Challenge. <i>Cancers</i> , 2020, 12, 2317. | 1.7 | 37 |
| 123 | A Model of a Zebrafish Avatar for Co-Clinical Trials. <i>Cancers</i> , 2020, 12, 677. | 1.7 | 36 |
| 124 | Metronomic 5-fluorouracil, oxaliplatin and irinotecan in colorectal cancer. <i>European Journal of Pharmacology</i> , 2009, 619, 8-14. | 1.7 | 35 |
| 125 | First-line therapy for mCRC – the influence of primary tumour location on the therapeutic algorithm. <i>Nature Reviews Clinical Oncology</i> , 2017, 14, 113-113. | 12.5 | 35 |
| 126 | DPYD*6 plays an important role in fluoropyrimidine toxicity in addition to DPYD*2A and c.2846A>T: a comprehensive analysis in 1254 patients. <i>Pharmacogenomics Journal</i> , 2019, 19, 556-563. | 0.9 | 35 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 127 | Analysis of HER-3, insulin growth factor-1, nuclear factor-kB and epidermal growth factor receptor gene copy number in the prediction of clinical outcome for K-RAS wild-type colorectal cancer patients receiving irinotecan+cetuximab. <i>Annals of Oncology</i> , 2012, 23, 1706-1712. | 0.6 | 34 |
| 128 | Clinical impact of anti-epidermal growth factor receptor monoclonal antibodies in first-line treatment of metastatic colorectal cancer. <i>Cancer</i> , 2012, 118, 1523-1532. | 2.0 | 34 |
| 129 | Results of a phase III randomized, double-blind, placebo-controlled, multicenter trial (CORRECT) of regorafenib plus best supportive care (BSC) versus placebo plus BSC in patients (pts) with metastatic colorectal cancer (mCRC) who have progressed after standard therapies.. <i>Journal of Clinical Oncology</i> , 2012, 30, LBA385-LBA385. | 0.8 | 34 |
| 130 | Docetaxel plus oral metronomic cyclophosphamide: A phase II study with pharmacodynamic and pharmacogenetic analyses in castration-resistant prostate cancer patients. <i>Cancer</i> , 2014, 120, 3923-3931. | 2.0 | 33 |
| 131 | MRI tumor volume reduction rate vs tumor regression grade in the pre-operative re-staging of locally advanced rectal cancer after chemo-radiotherapy. <i>European Journal of Radiology</i> , 2015, 84, 2438-2443. | 1.2 | 32 |
| 132 | Androgen receptor (AR) splice variant 7 and full-length AR expression is associated with clinical outcome: a translational study in patients with castration-resistant prostate cancer. <i>BJU International</i> , 2019, 124, 693-700. | 1.3 | 32 |
| 133 | Cetuximab plus irinotecan after irinotecan failure in elderly metastatic colorectal cancer patients: Clinical outcome according to KRAS and BRAF mutational status. <i>Critical Reviews in Oncology/Hematology</i> , 2011, 78, 243-251. | 2.0 | 31 |
| 134 | Application of the ESR iGuide clinical decision support system to the imaging pathway of patients with hepatocellular carcinoma and cholangiocarcinoma: preliminary findings. <i>Radiologia Medica</i> , 2020, 125, 531-537. | 4.7 | 31 |
| 135 | Suramin in combination with weekly epirubicin for patients with advanced hormone-refractory prostate carcinoma. , 1999, 86, 470-476. | | 30 |
| 136 | A multicenter phase II study of the combination of oxaliplatin, irinotecan and capecitabine in the first-line treatment of metastatic colorectal cancer. <i>British Journal of Cancer</i> , 2009, 100, 1720-1724. | 2.9 | 30 |
| 137 | Prognostic significance of <i>K-Ras</i> mutation rate in metastatic colorectal cancer patients. <i>Oncotarget</i> , 2015, 6, 31604-31612. | 0.8 | 30 |
| 138 | Liver-only metastatic colorectal cancer patients and thymidylate synthase polymorphisms for predicting response to 5-fluorouracil-based chemotherapy. <i>British Journal of Cancer</i> , 2008, 99, 716-721. | 2.9 | 29 |
| 139 | Could Interferon Still Play a Role in Metastatic Renal Cell Carcinoma? A Randomized Study of Two Schedules of Sorafenib Plus Interferon-Alpha 2a (RAPSODY). <i>European Urology</i> , 2013, 63, 254-261. | 0.9 | 29 |
| 140 | Variations in genes regulating tumor-associated macrophages (TAMs) to predict outcomes of bevacizumab-based treatment in patients with metastatic colorectal cancer: results from TRIBE and FIRE3 trials. <i>Annals of Oncology</i> , 2015, 26, 2450-2456. | 0.6 | 29 |
| 141 | Clinical, pharmacodynamic and pharmacokinetic results of a prospective phase II study on oral metronomic vinorelbine and dexamethasone in castration-resistant prostate cancer patients. <i>Investigational New Drugs</i> , 2016, 34, 760-770. | 1.2 | 29 |
| 142 | Serum LDH predicts benefit from bevacizumab beyond progression in metastatic colorectal cancer. <i>British Journal of Cancer</i> , 2017, 116, 318-323. | 2.9 | 29 |
| 143 | Consensus statement on essential patient characteristics in systemic treatment trials for metastatic colorectal cancer: Supported by the ARCAD Group. <i>European Journal of Cancer</i> , 2018, 100, 35-45. | 1.3 | 29 |
| 144 | Adjuvant Systemic Chemotherapy After Putative Curative Resection of Colorectal Liver and Lung Metastases. <i>Clinical Colorectal Cancer</i> , 2013, 12, 188-194. | 1.0 | 28 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 145 | Impact of genetic variations in the MAPK signaling pathway on outcome in metastatic colorectal cancer patients treated with first-line FOLFIRI and bevacizumab: data from FIRE-3 and TRIBE trials. <i>Annals of Oncology</i> , 2017, 28, 2780-2785. | 0.6 | 28 |
| 146 | TRIBE-2: a phase III, randomized, open-label, strategy trial in unresectable metastatic colorectal cancer patients by the GONO group. <i>BMC Cancer</i> , 2017, 17, 408. | 1.1 | 28 |
| 147 | Clinical Validation of a Machine-learningâ€‘derived Signature Predictive of Outcomes from First-line Oxaliplatin-based Chemotherapy in Advanced Colorectal Cancer. <i>Clinical Cancer Research</i> , 2021, 27, 1174-1183. | 3.2 | 28 |
| 148 | METRONOMIC CYCLOPHOSPHAMIDE IN ELDERLY PATIENTS WITH ADVANCED, CASTRATION-RESISTANT PROSTATE CANCER. <i>Journal of the American Geriatrics Society</i> , 2010, 58, 986-988. | 1.3 | 27 |
| 149 | Circulating endothelial cells and their apoptotic fraction are mutually independent predictive biomarkers in Bevacizumab-based treatment for advanced colorectal cancer. <i>Journal of Cancer Research and Clinical Oncology</i> , 2012, 138, 1187-1196. | 1.2 | 27 |
| 150 | EGFR ligands as pharmacodynamic biomarkers in metastatic colorectal cancer patients treated with cetuximab and irinotecan. <i>Targeted Oncology</i> , 2014, 9, 205-214. | 1.7 | 27 |
| 151 | Apatinib in Advanced Gastric Cancer: A Doubtful Step Forward. <i>Journal of Clinical Oncology</i> , 2016, 34, 3822-3823. | 0.8 | 27 |
| 152 | HER2 Overexpression as a Poor Prognostic Determinant in Resected Biliary Tract Cancer. <i>Oncologist</i> , 2020, 25, 886-893. | 1.9 | 27 |
| 153 | Homologous Recombination Deficiency Alterations in Colorectal Cancer: Clinical, Molecular, and Prognostic Implications. <i>Journal of the National Cancer Institute</i> , 2022, 114, 271-279. | 3.0 | 27 |
| 154 | Safety, efficacy and patient-reported outcomes with trifluridine/tipiracil in pretreated metastatic colorectal cancer: results of the PRECONNECT study. <i>ESMO Open</i> , 2020, 5, e000698. | 2.0 | 26 |
| 155 | Adjuvant chemotherapy in the treatment of colon cancer: randomized multicenter trial of the Italian National Intergroup of Adjuvant Chemotherapy in Colon Cancer (INTACC). <i>Annals of Oncology</i> , 2003, 14, 1365-1372. | 0.6 | 25 |
| 156 | Genes involved in pericyte-driven tumor maturation predict treatment benefit of first-line FOLFIRI plus bevacizumab in patients with metastatic colorectal cancer. <i>Pharmacogenomics Journal</i> , 2015, 15, 69-76. | 0.9 | 25 |
| 157 | Radiological imaging markers predicting clinical outcome in patients with metastatic colorectal carcinoma treated with regorafenib: post hoc analysis of the CORRECT phase III trial (RadioCORRECT) <i>Tj ETQq1 1 0z784314 rgBT /Ove</i> | | |
| 158 | Angiogenesis genotyping and clinical outcome during regorafenib treatment in metastatic colorectal cancer patients. <i>Scientific Reports</i> , 2016, 6, 25195. | 1.6 | 25 |
| 159 | Gene Polymorphisms in the CCL5/CCR5 Pathway as a Genetic Biomarker for Outcome and Handâ€‘Foot Skin Reaction in Metastatic Colorectal Cancer Patients Treated With Regorafenib. <i>Clinical Colorectal Cancer</i> , 2018, 17, e395-e414. | 1.0 | 25 |
| 160 | Phase II randomised study of maintenance treatment with bevacizumab or bevacizumab plus metronomic chemotherapy after first-line induction with FOLFOXIRI plus Bevacizumab for metastatic colorectal cancer patients: the MOMA trial. <i>European Journal of Cancer</i> , 2019, 109, 175-182. | 1.3 | 25 |
| 161 | Total neoadjuvant approach with FOLFOXIRI plus bevacizumab followed by chemoradiotherapy plus bevacizumab in locally advanced rectal cancer: the TRUST trial. <i>European Journal of Cancer</i> , 2019, 110, 32-41. | 1.3 | 25 |
| 162 | <i>DPYD</i> and <i>UGT1A1</i> genotyping to predict adverse events during first-line FOLFIRI or FOLFOXIRI plus bevacizumab in metastatic colorectal cancer. <i>Oncotarget</i> , 2018, 9, 7859-7866. | 0.8 | 25 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 163 | Improved Analysis of 5-Fluorouracil and 5,6-Dihydro-5-Fluorouracil by HPLC With Diode Array Detection for Determination of Cellular Dihydropyrimidine Dehydrogenase Activity and Pharmacokinetic Profiling. <i>Therapeutic Drug Monitoring</i> , 2005, 27, 362-368. | 1.0 | 24 |
| 164 | Bevacizumab in the pre-operative treatment of locally advanced rectal cancer: A systematic review. <i>World Journal of Gastroenterology</i> , 2014, 20, 6081. | 1.4 | 24 |
| 165 | Limited sampling model for the analysis of 5-fluorouracil pharmacokinetics in adjuvant chemotherapy for colorectal cancer. <i>Clinical Pharmacology and Therapeutics</i> , 2002, 72, 627-637. | 2.3 | 23 |
| 166 | A Phase I and Pharmacokinetic Study of Irinotecan Given as a 7-Day Continuous Infusion in Metastatic Colorectal Cancer Patients Pretreated with 5-Fluorouracil or Raltitrexed. <i>Clinical Cancer Research</i> , 2004, 10, 1657-1663. | 3.2 | 23 |
| 167 | Variations in the interleukin-1 receptor antagonist gene impact on survival of patients with advanced colorectal cancer. <i>Pharmacogenomics Journal</i> , 2009, 9, 78-84. | 0.9 | 23 |
| 168 | A Polymorphism within the Vitamin D Transporter Gene Predicts Outcome in Metastatic Colorectal Cancer Patients Treated with FOLFIRI/Bevacizumab or FOLFIRI/Cetuximab. <i>Clinical Cancer Research</i> , 2018, 24, 784-793. | 3.2 | 23 |
| 169 | The emerging role of liquid biopsy in diagnosis, prognosis and treatment monitoring of pancreatic cancer. <i>Pharmacogenomics</i> , 2019, 20, 49-68. | 0.6 | 23 |
| 170 | AXL is a predictor of poor survival and of resistance to anti-EGFR therapy in RAS wild-type metastatic colorectal cancer. <i>European Journal of Cancer</i> , 2020, 138, 1-10. | 1.3 | 23 |
| 171 | Immunogenic cell death pathway polymorphisms for predicting oxaliplatin efficacy in metastatic colorectal cancer. , 2020, 8, e001714. | | 23 |
| 172 | Use of zebrafish embryos as avatar of patients with pancreatic cancer: A new xenotransplantation model towards personalized medicine. <i>World Journal of Gastroenterology</i> , 2020, 26, 2792-2809. | 1.4 | 23 |
| 173 | Infusions of Fluorouracil and Leucovorin: Effects of the Timing and Semi-Intermittency of Drug Delivery. <i>Oncology</i> , 1999, 57, 195-201. | 0.9 | 22 |
| 174 | First-line 5-fluorouracil/folinic acid, oxaliplatin and irinotecan (FOLFOXIRI) does not impair the feasibility and the activity of second line treatments in metastatic colorectal cancer. <i>Annals of Oncology</i> , 2006, 17, 1249-1254. | 0.6 | 22 |
| 175 | Dihydropyrimidine dehydrogenase polymorphisms and fluoropyrimidine toxicity: ready for routine clinical application within personalized medicine?. <i>EPMA Journal</i> , 2010, 1, 495-502. | 3.3 | 22 |
| 176 | Multimodality treatment of locally advanced squamous cell carcinoma of the oesophagus: A comprehensive review and network meta-analysis. <i>Critical Reviews in Oncology/Hematology</i> , 2017, 114, 24-32. | 2.0 | 22 |
| 177 | Proxies of quality of life in metastatic colorectal cancer: analyses in the RECURSE trial. <i>ESMO Open</i> , 2017, 2, e000261. | 2.0 | 22 |
| 178 | Immune Checkpoint Inhibitors in Esophageal Cancers: Are We Finally Finding the Right Path in the Mist?. <i>International Journal of Molecular Sciences</i> , 2020, 21, 1658. | 1.8 | 22 |
| 179 | Impact of Cytidine Deaminase Polymorphisms on Toxicity After Gemcitabine: The Question Is Still Ongoing. <i>Journal of Clinical Oncology</i> , 2010, 28, e221-e222. | 0.8 | 21 |
| 180 | Metronomic Chemotherapy for Metastatic Prostate Cancer. <i>Drugs and Aging</i> , 2010, 27, 689-696. | 1.3 | 21 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 181 | Locally advanced gastro-oesophageal cancer: Recent therapeutic advances and research directions. <i>Cancer Treatment Reviews</i> , 2018, 69, 90-100. | 3.4 | 21 |
| 182 | Genetic interaction of P2X7 receptor and VEGFR-2 polymorphisms identifies a favorable prognostic profile in prostate cancer patients. <i>Oncotarget</i> , 2015, 6, 28743-28754. | 0.8 | 21 |
| 183 | Stereotactic Body Radiotherapy in Patients with Lung Oligometastases from Colorectal Cancer. <i>Anticancer Research</i> , 2017, 37, 315-320. | 0.5 | 21 |
| 184 | Recombinant β -2a Interferon Plus Vinblastine in the Treatment of Metastatic Renal Cell Carcinoma. <i>American Journal of Clinical Oncology: Cancer Clinical Trials</i> , 1989, 12, 43-45. | 0.6 | 20 |
| 185 | TRIPLETE: a randomised phase III study of modified FOLFOXIRI plus panitumumab versus mFOLFOX6 plus panitumumab as initial therapy for patients with unresectable RAS and BRAF wild-type metastatic colorectal cancer. <i>ESMO Open</i> , 2018, 3, e000403. | 2.0 | 20 |
| 186 | First-line trifluridine/tipiracil plus bevacizumab for unresectable metastatic colorectal cancer: SOLSTICE study design. <i>Future Oncology</i> , 2020, 16, 21-29. | 1.1 | 20 |
| 187 | Retreatment With Anti-EGFR Antibodies in Metastatic Colorectal Cancer Patients: A Multi-institutional Analysis. <i>Clinical Colorectal Cancer</i> , 2020, 19, 191-199.e6. | 1.0 | 20 |
| 188 | Intravenous azidothymidine with fluorouracil and leucovorin: a phase I-II study in previously untreated metastatic colorectal cancer patients. <i>Journal of Clinical Oncology</i> , 1996, 14, 729-736. | 0.8 | 19 |
| 189 | Second-Line Chemotherapy with a Modified Schedule of Docetaxel in Elderly Patients with Advanced-Stage Non-Small-Cell Lung Cancer. <i>Clinical Lung Cancer</i> , 2006, 7, 401-405. | 1.1 | 19 |
| 190 | Dicer and Drosha expression and response to Bevacizumab-based therapy in advanced colorectal cancer patients. <i>European Journal of Cancer</i> , 2013, 49, 1501-1508. | 1.3 | 19 |
| 191 | The role of primary tumour sidedness, EGFR gene copy number and EGFR promoter methylation in RAS/BRAF wild-type colorectal cancer patients receiving irinotecan/cetuximab. <i>British Journal of Cancer</i> , 2017, 117, 315-321. | 2.9 | 19 |
| 192 | Autophagy-related polymorphisms predict hypertension in patients with metastatic colorectal cancer treated with FOLFIRI and bevacizumab: Results from TRIBE and FIRE-3 trials. <i>European Journal of Cancer</i> , 2017, 77, 13-20. | 1.3 | 19 |
| 193 | Palliative treatment of unresectable metastatic colorectal cancer. <i>Expert Opinion on Pharmacotherapy</i> , 2010, 11, 63-77. | 0.9 | 18 |
| 194 | Phase II study of single-agent cetuximab in KRAS G13D mutant metastatic colorectal cancer. <i>Annals of Oncology</i> , 2015, 26, 2503. | 0.6 | 18 |
| 195 | Impact of a supportive care service for cancer outpatients: management and reduction of hospitalizations. Preliminary results of an integrated model of care. <i>Supportive Care in Cancer</i> , 2017, 25, 209-212. | 1.0 | 18 |
| 196 | Prognostic and Predictive Biomarkers in Patients with Metastatic Colorectal Cancer Receiving Regorafenib. <i>Molecular Cancer Therapeutics</i> , 2020, 19, 2146-2154. | 1.9 | 18 |
| 197 | Severe 5-fluorouracil toxicity associated with a marked alteration of pharmacokinetics of 5-fluorouracil and its catabolite 5-fluoro-5,6-dihydrouracil: a case report. <i>European Journal of Clinical Pharmacology</i> , 2002, 58, 593-595. | 0.8 | 17 |
| 198 | Capecitabine and Mitomycin c May be an Effective Treatment Option for Third-line Chemotherapy in Advanced Colorectal Cancer. <i>Tumori</i> , 2006, 92, 384-388. | 0.6 | 17 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 199 | Should Oncologists Be Aware in Their Clinical Practice of <i>KRAS</i> Molecular Analysis?. <i>Journal of Clinical Oncology</i> , 2011, 29, e206-e207. | 0.8 | 17 |
| 200 | Outcome of Second-Line Treatment After First-Line Chemotherapy With the GONO FOLFOXIRI Regimen. <i>Clinical Colorectal Cancer</i> , 2012, 11, 71-76. | 1.0 | 17 |
| 201 | Molecular analysis of cell-free circulating DNA for the diagnosis of somatic mutations associated with resistance to tyrosine kinase inhibitors in non-small-cell lung cancer. <i>Expert Review of Molecular Diagnostics</i> , 2014, 14, 453-468. | 1.5 | 17 |
| 202 | Differential histopathologic parameters in colorectal cancer liver metastases resected after triplets plus bevacizumab or cetuximab: a pooled analysis of five prospective trials. <i>British Journal of Cancer</i> , 2018, 118, 955-965. | 2.9 | 17 |
| 203 | Oligometastatic colorectal cancer: prognosis, role of locoregional treatments and impact of first-line chemotherapy—a pooled analysis of TRIBE and TRIBE2 studies by Gruppo Oncologico del Nord Ovest. <i>European Journal of Cancer</i> , 2020, 139, 81-89. | 1.3 | 17 |
| 204 | <i>BRAF</i> V600E Mutation in First-Line Metastatic Colorectal Cancer: An Analysis of Individual Patient Data From the ARCAD Database. <i>Journal of the National Cancer Institute</i> , 2021, 113, 1386-1395. | 3.0 | 17 |
| 205 | FOLFOXIRI plus bevacizumab (bev) versus FOLFIRI plus bev as first-line treatment of metastatic colorectal cancer (mCRC): Updated survival results of the phase III TRIBE trial by the GONO group.. <i>Journal of Clinical Oncology</i> , 2015, 33, 657-657. | 0.8 | 17 |
| 206 | Prognostic relevance of a T-type calcium channels gene signature in solid tumours: A correlation ready for clinical validation. <i>PLoS ONE</i> , 2017, 12, e0182818. | 1.1 | 17 |
| 207 | Triplet chemotherapy in combination with anti-EGFR agents for the treatment of metastatic colorectal cancer: Current evidence, advances, and future perspectives. <i>Cancer Treatment Reviews</i> , 2022, 102, 102301. | 3.4 | 17 |
| 208 | Suramin in patients with metastatic colorectal cancer pretreated with fluoropyrimidine-based chemotherapy. A phase II Study. <i>Cancer</i> , 1995, 75, 440-443. | 2.0 | 16 |
| 209 | Adjuvant sequential methotrexate + 5-fluorouracil vs 5-fluorouracil plus leucovorin in radically resected stage III and high-risk stage II colon cancer. <i>British Journal of Cancer</i> , 2005, 92, 24-29. | 2.9 | 16 |
| 210 | Pharmacogenetic interaction analysis of <i>VEGFR-2</i> and <i>IL-8</i> polymorphisms in advanced breast cancer patients treated with paclitaxel and bevacizumab. <i>Pharmacogenomics</i> , 2014, 15, 1985-1999. | 0.6 | 16 |
| 211 | EGFR and AKT1 overexpression are mutually exclusive and associated with a poor survival in resected gastric adenocarcinomas. <i>Cancer Biomarkers</i> , 2018, 21, 731-741. | 0.8 | 16 |
| 212 | Is a pharmacogenomic panel useful to estimate the risk of oxaliplatin-related neurotoxicity in colorectal cancer patients?. <i>Pharmacogenomics Journal</i> , 2019, 19, 465-472. | 0.9 | 16 |
| 213 | The Role of Anti-Angiogenics in Pre-Treated Metastatic BRAF-Mutant Colorectal Cancer: A Pooled Analysis. <i>Cancers</i> , 2020, 12, 1022. | 1.7 | 16 |
| 214 | Triplet Combination of Fluoropyrimidines, Oxaliplatin, and Irinotecan in the First-Line Treatment of Metastatic Colorectal Cancer. <i>Clinical Colorectal Cancer</i> , 2008, 7, 7-14. | 1.0 | 15 |
| 215 | Cigarettes smoking habit may reduce benefit from cetuximab-based treatment in advanced colorectal cancer patients. <i>Expert Opinion on Biological Therapy</i> , 2009, 9, 945-949. | 1.4 | 15 |
| 216 | A dose finding and pharmacokinetic study of capecitabine in combination with oxaliplatin and irinotecan in metastatic colorectal cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2009, 63, 965-969. | 1.1 | 15 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 217 | Angiogenesis Genotyping and Clinical Outcomes in Patients with Advanced Hepatocellular Carcinoma Receiving Sorafenib: The ALICE-2 Study. <i>Targeted Oncology</i> , 2020, 15, 115-126. | 1.7 | 15 |
| 218 | Biochemical Modulation by 5-Fluorouracil and 1-Folinic Acid of Tumor Uptake of Intra-Arterial 5-[¹²³ I]Iodo-2'-Deoxyuridine in Patients with Liver Metastases from Colorectal Cancer. <i>Acta Oncol</i> , 1996, 35, 941-945. | 0.8 | 14 |
| 219 | Protracted continuous infusion of 5-fluorouracil and low-dose leucovorin in patients with metastatic colorectal cancer resistant to 5-fluorouracil bolus-based chemotherapy: a phase II study. <i>Cancer Chemotherapy and Pharmacology</i> , 1999, 44, 159-163. | 1.1 | 14 |
| 220 | Cisplatin plus Gemcitabine as Adjuvant Chemotherapy for Radically Resected Non-Small-Cell Lung Cancer: A Pilot Study. <i>Clinical Lung Cancer</i> , 2009, 10, 53-57. | 1.1 | 14 |
| 221 | Liver metastases from colorectal cancer: how to best complement medical treatment with surgical approaches. <i>Future Oncology</i> , 2011, 7, 1299-1323. | 1.1 | 14 |
| 222 | Faithful Markers of Circulating Cancer Stem Cells: Is CD133 Sufficient for Validation in Clinics?. <i>Journal of Clinical Oncology</i> , 2011, 29, 3487-3488. | 0.8 | 14 |
| 223 | Second-line therapy for advanced pancreatic cancer: evaluation of prognostic factors and review of current literature. <i>Future Oncology</i> , 2016, 12, 901-908. | 1.1 | 14 |
| 224 | Benefit from anti-EGFRs in RAS and BRAF wild-type metastatic transverse colon cancer: a clinical and molecular proof of concept study. <i>ESMO Open</i> , 2019, 4, e000489. | 2.0 | 14 |
| 225 | Chemotherapeutic and antiangiogenic drugs beyond tumor progression in colon cancer: Evaluation of the effects of switched schedules and related pharmacodynamics. <i>Biochemical Pharmacology</i> , 2019, 164, 94-105. | 2.0 | 14 |
| 226 | Tumor Regression Grading Assessment in Locally Advanced Pancreatic Cancer After Neoadjuvant FOLFIRINOX: Interobserver Agreement and Prognostic Implications. <i>Frontiers in Oncology</i> , 2020, 10, 64. | 1.3 | 14 |
| 227 | Irinotecan in combination with thalidomide in patients with advanced solid tumors: a clinical study with pharmacodynamic and pharmacokinetic evaluation. <i>Cancer Chemotherapy and Pharmacology</i> , 2006, 58, 585-593. | 1.1 | 13 |
| 228 | Dissecting signaling pathways in hepatocellular carcinoma: new perspectives in medical therapy. <i>Future Oncology</i> , 2014, 10, 285-304. | 1.1 | 13 |
| 229 | Clinicopathological differences and survival outcomes with first-line therapy in patients with left-sided colon cancer and rectal cancer: Pooled analysis of 2879 patients from AGITC (MAX), COIN, FOCUS2, OPUS, CRYSTAL and COIN-B trials in the ARCAD database. <i>European Journal of Cancer</i> , 2018, 103, 205-213. | 1.3 | 13 |
| 230 | Validated Nomogram Predicting 6-Month Survival in Pancreatic Cancer Patients Receiving First-Line 5-Fluorouracil, Oxaliplatin, and Irinotecan. <i>Clinical Colorectal Cancer</i> , 2019, 18, e394-e401. | 1.0 | 13 |
| 231 | Tumour mutational burden, microsatellite instability, and actionable alterations in metastatic colorectal cancer: Next-generation sequencing results of TRIBE2 study. <i>European Journal of Cancer</i> , 2021, 155, 73-84. | 1.3 | 13 |
| 232 | Liquid biopsy to predict benefit from rechallenge with cetuximab (cet) + irinotecan (iri) in RAS/BRAF wild-type metastatic colorectal cancer patients (pts) with acquired resistance to first-line cet+iri: Final results and translational analyses of the CRICKET study by GONO.. <i>Journal of Clinical Oncology</i> , 2018, 36, 12007-12007. | 0.8 | 13 |
| 233 | Patients with NSCLC may display a low ratio of p.T790M vs. activating EGFR mutations in plasma at disease progression: implications for personalised treatment. <i>Oncotarget</i> , 2017, 8, 86056-86065. | 0.8 | 13 |
| 234 | EGF-receptor targeting with monoclonal antibodies in colorectal carcinomas: rationale for a pharmacogenomic approach. <i>Pharmacogenomics</i> , 2008, 9, 55-69. | 0.6 | 12 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|------|-----------|
| 235 | Randomised multicenter phase II study of two schedules of docetaxel and gemcitabine or cisplatin/gemcitabine followed by docetaxel as first line treatment for advanced non-small cell lung cancer. <i>Lung Cancer</i> , 2009, 66, 327-332. | 0.9 | 12 |
| 236 | Targeting Vascular Endothelial Growth Factor Pathway in First-Line Treatment of Metastatic Colorectal Cancer: State-of-the-Art and Future Perspectives in Clinical and Molecular Selection of Patients. <i>Current Cancer Drug Targets</i> , 2010, 10, 37-45. | 0.8 | 12 |
| 237 | KRAS Early Testing: Consensus Initiative and Cost-Effectiveness Evaluation for Metastatic Colorectal Patients in an Italian Setting. <i>PLoS ONE</i> , 2014, 9, e85897. | 1.1 | 12 |
| 238 | TAS-102 for the treatment of metastatic colorectal cancer. <i>Expert Review of Anticancer Therapy</i> , 2015, 15, 1283-1292. | 1.1 | 12 |
| 239 | Prognostic Value of ACVRL1 Expression in Metastatic Colorectal Cancer Patients Receiving First-line Chemotherapy With Bevacizumab: Results From the Triplet Plus Bevacizumab (TRIBE) Study. <i>Clinical Colorectal Cancer</i> , 2018, 17, e471-e488. | 1.0 | 12 |
| 240 | Integrated safety summary for trifluridine/tipiracil (TAS-102). <i>Anti-Cancer Drugs</i> , 2018, 29, 89-96. | 0.7 | 12 |
| 241 | Early Tumor Shrinkage and Depth of Response Evaluation in Metastatic Pancreatic Cancer Treated with First Line Chemotherapy: An Observational Retrospective Cohort Study. <i>Cancers</i> , 2019, 11, 939. | 1.7 | 12 |
| 242 | Prognostic Effect of Adenosine-related Genetic Variants in Metastatic Colorectal Cancer Treated With Bevacizumab-based Chemotherapy. <i>Clinical Colorectal Cancer</i> , 2019, 18, e8-e19. | 1.0 | 12 |
| 243 | Prognostic clinical factors in patients affected by non-small-cell lung cancer receiving Nivolumab. <i>Expert Opinion on Biological Therapy</i> , 2020, 20, 319-326. | 1.4 | 12 |
| 244 | Prognostic and Predictive Impact of Primary Tumor Sidedness for Previously Untreated Advanced Colorectal Cancer. <i>Journal of the National Cancer Institute</i> , 2021, 113, 1705-1713. | 3.0 | 12 |
| 245 | Rationale and Study Design of the PARERE Trial: Randomized phase II Study of Panitumumab Re-Treatment Followed by Regorafenib Versus the Reverse Sequence in RAS and BRAF Wild-Type Chemo-Refractory Metastatic Colorectal Cancer Patients. <i>Clinical Colorectal Cancer</i> , 2021, 20, 314-317. | 1.0 | 12 |
| 246 | FOLFOXIRI/bevacizumab (bev) versus doublets/bev as initial therapy of unresectable metastatic colorectal cancer (mCRC): A meta-analysis of individual patient data (IPD) from five randomized trials.. <i>Journal of Clinical Oncology</i> , 2020, 38, 4015-4015. | 0.8 | 12 |
| 247 | Surrogate Endpoints in Second-Line Trials of Targeted Agents in Metastatic Colorectal Cancer: A Literature-Based Systematic Review and Meta-Analysis. <i>Cancer Research and Treatment</i> , 2017, 49, 834-845. | 1.3 | 12 |
| 248 | Alpha-2B-interferon plus floxuridine in metastatic renal cell carcinoma a phase I-II study. <i>Cancer</i> , 1993, 72, 564-568. | 2.0 | 11 |
| 249 | First-line chemotherapy in metastatic colorectal cancer: new approaches and therapeutic algorithms. Always hit hard first?. <i>Current Opinion in Oncology</i> , 2008, 20, 459-465. | 1.1 | 11 |
| 250 | Second-line Treatment for Non-“Small-Cell Lung Cancer: One Size Does Not Fit All. <i>Clinical Lung Cancer</i> , 2010, 11, 320-327. | 1.1 | 11 |
| 251 | Prospective study of EGFR intron 1 (CA) _n repeats variants as predictors of benefit from cetuximab and irinotecan in chemo-refractory metastatic colorectal cancer (mCRC) patients. <i>Pharmacogenomics Journal</i> , 2014, 14, 322-327. | 0.9 | 11 |
| 252 | FOLFOXIRI and Bevacizumab for Metastatic Colorectal Cancer. <i>New England Journal of Medicine</i> , 2015, 372, 290-292. | 13.9 | 11 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 253 | Third-Line Chemotherapy with Irinotecan plus 5-Fluorouracil in Caucasian Metastatic Gastric Cancer Patients. <i>Oncology</i> , 2016, 91, 311-316. | 0.9 | 11 |
| 254 | Cardiac safety of adjuvant non-pegylated liposomal doxorubicin combined with cyclophosphamide and followed by paclitaxel in older breast cancer patients. <i>Breast</i> , 2017, 31, 186-191. | 0.9 | 11 |
| 255 | Circulating Tumor DNA Analysis in Colorectal Cancer: From Dream to Reality. <i>JCO Precision Oncology</i> , 2019, 3, 1-14. | 1.5 | 11 |
| 256 | Validated clinico-pathologic nomogram in the prediction of HER2 status in gastro-oesophageal cancer. <i>British Journal of Cancer</i> , 2019, 120, 522-526. | 2.9 | 11 |
| 257 | First-line gemcitabine plus nab-paclitaxel for elderly patients with metastatic pancreatic cancer: Crossing the frontier of age?. <i>European Journal of Cancer</i> , 2020, 137, 108-116. | 1.3 | 11 |
| 258 | Immune Profiling of Deficient Mismatch Repair Colorectal Cancer Tumor Microenvironment Reveals Different Levels of Immune System Activation. <i>Journal of Molecular Diagnostics</i> , 2020, 22, 685-698. | 1.2 | 11 |
| 259 | Clinical outcomes of NSCLC patients experiencing early immune-related adverse events to PD-1/PD-L1 checkpoint inhibitors leading to treatment discontinuation. <i>Cancer Immunology, Immunotherapy</i> , 2022, 71, 865-874. | 2.0 | 11 |
| 260 | Electrochemotherapy and its controversial results in patients with head and neck cancer. <i>Anticancer Research</i> , 2014, 34, 967-72. | 0.5 | 11 |
| 261 | Pharmacokinetic Optimisation of the Treatment of Cancer with High Dose Zidovudine. <i>Clinical Pharmacokinetics</i> , 1998, 34, 173-180. | 1.6 | 10 |
| 262 | Molecular and pathological characterization of the EZH2 rs3757441 single nucleotide polymorphism in colorectal cancer. <i>BMC Cancer</i> , 2015, 15, 874. | 1.1 | 10 |
| 263 | Single nucleotide polymorphisms in the IGF1R pathway are associated with outcome in mCRC patients enrolled in the FIRE trial. <i>International Journal of Cancer</i> , 2017, 141, 383-392. | 2.3 | 10 |
| 264 | Vinorelbine in BRAF V600E mutated metastatic colorectal cancer: a prospective multicentre phase II clinical study. <i>ESMO Open</i> , 2017, 2, e000241. | 2.0 | 10 |
| 265 | Prognostic impact of immune-microenvironment in colorectal liver metastases resected after triplets plus a biologic agent: A pooled analysis of five prospective trials. <i>European Journal of Cancer</i> , 2020, 135, 78-88. | 1.3 | 10 |
| 266 | <i>PNN</i> and <i>KCNQ1OT1</i> Can Predict the Efficacy of Adjuvant Fluoropyrimidine-Based Chemotherapy in Colorectal Cancer Patients. <i>Oncology Research</i> , 2020, 28, 631-644. | 0.6 | 10 |
| 267 | 5-Fluorouracil Administered as a 48-Hour Chronomodulated Infusion in Combination with Leucovorin and Cisplatin: A Randomized Phase II Study in Metastatic Colorectal Cancer. <i>Oncology</i> , 2001, 61, 28-35. | 0.9 | 9 |
| 268 | Pharmacokinetics, a main actor in a many-sided approach to severe 5-FU toxicity prediction. <i>British Journal of Clinical Pharmacology</i> , 2009, 67, 132-134. | 1.1 | 9 |
| 269 | Radium 223 dichloride: a multidisciplinary approach to metastatic castration-resistant prostate cancer. <i>Future Oncology</i> , 2015, 11, 323-331. | 1.1 | 9 |
| 270 | Clinical Significance of <i>TLR1</i> I602S Polymorphism for Patients with Metastatic Colorectal Cancer Treated with FOLFIRI plus Bevacizumab. <i>Molecular Cancer Therapeutics</i> , 2016, 15, 1740-1745. | 1.9 | 9 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 271 | An Italian cost-effectiveness analysis of paclitaxel albumin (nab-paclitaxel) + gemcitabine vs gemcitabine alone for metastatic pancreatic cancer patients: the APICE study. <i>Expert Review of Pharmacoeconomics and Outcomes Research</i> , 2018, 18, 435-446. | 0.7 | 9 |
| 272 | A polymorphism within the R-spondin 2 gene predicts outcome in metastatic colorectal cancer patients treated with FOLFIRI/bevacizumab: data from FIRE-3 and TRIBE trials. <i>European Journal of Cancer</i> , 2020, 131, 89-97. | 1.3 | 9 |
| 273 | FOLFOXIRI-Bevacizumab or FOLFOX-Panitumumab in Patients with Left-Sided <i>RAS/BRAF</i> Wild-Type Metastatic Colorectal Cancer: A Propensity Score-Based Analysis. <i>Oncologist</i> , 2021, 26, 302-309. | 1.9 | 9 |
| 274 | CEA increase as a marker of disease progression after first-line induction therapy in metastatic colorectal cancer patients. A pooled analysis of TRIBE and TRIBE2 studies. <i>British Journal of Cancer</i> , 2021, 125, 839-845. | 2.9 | 9 |
| 275 | Phase II Study of Oral Doxifluridine in Elderly Patients with Advanced Non-Small-Cell Lung Cancer. <i>American Journal of Clinical Oncology: Cancer Clinical Trials</i> , 1996, 19, 592-594. | 0.6 | 9 |
| 276 | Moving-Strip Abdomino-Pelvic Radiotherapy After cis-Platinum-Based Chemotherapy and Second-Look Operation A Feasibility Study in Advanced Ovarian Cancer. <i>American Journal of Clinical Oncology: Cancer Clinical Trials</i> , 1988, 11, 16-20. | 0.6 | 8 |
| 277 | Correlation of basal EGFR expression with pancreatic cancer grading but not with clinical outcome after gemcitabine-based treatment. <i>Annals of Oncology</i> , 2011, 22, 482-484. | 0.6 | 8 |
| 278 | Biomarkers and Response to Bevacizumab Letter. <i>Clinical Cancer Research</i> , 2014, 20, 1056-1057. | 3.2 | 8 |
| 279 | Tandem repeat variation near the <i>HIC1</i> (hypermethylated in cancer 1) promoter predicts outcome of oxaliplatin-based chemotherapy in patients with metastatic colorectal cancer. <i>Cancer</i> , 2017, 123, 4506-4514. | 2.0 | 8 |
| 280 | Pharmacokinetic analysis of metronomic capecitabine in refractory metastatic colorectal cancer patients. <i>Investigational New Drugs</i> , 2018, 36, 709-714. | 1.2 | 8 |
| 281 | Potential role of PIN1 genotypes in predicting benefit from oxaliplatin-based and irinotecan-based treatment in patients with metastatic colorectal cancer. <i>Pharmacogenomics Journal</i> , 2018, 18, 623-632. | 0.9 | 8 |
| 282 | Body Mass Index and Hormone Receptor Status Influence Recurrence Risk in HER2-Positive Early Breast Cancer Patients. <i>Clinical Breast Cancer</i> , 2020, 20, e89-e98. | 1.1 | 8 |
| 283 | Zebrafish Patient-Derived Xenografts Identify Chemo-Response in Pancreatic Ductal Adenocarcinoma Patients. <i>Cancers</i> , 2021, 13, 4131. | 1.7 | 8 |
| 284 | FOLFOXIRI plus bevacizumab versus FOLFIRI plus bevacizumab as initial treatment for metastatic colorectal cancer (TRIBE study): Updated survival results and final molecular subgroups analyses.. <i>Journal of Clinical Oncology</i> , 2015, 33, 3510-3510. | 0.8 | 8 |
| 285 | Oral Doxifluridine in Advanced Hepatocellular Carcinoma: A Phase II Study. <i>Oncology</i> , 2000, 59, 204-209. | 0.9 | 7 |
| 286 | First line chemotherapy with planned sequential administration of gemcitabine followed by docetaxel in elderly advanced non-small-cell lung cancer patients: a multicenter phase II study. <i>British Journal of Cancer</i> , 2008, 98, 558-563. | 2.9 | 7 |
| 287 | Review: Beyond KRAS: perspectives on new potential markers of intrinsic and acquired resistance to epidermal growth factor receptor inhibitors in metastatic colorectal cancer. <i>Therapeutic Advances in Medical Oncology</i> , 2009, 1, 167-181. | 1.4 | 7 |
| 288 | Discordant somatic and germline <i>VEGF-A</i> genotype in a cancer patient resistant to paclitaxel/bevacizumab with chemosensitive hepatic metastasis. <i>Pharmacogenomics</i> , 2009, 10, 1225-1229. | 0.6 | 7 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 289 | A Phase II Trial of Fixed-Dose Rate Gemcitabine plus Capecitabine in Metastatic/Advanced Biliary Tract Cancer Patients. <i>Oncology</i> , 2012, 82, 75-82. | 0.9 | 7 |
| 290 | EZH2 polymorphism and benefit from bevacizumab in colorectal cancer: another piece to the puzzle. <i>Annals of Oncology</i> , 2012, 23, 1370-1371. | 0.6 | 7 |
| 291 | Supportive care and not only palliative care in the route of cancer patients. <i>Supportive Care in Cancer</i> , 2013, 21, 657-658. | 1.0 | 7 |
| 292 | KRAS and BRAF genotyping of synchronous colorectal carcinomas. <i>Oncology Letters</i> , 2014, 7, 1532-1536. | 0.8 | 7 |
| 293 | Dedicated supportive care team at the oncology unit: a model of simultaneous care for cancer patients. <i>Supportive Care in Cancer</i> , 2014, 22, 867-868. | 1.0 | 7 |
| 294 | Lack of Benefit From Anti-EGFR Treatment in RAS and BRAF Wild-type Metastatic Colorectal Cancer With Mucinous Histology or Mucinous Component. <i>Clinical Colorectal Cancer</i> , 2019, 18, 116-124. | 1.0 | 7 |
| 295 | Clinical impact of first-line bevacizumab plus chemotherapy in metastatic colorectal cancer of mucinous histology: a multicenter, retrospective analysis on 685 patients. <i>Journal of Cancer Research and Clinical Oncology</i> , 2020, 146, 493-501. | 1.2 | 7 |
| 296 | Treatments after progression to first-line FOLFOXIRI and bevacizumab in metastatic colorectal cancer: a pooled analysis of TRIBE and TRIBE2 studies by GONO. <i>British Journal of Cancer</i> , 2021, 124, 183-190. | 2.9 | 7 |
| 297 | Capecitabine and mitomycin C may be an effective treatment option for third-line chemotherapy in advanced colorectal cancer. <i>Tumori</i> , 2006, 92, 384-8. | 0.6 | 7 |
| 298 | Cholinergic toxic syndrome by the anticancer drug irinotecan: Acetylcholinesterase does not play a major role. <i>Clinical Pharmacology and Therapeutics</i> , 2002, 71, 263-271. | 2.3 | 6 |
| 299 | Epirubicin/paclitaxel/etoposide in extensive-stage small-cell lung cancer: a phase II study. <i>British Journal of Cancer</i> , 2006, 94, 1263-1266. | 2.9 | 6 |
| 300 | Is erlotinib really active after failure of gefitinib in advanced non-small-cell lung cancer patients?. <i>Annals of Oncology</i> , 2009, 20, 790-791. | 0.6 | 6 |
| 301 | Host genetic variants in the IGF binding protein-3 impact on survival of patients with advanced gastric cancer treated with palliative chemotherapy. <i>Pharmacogenomics</i> , 2010, 11, 1247-1256. | 0.6 | 6 |
| 302 | Real-world dosing of regorafenib in metastatic colorectal cancer (mCRC): Interim analysis from the prospective, observational CORRELATE study. <i>Annals of Oncology</i> , 2017, 28, iii10. | 0.6 | 6 |
| 303 | Evaluation of Continuous Tumor-Size-Based End Points as Surrogates for Overall Survival in Randomized Clinical Trials in Metastatic Colorectal Cancer. <i>JAMA Network Open</i> , 2019, 2, e1911750. | 2.8 | 6 |
| 304 | Italian results of the PRECONNECT study: safety and efficacy of trifluridine/tipiracil in metastatic colorectal cancer. <i>Future Oncology</i> , 2021, 17, 2315-2324. | 1.1 | 6 |
| 305 | Prospective study of EGFR intron 1 CA tandem repeats to predict factor benefit from cetuximab and irinotecan. <i>Journal of Clinical Oncology</i> , 2012, 30, 3540-3540. | 0.8 | 6 |
| 306 | Phase II study of sequential chemotherapy with docetaxel-estramustine followed by mitoxantrone-prednisone in patients with advanced hormone-refractory prostate cancer. <i>British Journal of Cancer</i> , 2007, 97, 1613-1617. | 2.9 | 5 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|------|-----------|
| 307 | Use of pegfilgrastim support on day 9 to maintain relative dose intensity of chemotherapy in breast cancer patients receiving a day 1 and 8 CMF regimen. <i>Clinical and Translational Oncology</i> , 2009, 11, 842-848. | 1.2 | 5 |
| 308 | Refractory neuroendocrine tumor response to liposomal doxorubicin and capecitabine. <i>Nature Reviews Clinical Oncology</i> , 2009, 6, 670-674. | 12.5 | 5 |
| 309 | Phase II study of sequential cisplatin plus 5-fluorouracil/leucovorin (5-FU/LV) followed by irinotecan plus 5-FU/LV followed by docetaxel plus 5-FU/LV in patients with metastatic gastric or gastro-oesophageal junction adenocarcinoma. <i>Cancer Chemotherapy and Pharmacology</i> , 2010, 66, 559-566. | 1.1 | 5 |
| 310 | Capecitabine after gastrectomy for advanced gastric cancer: have we got the patient right?. <i>Annals of Oncology</i> , 2010, 21, 181. | 0.6 | 5 |
| 311 | PML as a potential predictive factor of oxaliplatin/fluoropyrimidine-based first line chemotherapy efficacy in colorectal cancer patients. <i>Journal of Cellular Physiology</i> , 2012, 227, 927-933. | 2.0 | 5 |
| 312 | Body mass index and impaired fasting blood glucose as predictive factor of time to progression (TTP) in cetuximab-based colorectal cancer treatment. <i>Cancer Biology and Therapy</i> , 2013, 14, 467-468. | 1.5 | 5 |
| 313 | FOLFOXIRI/Bevacizumab Versus FOLFIRI/Bevacizumab as First-Line Treatment in Unresectable Metastatic Colorectal Cancer: Results of Phase III Tribe Trial by Gono Group. <i>Annals of Oncology</i> , 2013, 24, iv21. | 0.6 | 5 |
| 314 | Small-Bowel Neuroendocrine Tumor and Retroperitoneal Fibrosis: Efficacy of Octreotide and Tamoxifen. <i>Tumori</i> , 2015, 101, e24-e28. | 0.6 | 5 |
| 315 | Topoisomerase 1 Promoter Variants and Benefit from Irinotecan in Metastatic Colorectal Cancer Patients. <i>Oncology</i> , 2016, 91, 283-288. | 0.9 | 5 |
| 316 | NOS2 polymorphisms in prediction of benefit from first-line chemotherapy in metastatic colorectal cancer patients. <i>PLoS ONE</i> , 2018, 13, e0193640. | 1.1 | 5 |
| 317 | Early modifications of circulating microRNAs levels in metastatic colorectal cancer patients treated with regorafenib. <i>Pharmacogenomics Journal</i> , 2019, 19, 455-464. | 0.9 | 5 |
| 318 | Pharmacodynamic biomarkers in metronomic chemotherapy: multiplex cytokine measurements in gastrointestinal cancer patients. <i>Clinical and Experimental Medicine</i> , 2021, 21, 149-159. | 1.9 | 5 |
| 319 | Exploring clinical and gene expression markers of benefit from FOLFOXIRI/bevacizumab in patients with BRAF-mutated metastatic colorectal cancer: Subgroup analyses of the TRIBE2 study. <i>European Journal of Cancer</i> , 2021, 153, 16-26. | 1.3 | 5 |
| 320 | In Pancreatic Adenocarcinoma Alpha-Synuclein Increases and Marks Peri-Neural Infiltration. <i>International Journal of Molecular Sciences</i> , 2022, 23, 3775. | 1.8 | 5 |
| 321 | Liver Transplantation for Metastatic Sinonasal Undifferentiated Carcinoma: A Case Report. <i>Transplantation Proceedings</i> , 2008, 40, 3821-3822. | 0.3 | 4 |
| 322 | Optimal approach to potentially resectable liver metastases from colorectal cancer. <i>Expert Review of Anticancer Therapy</i> , 2008, 8, 1533-1539. | 1.1 | 4 |
| 323 | Predictors of Benefit in Colorectal Cancer Treated With Cetuximab: Are We Getting "Lost in Translation"? <i>Journal of Clinical Oncology</i> , 2010, 28, e173-e174. | 0.8 | 4 |
| 324 | Pharmacogenomics of cetuximab in metastatic colorectal carcinoma. <i>Pharmacogenomics</i> , 2014, 15, 1701-1715. | 0.6 | 4 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|------|-----------|
| 325 | Not only chemotherapy in the second-line treatment of metastatic gastric cancer. <i>Annals of Oncology</i> , 2014, 25, 544-545. | 0.6 | 4 |
| 326 | Polymorphisms in Genes Involved in EGFR Turnover Are Predictive for Cetuximab Efficacy in Colorectal Cancer. <i>Molecular Cancer Therapeutics</i> , 2015, 14, 2374-2381. | 1.9 | 4 |
| 327 | The Winding Roadmap of Biomarkers toward Clinic: Lessons from Predictors of Resistance to Anti-EGFRs in Metastatic Colorectal Cancer. <i>International Journal of Molecular Sciences</i> , 2018, 19, 2298. | 1.8 | 4 |
| 328 | Health-related Quality of Life in the Phase III LUME-Colon 1 Study: Comparison and Interpretation of Results From EORTC QLQ-C30 Analyses. <i>Clinical Colorectal Cancer</i> , 2019, 18, 269-279.e5. | 1.0 | 4 |
| 329 | Impact of polymorphisms within genes involved in regulating DNA methylation in patients with metastatic colorectal cancer enrolled in three independent, randomised, open-label clinical trials: a meta-analysis from TRIBE, MAVERICC and FIRE-3. <i>European Journal of Cancer</i> , 2019, 111, 138-147. | 1.3 | 4 |
| 330 | AMPK variant, a candidate of novel predictor for chemotherapy in metastatic colorectal cancer: A meta-analysis using TRIBE, MAVERICC and FIRE3. <i>International Journal of Cancer</i> , 2019, 145, 2082-2090. | 2.3 | 4 |
| 331 | Polymorphisms within Immune Regulatory Pathways Predict Cetuximab Efficacy and Survival in Metastatic Colorectal Cancer Patients. <i>Cancers</i> , 2020, 12, 2947. | 1.7 | 4 |
| 332 | Single Nucleotide Polymorphisms in MiRNA Binding Sites of Nucleotide Excision Repair-Related Genes Predict Clinical Benefit of Oxaliplatin in FOLFOXIRI Plus Bevacizumab: Analysis of the TRIBE Trial. <i>Cancers</i> , 2020, 12, 1742. | 1.7 | 4 |
| 333 | Modified FOLFOXIRI plus cetuximab (cet) as induction treatment in unresectable metastatic colorectal cancer (mCRC) patients (pts): Preliminary results of the phase II randomized Macbeth trial by GONO group.. <i>Journal of Clinical Oncology</i> , 2014, 32, 3596-3596. | 0.8 | 4 |
| 334 | Regional Pharmacokinetic Selectivity of Intraperitoneal Cisplatin in Ovarian Cancer. <i>Oncology</i> , 1988, 45, 69-73. | 0.9 | 3 |
| 335 | Double 5-Fluorouracil Modulation with Folinic Acid and Recombinant Alpha-2B-Interferon. <i>American Journal of Clinical Oncology: Cancer Clinical Trials</i> , 1994, 17, 210-214. | 0.6 | 3 |
| 336 | Cancer Pharmacogenomics: Germline DNA, Tumor DNA, or Both?. <i>Current Pharmacogenomics and Personalized Medicine: the International Journal for Expert Reviews in Pharmacogenomics</i> , 2007, 5, 87-101. | 0.3 | 3 |
| 337 | How useful is adjuvant irinotecan in stage IV CRC?. <i>Nature Reviews Clinical Oncology</i> , 2010, 7, 190-191. | 12.5 | 3 |
| 338 | Cytotoxic triplets plus a biologic: state-of-the-art in maximizing the potential of up-front medical treatment of metastatic colorectal cancer. <i>Expert Opinion on Biological Therapy</i> , 2011, 11, 519-531. | 1.4 | 3 |
| 339 | Reply: Comment on "Histopathologic evaluation of liver metastases from colorectal cancer patients treated with FOLFOXIRI plus bevacizumab". <i>British Journal of Cancer</i> , 2013, 109, 3129-3130. | 2.9 | 3 |
| 340 | Adjuvant Chemoradiotherapy (Gemcitabine-based) in Pancreatic Adenocarcinoma: The Pisa University Experience. <i>Tumori</i> , 2017, 103, 577-582. | 0.6 | 3 |
| 341 | A retrospective study of trifluridine/tipiracil in pretreated metastatic colorectal cancer patients in clinical practice. <i>Colorectal Cancer</i> , 2018, 7, CRC01. | 0.8 | 3 |
| 342 | Clinical and molecular determinants of extrahepatic disease progression in patients with metastatic colorectal cancer with liver-limited metastases deemed initially unresectable. <i>ESMO Open</i> , 2019, 4, e000496. | 2.0 | 3 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 343 | Pharmacological effects of the simultaneous and sequential combinations of trifluridine/tipiracil (TAS-102) and 5-fluorouracil in fluoropyrimidine-sensitive colon cancer cells. <i>Investigational New Drugs</i> , 2020, 38, 92-98. | 1.2 | 3 |
| 344 | Management of Peritoneal Carcinomatosis With Cytoreductive Surgery Combined With Intraperitoneal Chemohyperthermia at a Novel Italian Center. <i>In Vivo</i> , 2020, 34, 2061-2066. | 0.6 | 3 |
| 345 | Duration of oxaliplatin-based adjuvant chemotherapy in patients with Stage III or high-risk Stage II resected colon cancer. <i>International Journal of Cancer</i> , 2020, 146, 2652-2654. | 2.3 | 3 |
| 346 | Clinical significance of enterocyte-specific gene polymorphisms as candidate markers of oxaliplatin-based treatment for metastatic colorectal cancer. <i>Pharmacogenomics Journal</i> , 2021, 21, 285-295. | 0.9 | 3 |
| 347 | Management of Thyrotoxicosis Induced by PD1 or PD-L1 Blockade. <i>Journal of the Endocrine Society</i> , 2021, 5, bvab093. | 0.1 | 3 |
| 348 | Random survival forests identify pathways with polymorphisms predictive of survival in KRAS mutant and KRAS wild-type metastatic colorectal cancer patients. <i>Scientific Reports</i> , 2021, 11, 12191. | 1.6 | 3 |
| 349 | Novel prognostic markers for epithelioid malignant pleural mesothelioma. <i>Journal of Clinical Oncology</i> , 2017, 35, e20028-e20028. | 0.8 | 3 |
| 350 | A pharmacogenetic interaction analysis of bevacizumab with paclitaxel in advanced breast cancer patients. <i>Npj Breast Cancer</i> , 2022, 8, 33. | 2.3 | 3 |
| 351 | Genetic variants involved in the cGAS-STING pathway predict outcome in patients with metastatic colorectal cancer: Data from FIRE-3 and TRIBE trials. <i>European Journal of Cancer</i> , 2022, 172, 22-30. | 1.3 | 3 |
| 352 | 5-Fluorouracil Administered as a 48-Hour Semiintermittent Infusion in Combination With Leucovorin, Cisplatin and Epirubicin. <i>American Journal of Clinical Oncology: Cancer Clinical Trials</i> , 2004, 27, 101-105. | 0.6 | 2 |
| 353 | Molecular predictive factors of response to anti-EGFR antibodies in colorectal cancer patients. <i>European Journal of Cancer, Supplement</i> , 2008, 6, 86-90. | 2.2 | 2 |
| 354 | Do we need biopsies of metastases for colorectal cancer patients?. <i>British Journal of Cancer</i> , 2009, 101, 374-375. | 2.9 | 2 |
| 355 | Treatments after first progression in metastatic colorectal cancer. A literature review and evidence-based algorithm. <i>Cancer Treatment Reviews</i> , 2021, 92, 102135. | 3.4 | 2 |
| 356 | Beyond the Guidelines: The Grey Zones of the Management of Gastric Cancer. Consensus Statements from the Gastric Cancer Italian Network (GAIN). <i>Cancers</i> , 2021, 13, 1304. | 1.7 | 2 |
| 357 | Detailing the ultrastructure's increase of prion protein in pancreatic adenocarcinoma. <i>World Journal of Gastroenterology</i> , 2021, 27, 7324-7339. | 1.4 | 2 |
| 358 | Reply: KRAS status analysis and anti-EGFR therapies: is comprehensiveness a biologist's fancy or a clinical necessity?. <i>British Journal of Cancer</i> , 2010, 102, 1076-1077. | 2.9 | 1 |
| 359 | Pharmacogenetic Concerns in Metastatic Colorectal Cancer Therapy. <i>Current Colorectal Cancer Reports</i> , 2012, 8, 263-271. | 1.0 | 1 |
| 360 | O-0023 Phase 3 Correct Trial of Regorafenib in Metastatic Colorectal Cancer (mCRC). <i>Annals of Oncology</i> , 2012, 23, iv15-iv16. | 0.6 | 1 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 361 | Long-Survivors with Lung Metastases and Kras Mutations Have an Increased Risk to Develop Brain Metastases From Colorectal Cancer. <i>Annals of Oncology</i> , 2013, 24, iv15. | 0.6 | 1 |
| 362 | Oral multikinase inhibitor regorafenib for the treatment of patients with metastatic colorectal cancer. <i>Colorectal Cancer</i> , 2013, 2, 411-417. | 0.8 | 1 |
| 363 | Comment on: "Nab-paclitaxel plus gemcitabine for metastatic pancreatic adenocarcinoma after Folfirinox failure: an AGEO prospective multicentre cohort"™. <i>British Journal of Cancer</i> , 2016, 114, e8-e8. | 2.9 | 1 |
| 364 | Optimization of biomarkers-based classification scores as progression-free survival predictors: an intuitive graphical representation. <i>Future Science OA</i> , 2018, 4, FSO346. | 0.9 | 1 |
| 365 | BRAF mutant metastatic colorectal cancers: new arrows in our quiver. <i>Annals of Translational Medicine</i> , 2019, 7, S367-S367. | 0.7 | 1 |
| 366 | Atypical KRAS Mutations in Metastatic Colorectal Cancer. <i>JCO Precision Oncology</i> , 2019, 3, 1-11. | 1.5 | 1 |
| 367 | RNA-Binding Protein Polymorphisms as Novel Biomarkers to Predict Outcomes of Metastatic Colorectal Cancer: A Meta-analysis from TRIBE, FIRE-3, and MAVERICC. <i>Molecular Cancer Therapeutics</i> , 2021, 20, 1153-1160. | 1.9 | 1 |
| 368 | Germ line polymorphisms of genes involved in pluripotency transcription factors predict efficacy of cetuximab in metastatic colorectal cancer. <i>European Journal of Cancer</i> , 2021, 150, 133-142. | 1.3 | 1 |
| 369 | Variations of circulating KRAS amount as a biomarker to monitor chemotherapy response in pancreatic cancer.. <i>Journal of Clinical Oncology</i> , 2017, 35, e15794-e15794. | 0.8 | 1 |
| 370 | FLOaTing toward new standards in locally advanced resectable gastroesophageal cancer. <i>Journal of Thoracic Disease</i> , 2019, 11, 5694-5700. | 0.6 | 1 |
| 371 | Reply to the Letter to the Editor on "Cost-opportunity analysis in clinical oncology: from the "wild west" to a correct integration of the disciplines, avoiding the "war of the worlds"™, by D. Tassinari et al. (<i>Ann Oncol</i> 2006; 17: 876). <i>Annals of Oncology</i> , 2006, 17, 877-878. | 0.6 | 0 |
| 372 | A Pilot Study of a Day One and Eight Every Three Weeks Administration of Docetaxel in Metastatic Cancer Patients. <i>Tumori</i> , 2007, 93, 145-149. | 0.6 | 0 |
| 373 | Chemotherapy intensification. <i>Current Colorectal Cancer Reports</i> , 2007, 3, 116-122. | 1.0 | 0 |
| 374 | Long-Term Outcome of Unresectable Metastatic Colorectal Cancer: Does "Adjuvant" Chemotherapy Play a Role After Resection?. <i>Annals of Surgery</i> , 2009, 250, 655. | 2.1 | 0 |
| 375 | Prognostic Value of CD133 Caused by Mutant K-Ras and B-Raf"™ Letter. <i>Clinical Cancer Research</i> , 2012, 18, 4473-4473. | 3.2 | 0 |
| 376 | Resectable liver metastases from colorectal cancer: where we are now and where do we go from here?. <i>Colorectal Cancer</i> , 2012, 1, 397-411. | 0.8 | 0 |
| 377 | Upfront Chemotherapy Regimens in Unresectable Disease: One, Two, or Three Cytotoxics?. <i>Current Colorectal Cancer Reports</i> , 2012, 8, 153-160. | 1.0 | 0 |
| 378 | PD-0009 Primary Tumor Location is a Major Independent Prognostic Factor for Mrcr Patients. <i>Annals of Oncology</i> , 2012, 23, iv22. | 0.6 | 0 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 379 | P-0262 Prospective Evaluation of Candidate Snps of Vegf/Vegfr Pathway in Metastatic Colorectal Cancer Patients Treated with First-Line Folfiri Plus Bevacizumab (BV). <i>Annals of Oncology</i> , 2012, 23, iv105-iv106. | 0.6 | 0 |
| 380 | P-0263 Prospective Study of Egfr Intron 1 CA Tandem Repeats as Predictive Factor of Benefit from Cetuximab and Irinotecan. <i>Annals of Oncology</i> , 2012, 23, iv106. | 0.6 | 0 |
| 381 | Prospective Analysis of the Early Modulation of Plasma Amphiregulin During Treatment with Cetuximab and Irinotecan in Metastatic Colorectal Cancer Patients. <i>Annals of Oncology</i> , 2013, 24, iv28. | 0.6 | 0 |
| 382 | Conference Scene: Annual Meeting of the American Society of Clinical Oncology. <i>Colorectal Cancer</i> , 2013, 2, 401-404. | 0.8 | 0 |
| 383 | TRIBE study: are all three cytotoxic drugs crucial? â€œ Authors' reply. <i>Lancet Oncology</i> , The, 2015, 16, e578-e579. | 5.1 | 0 |
| 384 | A still missing piece of the FIRE-3 puzzle. <i>Lancet Oncology</i> , The, 2016, 17, e515. | 5.1 | 0 |
| 385 | RET rearrangements define a new and rare molecular subtype of metastatic colorectal cancer (mCRC). <i>Annals of Oncology</i> , 2017, 28, iii140-iii141. | 0.6 | 0 |
| 386 | Prognostic and predictive role of neutrophils/lymphocytes ratio in metastatic colorectal cancer: A retrospective analysis of the TRIBE study by Gono. <i>Annals of Oncology</i> , 2017, 28, iii141-iii142. | 0.6 | 0 |
| 387 | Regorafenib: lights and shadows of antiangiogenic therapies in gastric cancer. <i>Translational Gastroenterology and Hepatology</i> , 2017, 2, 11-11. | 1.5 | 0 |
| 388 | Reply to Ugo De Giorgi, Vincenza Conteduca, and Emanuela Scarpi's Letter to the Editor re: Marzia Del Re, Elisa Biasco, Stefania Crucitta, et al. The Detection of Androgen Receptor Splice Variant 7 in Plasma-derived Exosomal RNA Strongly Predicts Resistance to Hormonal Therapy in Metastatic Prostate Cancer Patients. <i>Eur Urol</i> 2017;71:680â€œ7. <i>European Urology</i> , 2018, 73, e11-e12. | 0.9 | 0 |
| 389 | <i>BRAF</i> V600E Mutation as a Negative Prognostic Determinant in Resected Colorectal Liver Metastases. <i>JAMA Surgery</i> , 2018, 153, 1162. | 2.2 | 0 |
| 390 | Perioperative Morbidity Following Cytoreductive Surgery Combined with Intraperitoneal Chemohyperthermia in a Novel Italian Centre. <i>European Journal of Surgical Oncology</i> , 2020, 46, e160-e161. | 0.5 | 0 |
| 391 | TRIBE2 results and toxicity â€œ Authors' reply. <i>Lancet Oncology</i> , The, 2020, 21, e300-e301. | 5.1 | 0 |
| 392 | First-Line Systemic Chemotherapy with Folfoxiri Followed by Radical Surgical Resection of Metastases for the Treatment of Unresectable Metastatic Colorectal Cancer Patients. , 2009, , 285-293. | | 0 |
| 393 | The Role of Metronomic Chemotherapy in the Treatment of Metastatic Colorectal Cancer Patients. , 2014, , 135-142. | | 0 |
| 394 | Bevacizumab as maintenance therapy in mCRC: Interpreting results of the MOMA trial. <i>Oncotarget</i> , 2019, 10, 2791-2792. | 0.8 | 0 |
| 395 | Clinical and economic effect of administration of red blood product transfusions in an outpatient supportive care cancer service. <i>Biomedical Reports</i> , 2020, 12, 199-203. | 0.9 | 0 |