

Colin Dourish

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4025383/colin-dourish-publications-by-year.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

188
papers

9,081
citations

50
h-index

88
g-index

218
ext. papers

9,554
ext. citations

5.2
avg, IF

5.63
L-index

#	Paper	IF	Citations
188	The effects of lisdexamfetamine dimesylate on eating behaviour and homeostatic, reward and cognitive processes in women with binge-eating symptoms: an experimental medicine study.. <i>Translational Psychiatry</i> , 2022 , 12, 9	8.6	2
187	Validation of the P1vital [®] Faces Set for Use as Stimuli in Tests of Facial Emotion Recognition.. <i>Frontiers in Psychiatry</i> , 2022 , 13, 663763	5	
186	Utility of an experimental medicine model to evaluate efficacy, side-effects and mechanism of action of novel treatments for obesity and binge-eating disorder.. <i>Appetite</i> , 2022 , 106087	4.5	
185	The clinical effectiveness of using a predictive algorithm to guide antidepressant treatment in primary care (PReDicT): an open-label, randomised controlled trial. <i>Neuropsychopharmacology</i> , 2021 , 46, 1307-1314	8.7	11
184	Lisdexamfetamine and binge-eating disorder: A systematic review and meta-analysis of the preclinical and clinical data with a focus on mechanism of drug action in treating the disorder. <i>European Neuropsychopharmacology</i> , 2021 , 53, 49-78	1.2	3
183	Associations between inattention and impulsivity ADHD symptoms and disordered eating risk in a community sample of young adults. <i>Psychological Medicine</i> , 2020 , 1-10	6.9	2
182	T108. AUT00206, A NOVEL KV3 CHANNEL MODULATOR, REDUCES KETAMINE-INDUCED BOLD SIGNALLING IN HEALTHY MALE VOLUNTEERS: A RANDOMISED PLACEBO-CONTROLLED CROSSOVER TRIAL. <i>Schizophrenia Bulletin</i> , 2019 , 45, S245-S246	1.3	5
181	Interoception and disordered eating: A systematic review. <i>Neuroscience and Biobehavioral Reviews</i> , 2019 , 107, 166-191	9	24
180	Value of monitoring negative emotional bias in primary care in England for personalised antidepressant treatment: a modelling study. <i>Evidence-Based Mental Health</i> , 2019 , 22, 145-152	11.1	3
179	Top-down guidance of attention to food cues is enhanced in individuals with overweight/obesity and predicts change in weight at one-year follow up. <i>International Journal of Obesity</i> , 2019 , 43, 1849-1858	5.5	6
178	Predicting treatment response to antidepressant medication using early changes in emotional processing. <i>European Neuropsychopharmacology</i> , 2019 , 29, 66-75	1.2	24
177	The 5-HT receptor agonist meta-chlorophenylpiperazine (mCPP) reduces palatable food consumption and BOLD fMRI responses to food images in healthy female volunteers. <i>Psychopharmacology</i> , 2018 , 235, 257-267	4.7	8
176	Associations Between Core Symptoms of Attention Deficit Hyperactivity Disorder and Both Binge and Restrictive Eating. <i>Frontiers in Psychiatry</i> , 2018 , 9, 103	5	18
175	Attention Deficit Hyperactivity Disorder (ADHD) and disordered eating behaviour: A systematic review and a framework for future research. <i>Clinical Psychology Review</i> , 2017 , 53, 109-121	10.8	47
174	Interactions between metabolic, reward and cognitive processes in appetite control: Implications for novel weight management therapies. <i>Journal of Psychopharmacology</i> , 2017 , 31, 1460-1474	4.6	35
173	The effects of using the PReDicT Test to guide the antidepressant treatment of depressed patients: study protocol for a randomised controlled trial. <i>Trials</i> , 2017 , 18, 558	2.8	20
172	Multidisciplinary approaches to the study of eating disorders and obesity: Recent progress in research and development and future prospects. <i>Journal of Psychopharmacology</i> , 2017 , 31, 1383-1387	4.6	

171	Investigating virtual reality navigation in amnesic mild cognitive impairment using fMRI. <i>Aging, Neuropsychology, and Cognition</i> , 2016 , 23, 196-217	2.1	15
170	Cognitive and oculomotor performance in subjects with low and high schizotypy: implications for translational drug development studies. <i>Translational Psychiatry</i> , 2016 , 6, C	8.6	11
169	Test-retest reliability and effects of repeated testing and satiety on performance of an Emotional Test Battery. <i>Journal of Clinical and Experimental Neuropsychology</i> , 2016 , 38, 416-33	2.1	14
168	Comparing the actions of lanicemine and ketamine in depression: key role of the anterior cingulate. <i>European Neuropsychopharmacology</i> , 2016 , 26, 994-1003	1.2	76
167	Satiation attenuates BOLD activity in brain regions involved in reward and increases activity in dorsolateral prefrontal cortex: an fMRI study in healthy volunteers. <i>American Journal of Clinical Nutrition</i> , 2015 , 101, 697-704	7	44
166	Effects of awareness that food intake is being measured by a universal eating monitor on the consumption of a pasta lunch and a cookie snack in healthy female volunteers. <i>Appetite</i> , 2015 , 92, 247-54	4.5	14
165	P.2.f.013 A precision medicine approach to antidepressant treatment in depression. <i>European Neuropsychopharmacology</i> , 2015 , 25, S441	1.2	1
164	Alterations in working memory networks in amnesic mild cognitive impairment. <i>Aging, Neuropsychology, and Cognition</i> , 2015 , 22, 106-27	2.1	25
163	Effects of the 5-HT _{2C} receptor agonist meta-chlorophenylpiperazine on appetite, food intake and emotional processing in healthy volunteers. <i>Psychopharmacology</i> , 2014 , 231, 2449-59	4.7	21
162	Effects of risperidone, amisulpride and nicotine on eye movement control and their modulation by schizotypy. <i>Psychopharmacology</i> , 2013 , 227, 331-45	4.7	30
161	Neuroimaging Approaches to the Understanding of Depression and the Identification of Novel Antidepressants 2013 , 343-411		1
160	Combined NK ₁ antagonism and serotonin reuptake inhibition: effects on emotional processing in humans. <i>Journal of Psychopharmacology</i> , 2013 , 27, 435-43	4.6	16
159	Evaluation of the effects of venlafaxine and pregabalin on the carbon dioxide inhalation models of Generalised Anxiety Disorder and panic. <i>Journal of Psychopharmacology</i> , 2013 , 27, 135-45	4.6	10
158	Changes in cardiovascular function after venlafaxine but not pregabalin in healthy volunteers: a double-blind, placebo-controlled study of orthostatic challenge, blood pressure and heart rate. <i>Human Psychopharmacology</i> , 2013 , 28, 562-75	2.3	10
157	The effects of ketamine and risperidone on eye movement control in healthy volunteers. <i>Translational Psychiatry</i> , 2013 , 3, e334	8.6	19
156	The effect of a clinically effective and non-effective dose of lorazepam on 7.5% CO ₂ induced anxiety. <i>Human Psychopharmacology</i> , 2012 , 27, 540-8	2.3	8
155	A validation of cognitive biomarkers for the early identification of cognitive enhancing agents in schizotypy: a three-center double-blind placebo-controlled study. <i>European Neuropsychopharmacology</i> , 2012 , 22, 469-81	1.2	32
154	Validation of experimental medicine methods in psychiatry: the P1vital approach and experience. <i>Biochemical Pharmacology</i> , 2011 , 81, 1435-41	6	11

153	Preliminary evidence of anxiolytic effects of the CRF(1) receptor antagonist R317573 in the 7.5% CO(2) proof-of-concept experimental model of human anxiety. <i>Journal of Psychopharmacology</i> , 2011 , 25, 1199-206	4.6	47
152	Validating the inhalation of 7.5% CO(2) in healthy volunteers as a human experimental medicine: a model of generalized anxiety disorder (GAD). <i>Journal of Psychopharmacology</i> , 2011 , 25, 1192-8	4.6	30
151	Agomelatine facilitates positive versus negative affective processing in healthy volunteer models. <i>Journal of Psychopharmacology</i> , 2011 , 25, 1159-67	4.6	50
150	Effects of risperidone, amisulpride and nicotine on eye movement control and their modulation by schizotypy. <i>Pharmacopsychiatry</i> , 2011 , 44,	2	2
149	The effects of sibutramine on the microstructure of eating behaviour and energy expenditure in obese women. <i>Journal of Psychopharmacology</i> , 2010 , 24, 99-109	4.6	30
148	P.2.c.021 Agomelatine facilitates positive emotional processing in healthy volunteers. <i>European Neuropsychopharmacology</i> , 2010 , 20, S379-S380	1.2	
147	Anti-obesity Drugs: From Animal Models to Clinical Efficacy 2008 , 271-315		4
146	Antagonists of the human adenosine A2A receptor. Part 3: Design and synthesis of pyrazolo[3,4-d]pyrimidines, pyrrolo[2,3-d]pyrimidines and 6-arylpurines. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2008 , 18, 2924-9	2.9	54
145	Antagonists of the human adenosine A2A receptor. Part 1: Discovery and synthesis of thieno[3,2-d]pyrimidine-4-methanone derivatives. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2008 , 18, 2916-9	2.9	36
144	Antagonists of the human adenosine A2A receptor. Part 2: Design and synthesis of 4-arylthieno[3,2-d]pyrimidine derivatives. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2008 , 18, 2920-3	2.9	26
143	Serotonin 1B and 2C receptor interactions in the modulation of feeding behaviour in the mouse. <i>Psychopharmacology</i> , 2006 , 185, 45-57	4.7	36
142	Pyrrolo(iso)quinoline derivatives as 5-HT(2C) receptor agonists. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2006 , 16, 677-80	2.9	22
141	Synthesis and biological evaluation of novel hexahydro-pyrido[3,2-b:4,5]pyrrolo[1,2-a]pyrazines as potent and selective 5-HT(2C) receptor agonists. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2006 , 16, 1207-11	2.9	36
140	Identification of 4-methyl-1,2,3,4,10,10a-hexahydropyrazino[1,2-a]indoles as 5-HT2C receptor agonists. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2005 , 15, 3604-8	2.9	21
139	Tonic regulation of satiety by 5-HT receptors in the mouse: converging evidence from behavioural and c-fos immunoreactivity studies?. <i>European Journal of Neuroscience</i> , 2004 , 19, 3017-25	3.5	27
138	Indoline derivatives as 5-HT(2C) receptor agonists. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2004 , 14, 2367-70	2.9	42
137	Reduced hypophagic effects of d-fenfluramine and the 5-HT2C receptor agonist mCPP in 5-HT1B receptor knockout mice. <i>Psychopharmacology</i> , 2004 , 176, 39-49	4.7	50
136	mCPP-induced hyperactivity in 5-HT2C receptor mutant mice is mediated by activation of multiple 5-HT receptor subtypes. <i>Neuropharmacology</i> , 2004 , 46, 663-71	5.5	41

135	Serotonin receptor ligands and the treatment of obesity. <i>Current Opinion in Investigational Drugs</i> , 2004 , 5, 377-88		35
134	Oral administration of the 5-HT _{2C} receptor agonist, mCPP, reduces body weight gain in rats over 28 days as a result of maintained hypophagia. <i>Psychopharmacology</i> , 2003 , 167, 274-80	4.7	62
133	Preferential effects of the cannabinoid CB ₁ receptor antagonist, SR 141716, on food intake and body weight gain of obese (fa/fa) compared to lean Zucker rats. <i>Psychopharmacology</i> , 2003 , 167, 103-114	4.7	177
132	5-HT _{1B} receptor knockout mice show a compensatory reduction in 5-HT _{2C} receptor function. <i>European Journal of Neuroscience</i> , 2003 , 17, 185-90	3.5	34
131	Potential for antipsychotic and psychotomimetic effects of A _{2A} receptor modulation. <i>Neurology</i> , 2003 , 61, S88-93	6.5	13
130	Discovery of nonxanthine adenosine A _{2A} receptor antagonists for the treatment of Parkinson's disease. <i>Neurology</i> , 2003 , 61, S101-6	6.5	68
129	Serotonin 2C receptor agonists and the behavioural satiety sequence in mice. <i>Pharmacology Biochemistry and Behavior</i> , 2002 , 71, 691-700	3.9	103
128	5-HT _{1B} receptors modulate components of satiety in the rat: behavioural and pharmacological analyses of the selective serotonin 1B agonist CP-94,253. <i>Psychopharmacology</i> , 2002 , 164, 49-60	4.7	56
127	Case series: amantadine open-label treatment of impulsive and aggressive behavior in hospitalized children with developmental disabilities. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2001 , 40, 654-7	7.2	20
126	Evidence that hypophagia induced by d-fenfluramine and d-norfenfluramine in the rat is mediated by 5-HT _{2C} receptors. <i>Neuropharmacology</i> , 2001 , 41, 200-9	5.5	179
125	Double-blind, placebo-controlled study of amantadine hydrochloride in the treatment of children with autistic disorder. <i>Journal of the American Academy of Child and Adolescent Psychiatry</i> , 2001 , 40, 658-65	7.2	147
124	Comparative effects of continuous infusion of mCPP, Ro 60-0175 and d-fenfluramine on food intake, water intake, body weight and locomotor activity in rats. <i>British Journal of Pharmacology</i> , 2000 , 130, 1305-14	8.6	64
123	Similarities in the action of Ro 60-0175, a 5-HT _{2C} receptor agonist and d-fenfluramine on feeding patterns in the rat. <i>Psychopharmacology</i> , 2000 , 152, 256-67	4.7	125
122	Measurement of anxiety in transgenic mice. <i>Reviews in the Neurosciences</i> , 2000 , 11, 59-74	4.7	40
121	5-HT _{2C} receptor modulation and the treatment of obesity. <i>Diabetes, Obesity and Metabolism</i> , 1999 , 1, 207-14	6.7	37
120	Reduced satiating effect of d-fenfluramine in serotonin 5-HT _{2C} receptor mutant mice. <i>Psychopharmacology</i> , 1999 , 143, 309-14	4.7	244
119	Utility of ethological analysis to overcome locomotor confounds in elevated maze models of anxiety. <i>Neuroscience and Biobehavioral Reviews</i> , 1998 , 23, 265-71	9	154
118	Child and learning disability psychopharmacology. <i>Journal of Psychopharmacology</i> , 1997 , 11, 291-4	4.6	19

117	Devazepide attenuates dl-fenfluramine-induced suppression of gastric emptying but not food intake in the 17 h food-deprived rat. <i>Physiology and Behavior</i> , 1997 , 62, 545-50	3.5	9
116	WAY100635 and latent inhibition in the rat: selective effects at preexposure. <i>Behavioural Brain Research</i> , 1997 , 88, 51-7	3.4	15
115	Behavioural and pharmacological characterisation of the canopy stretched attend posture test as a model of anxiety in mice and rats. <i>Psychopharmacology</i> , 1997 , 133, 29-38	4.7	65
114	Blockade of pre- and post-synaptic 5-HT _{1A} receptors does not modify the effect of fluoxetine or 5-hydroxytryptophan on ethanol and food intake in rats. <i>Psychopharmacology</i> , 1997 , 134, 55-63	4.7	28
113	Electrophysiological, biochemical, neurohormonal and behavioural studies with WAY-100635, a potent, selective and silent 5-HT _{1A} receptor antagonist. <i>Behavioural Brain Research</i> , 1996 , 73, 337-53	3.4	404
112	Lack of interaction between devazepide and 8-OH-DPAT-induced hyperphagia in the rat. <i>Physiology and Behavior</i> , 1996 , 60, 1337-40	3.5	
111	Effects of 5-HT _{1A} receptor agonists, partial agonists and a silent antagonist on the performance of the conditioned emotional response test in the rat. <i>Psychopharmacology</i> , 1996 , 128, 293-303	4.7	47
110	Effects of the selective angiotensin II receptor antagonists losartan and PD123177 in animal models of anxiety and memory. <i>Psychopharmacology</i> , 1996 , 126, 206-18	4.7	28
109	Behavioural evidence that d-fenfluramine-induced anorexia in the rat is not mediated by the 5-HT _{1A} receptor subtype. <i>Psychopharmacology</i> , 1996 , 125, 168-75	4.7	34
108	Dissociation between cognitive and motor/motivational deficits in the delayed matching to position test: effects of scopolamine, 8-OH-DPAT and EAA antagonists. <i>Psychopharmacology</i> , 1995 , 122, 268-80	4.7	78
107	Analysis of glutamate in striatal microdialysates using capillary electrophoresis and laser-induced fluorescence detection. <i>Journal of Chromatography A</i> , 1995 , 700, 81-87	4.5	28
106	Comparisons between the effects of 5-HT and DL-fenfluramine on food intake and gastric emptying in the rat. <i>Pharmacology Biochemistry and Behavior</i> , 1995 , 50, 581-5	3.9	17
105	Multiple serotonin receptors: opportunities for new treatments for obesity?. <i>Obesity</i> , 1995 , 3 Suppl 4, 449S-462S		84
104	Differential effects of WAY-100135 on the decrease in 5-hydroxytryptamine release induced by buspirone and NAN-190. <i>European Journal of Pharmacology</i> , 1995 , 276, 281-4	5.3	12
103	Benefits of ethological analysis of behaviour. <i>Trends in Pharmacological Sciences</i> , 1995 , 16, 260-1	13.2	4
102	Behavioural and pharmacological characterisation of the elevated "zero-maze" as an animal model of anxiety. <i>Psychopharmacology</i> , 1994 , 116, 56-64	4.7	519
101	An examination of the behavioural specificity of hypophagia induced by 5-HT _{1B} , 5-HT _{1C} and 5-HT ₂ receptor agonists using the post-prandial satiety sequence in rats. <i>Psychopharmacology</i> , 1994 , 113, 369-477	4.7	89
100	Effects of the selective 5-HT _{1A} receptor antagonist WAY100135 and its enantiomers on 8-OH-DPAT-induced hyperglycaemia in conscious rats. <i>European Journal of Pharmacology</i> , 1994 , 254, 133-9	5.3	11

99	Inhibition of 8-OH-DPAT-induced elevation of plasma corticotrophin by the 5-HT1A receptor antagonist WAY100635. <i>European Journal of Pharmacology</i> , 1994 , 264, 95-7	5.3	31
98	The influence of 5-hydroxytryptamine re-uptake blockade on CCK receptor antagonist effects in the rat elevated zero-maze. <i>European Journal of Pharmacology</i> , 1994 , 271, 403-11	5.3	34
97	In vivo characterization of the putative 5-HT1A receptor antagonist SDZ 216,525 using two models of somatodendritic 5-HT1A receptor function. <i>Neuropharmacology</i> , 1994 , 33, 359-66	5.5	15
96	The angiotensin converting enzyme inhibitors captopril and enalapril inhibit apomorphine-induced oral stereotypy in the rat. <i>Neuroscience</i> , 1994 , 58, 799-805	3.9	18
95	Implications of Eshkol-Wachman movement notation for behavioural pharmacology. <i>Behavioral and Brain Sciences</i> , 1994 , 17, 754-754	0.9	
94	Neurochemical profile of the selective and silent 5-HT1A receptor antagonist WAY100135: an in vivo microdialysis study. <i>European Journal of Pharmacology</i> , 1993 , 239, 195-202	5.3	76
93	Na ⁺ channels. <i>Trends in Pharmacological Sciences</i> , 1993 , 14, 41	13.2	34
92	Silent 5-HT1A receptor antagonists: utility as research tools and therapeutic agents. <i>Trends in Pharmacological Sciences</i> , 1993 , 14, 41-8	13.2	161
91	The effects of CCKA and CCKB antagonists on activity in the black/white exploration model of anxiety in mice. <i>Physiology and Behavior</i> , 1993 , 54, 689-93	3.5	46
90	Drinking induced by subcutaneous injection of angiotensin II in the rat is blocked by the selective AT1 receptor antagonist DuP 753 but not by the selective AT2 receptor antagonist WL 19. <i>European Journal of Pharmacology</i> , 1992 , 211, 113-6	5.3	34
89	Effects of the putative 5-HT1A receptor antagonist NAN-190 on free feeding and on feeding induced by the 5-HT1A receptor agonist 8-OH-DPAT in the rat. <i>European Journal of Pharmacology</i> , 1992 , 219, 105-12	5.3	16
88	CCK antagonists and CCK-monoamine interactions in the control of satiety. <i>American Journal of Clinical Nutrition</i> , 1992 , 55, 291S-295S	7	36
87	L-683,877: Pharmacological profile of a novel 5-HT3 receptor antagonist. <i>Drug Development Research</i> , 1992 , 25, 17-28	5.1	5
86	Cholecystokinin receptors: synthetic antagonists with selectivity for receptor subtypes and possible clinical applications. <i>Biochemical Society Transactions</i> , 1991 , 19, 913-5	5.1	10
85	Evidence for an involvement of D1 and D2 dopamine receptors in mediating nicotine-induced hyperactivity in rats. <i>Psychopharmacology</i> , 1991 , 104, 343-50	4.7	85
84	Hypolocomotion induced by peripheral or central injection of CCK in the mouse is blocked by the CCKA receptor antagonist devazepide but not by the CCKB receptor antagonist L-365,260. <i>European Journal of Pharmacology</i> , 1991 , 193, 203-8	5.3	35
83	Fluoxetine reduces food intake by a cholecystokinin-independent mechanism. <i>Pharmacology Biochemistry and Behavior</i> , 1990 , 35, 51-4	3.9	27
82	Cholecystokinin and anxiety. <i>Trends in Pharmacological Sciences</i> , 1990 , 11, 271-3	13.2	140

81	The selective CCK-B receptor antagonist L-365,260 enhances morphine analgesia and prevents morphine tolerance in the rat. <i>European Journal of Pharmacology</i> , 1990 , 176, 35-44	5.3	220
80	Blockade of CCK-B receptors by L-365,260 induces analgesia in the squirrel monkey. <i>Brain Research</i> , 1990 , 534, 287-90	3.7	22
79	Pharmacological characterization of the behavioural syndrome induced by the NK-3 tachykinin agonist senktide in rodents: evidence for mediation by endogenous 5-HT. <i>Brain Research</i> , 1990 , 517, 111-6	3.7	39
78	Reversal of the anorectic effect of (+)-fenfluramine in the rat by the selective cholecystokinin receptor antagonist MK-329. <i>British Journal of Pharmacology</i> , 1990 , 99, 65-70	8.6	55
77	Multiple cholecystokinin (CCK) receptors and CCK-monoamine interactions are instrumental in the control of feeding. <i>Physiology and Behavior</i> , 1990 , 48, 849-57	3.5	54
76	The cholecystokinin receptor antagonist devazepide enhances morphine-induced analgesia but not morphine-induced respiratory depression in the squirrel monkey. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 1990 , 255, 1158-65	4.7	23
75	Postponement of satiety by blockade of brain cholecystokinin (CCK-B) receptors. <i>Science</i> , 1989 , 245, 1509-11	33.3	219
74	Evidence that blockade of post-synaptic 5-HT ₁ receptors elicits feeding in satiated rats. <i>Psychopharmacology</i> , 1989 , 97, 54-8	4.7	90
73	Chronic neuroleptic-induced mouth movements in the rat: suppression by CCK and selective dopamine D ₁ and D ₂ receptor antagonists. <i>Psychopharmacology</i> , 1989 , 98, 372-9	4.7	51
72	The role of 5-HT _{1B} receptors in the paraventricular nucleus of the hypothalamus in the control of feeding. <i>Neurobiology of Aging</i> , 1989 , 10, 209	5.6	2
71	MK-212 increases rat plasma ACTH concentration by activation of the 5-HT _{1C} receptor subtype. <i>Neuroscience Letters</i> , 1989 , 105, 174-6	3.3	50
70	The role of CCK caerulein, and CCK antagonists in nociception. <i>Pain</i> , 1989 , 39, 307-328	8	194
69	Blockade of apomorphine-induced yawning in rats by the dopamine autoreceptor antagonist (+)-AJ 76. <i>Neuropharmacology</i> , 1989 , 28, 1423-5	5.5	11
68	Morphine-induced analgesia in the rat paw pressure test is blocked by CCK and enhanced by the CCK antagonist MK-329. <i>Neuropharmacology</i> , 1989 , 28, 243-7	5.5	73
67	Evidence that decreased feeding induced by systemic injection of cholecystokinin is mediated by CCK-A receptors. <i>European Journal of Pharmacology</i> , 1989 , 173, 233-4	5.3	98
66	Effects of ageing on the behavioural responses to dopamine agonists: decreased yawning and locomotion, but increased stereotypy. <i>Brain Research</i> , 1989 , 495, 20-30	3.7	39
65	The 5-HT _{1A} agonist 8-OH-DPAT increases consumption of palatable wet mash and liquid diets in the rat. <i>Psychopharmacology</i> , 1988 , 94, 58-63	4.7	51
64	The NK-3 tachykinin agonist senktide elicits yawning and chewing mouth movements following subcutaneous administration in the rat. Evidence for cholinergic mediation. <i>Psychopharmacology</i> , 1988 , 95, 502-6	4.7	16

63	8-OH-DPAT elicits feeding and not chewing: evidence from liquid diet studies and a diet choice test. <i>Psychopharmacology</i> , 1988 , 95, 185-8	4.7	19
62	Role of dopamine D-1 and D-2 receptor subtypes in mediating dopamine agonist effects on food consumption in rats. <i>Psychopharmacology</i> , 1988 , 96, 370-4	4.7	58
61	Evidence that the hyperphagic response to 8-OH-DPAT is mediated by 5-HT _{1A} receptors. <i>European Journal of Pharmacology</i> , 1988 , 150, 361-6	5.3	98
60	Enhancement of morphine analgesia and prevention of morphine tolerance in the rat by the cholecystokinin antagonist L-364,718. <i>European Journal of Pharmacology</i> , 1988 , 147, 469-72	5.3	129
59	CCK-A receptors in the rat interpeduncular nucleus: evidence for a presynaptic location. <i>Brain Research</i> , 1988 , 454, 101-5	3.7	19
58	Antinociceptive effects of novel, selective tachykinin receptor antagonists in thermal and chemical analgesia tests. <i>Regulatory Peptides</i> , 1988 , 22, 58		10
57	The role of NK-3 and NK-4 receptors in the mediation of reciprocal hindlimb scratching induced by tachykinin receptor agonists. <i>Regulatory Peptides</i> , 1988 , 22, 59		3
56	Behavioural evidence for CNS tachykinin-cholinergic interactions. <i>Regulatory Peptides</i> , 1988 , 22, 178		3
55	The NK-3 tachykinin receptor agonist senktide elicits 5-HT-mediated behaviour following central or peripheral administration in mice and rats. <i>British Journal of Pharmacology</i> , 1988 , 94, 285-7	8.6	58
54	Relationship of increased food intake and plasma ACTH levels to 5-HT _{1A} receptor activation in rats. <i>Psychoneuroendocrinology</i> , 1988 , 13, 471-8	5	56
53	The 5-HT _{1A} agonists 8-OH-DPAT, buspirone and ipsapirone attenuate stress-induced anorexia in rats. <i>Journal of Psychopharmacology</i> , 1987 , 1, 23-30	4.6	13
52	Single administration of 5-HT _{1A} agonists decreases 5-HT _{1A} presynaptic, but not postsynaptic receptor-mediated responses: relationship to antidepressant-like action. <i>European Journal of Pharmacology</i> , 1987 , 138, 53-60	5.3	139
51	Antidepressant-like action of 5-HT _{1A} agonists and conventional antidepressants in an animal model of depression. <i>European Journal of Pharmacology</i> , 1987 , 134, 265-74	5.3	184
50	5-HT _{1B} agonists induce anorexia at a postsynaptic site. <i>European Journal of Pharmacology</i> , 1987 , 141, 429-35	5.3	145
49	Classification and function of CCK receptors. <i>Trends in Pharmacological Sciences</i> , 1987 , 8, 207-208	13.2	112
48	Senktide, a selective neurokinin B-like agonist, elicits serotonin-mediated behaviour following intracisternal administration in the mouse. <i>Neuroscience Letters</i> , 1987 , 80, 321-6	3.3	33
47	Apomorphine-induced yawning in rats is abolished by bilateral 6-hydroxydopamine lesions of the substantia nigra. <i>Psychopharmacology</i> , 1987 , 93, 336-42	4.7	36
46	Effects of the novel anxiolytics gepirone, buspirone and ipsapirone on free feeding and on feeding induced by 8-OH-DPAT. <i>Psychopharmacology</i> , 1987 , 93, 349-52	4.7	76

45	Effects of Drugs on Spontaneous Motor Activity 1987 , 153-211		
44	Para-chlorophenylalanine prevents feeding induced by the serotonin agonist 8-hydroxy-2-(di-n-propylamino) tetralin (8-OH-DPAT). <i>Psychopharmacology</i> , 1986 , 89, 467-71	4.7	101
43	Neurochemical and behavioural evidence for mediation of the hyperphagic action of 8-OH-DPAT by 5-HT cell body autoreceptors. <i>European Journal of Pharmacology</i> , 1986 , 129, 347-52	5.3	182
42	Monitoring 5HT metabolism in the brain of the freely moving rat. <i>Annals of the New York Academy of Sciences</i> , 1986 , 473, 321-36	6.5	12
41	8-OH-DPAT-induced hyperphagia: its neural basis and possible therapeutic relevance. <i>Appetite</i> , 1986 , 7 Suppl, 127-40	4.5	176
40	Putative anxiolytics 8-OH-DPAT, buspirone and TVX Q 7821 are agonists at 5-HT _{1A} autoreceptors in the raphae nuclei. <i>Trends in Pharmacological Sciences</i> , 1986 , 7, 212-214	13.2	108
39	Yawning elicited by systemic and intrastriatal injection of pibedil and apomorphine in the rat. <i>Psychopharmacology</i> , 1985 , 86, 175-81	4.7	51
38	Low doses of the putative serotonin agonist 8-hydroxy-2-(di-n-propylamino) tetralin (8-OH-DPAT) elicit feeding in the rat. <i>Psychopharmacology</i> , 1985 , 86, 197-204	4.7	283
37	Local application of beta-phenylethylamine to the caudate nucleus of the rat elicits locomotor stimulation. <i>Pharmacology Biochemistry and Behavior</i> , 1985 , 22, 159-62	3.9	15
36	Characteristics of feeding induced by the serotonin agonist 8-hydroxy-2-(di-n-propylamino) tetralin (8-OH-DPAT). <i>Brain Research Bulletin</i> , 1985 , 15, 377-84	3.9	141
35	Pharmacological and biochemical analysis of feeding behaviour. <i>Brain Research Bulletin</i> , 1985 , 15, 369-70	3.9	1
34	Bilateral lesions of the striatum induced with 6-hydroxydopamine abolish apomorphine-induced yawning in rats. <i>Neuropharmacology</i> , 1985 , 24, 1051-5	5.5	38
33	Behavioural evidence for the existence of dopamine autoreceptors. <i>Trends in Pharmacological Sciences</i> , 1985 , 6, 17-18	13.2	16
32	Hypodipsia, stereotypy and hyperactivity induced by beta-phenylethylamine in the water-deprived rat. <i>Pharmacology Biochemistry and Behavior</i> , 1984 , 20, 1-7	3.9	13
31	Differential aversive stimulus properties of beta-phenylethylamine and of d-amphetamine. <i>Psychopharmacology</i> , 1984 , 82, 189-93	4.7	18
30	Environmental experience produces qualitative changes in the stimulant effects of beta-phenylethylamine in rats. <i>Psychopharmacology</i> , 1984 , 84, 132-5	4.7	12
29	Effects of pibedil and of d-amphetamine on locomotor activity in immature rats. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 1984 , 8, 249-55	5.5	1
28	Potentiation of total horizontal activity and ambulation in rats treated with combinations of beta-phenylethylamine and naloxone. <i>Neuropharmacology</i> , 1984 , 23, 1059-64	5.5	7

27	Locomotor Stimulant Effect of β -Phenylethylamine-Naloxone Mixtures 1984 , 449-454		1
26	Taste-Dependent Effects of Low Doses of β -Phenylethylamine and d-Amphetamine on Drinking in the Rat 1984 , 435-440		1
25	Studies on the Mechanism of Action of β -Phenylethylamine Stereotypy in Rodents: Implications for a β -Phenylethylamine Animal Model of Schizophrenia 1984 , 389-411		7
24	Effects of Tryptamine and 5-Hydroxytryptamine on Food Intake in the Rat 1984 , 415-421		1
23	β -Phenylethylamine And d-Amphetamine: Differential Potency in the Conditioned Taste Aversion Paradigm 1984 , 441-447		2
22	Potentiation of the behavioural effects of the antidepressant phenelzine by deuterium substitution. <i>Psychopharmacology</i> , 1983 , 81, 122-5	4-7	25
21	Pharmacology of beta-phenylethylamine-induced seizures in mice. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 1983 , 7, 787-90	5-5	7
20	Piribedil: behavioural, neurochemical and clinical profile of a dopamine agonist. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 1983 , 7, 3-27	5-5	30
19	Dopaminergic involvement in the control of drinking behaviour: a brief review. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 1983 , 7, 487-93	5-5	45
18	Deuterium substitution enhances the effects of beta-phenylethylamine on spontaneous motor activity in the rat. <i>Pharmacology Biochemistry and Behavior</i> , 1983 , 19, 471-5	3-9	31
17	Dopamine agonist-induced restoration of drinking in response to hypertonic saline in adipsic dopamine denervated rats. <i>Brain Research Bulletin</i> , 1982 , 8, 375-9	3-9	19
16	Behavioural and neurochemical effects of meta-tyrosine in the rat. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 1982 , 6, 389-94	5-5	1
15	Biphasic behavioural stimulation induced by a monoamine oxidase-inhibiting antidepressant. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 1982 , 6, 383-8	5-5	7
14	An observational analysis of the behavioural effects of beta-phenylethylamine in isolated and grouped mice. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 1982 , 6, 143-58	5-5	18
13	Putative PEA receptors?. <i>Psychiatry Research</i> , 1982 , 7, 387-8	9-9	2
12	Variation in response to stimulation of central 5-hydroxytryptamine mechanisms in two strains of albino rat. <i>Brain Research</i> , 1982 , 247, 172-6	3-7	10
11	Suppression of drinking and induction of sedation by a dopamine agonist is blocked by small doses of spiperone. <i>Neuropharmacology</i> , 1982 , 21, 69-72	5-5	9
10	The role of catecholamines, 5-hydroxytryptamine and m-tyramine in the behavioural effects of m-tyrosine in the rat. <i>European Journal of Pharmacology</i> , 1982 , 84, 139-49	5-3	12

9	A pharmacological analysis of the hyperactivity syndrome induced by beta-phenylethylamine in the mouse. <i>British Journal of Pharmacology</i> , 1982 , 77, 129-39	8.6	44
8	Alterations in trace amine and trace acid concentrations in isolated aggressive mice. <i>Pharmacology Biochemistry and Behavior</i> , 1982 , 17, 1291-4	3.9	14
7	Behavioural effects of acute and chronic beta-phenylethylamine administration in the rat: evidence for the involvement of 5-hydroxytryptamine. <i>Neuropharmacology</i> , 1981 , 20, 1067-72	5.5	48
6	Single or repeated administration of small doses of apomorphine on water intake and activity in water-deprived rats. <i>Neuropharmacology</i> , 1981 , 20, 257-60	5.5	13
5	The effects of acute and chronic administration of beta-phenylethylamine on food intake and body weight in rats. <i>Progress in Neuro-Psychopharmacology & Biological Psychiatry</i> , 1981 , 5, 411-4		24
4	Effects of acute or chronic administration of low doses of a dopamine agonist on drinking and locomotor activity in the rat. <i>Psychopharmacology</i> , 1981 , 72, 197-202	4.7	19
3	Sedation, stereotypy and the inhibition of drinking after administration of ET 495 in the rat [proceedings]. <i>British Journal of Pharmacology</i> , 1979 , 66, 481P	8.6	1
2	Effects of ET 495 (piribedil) on water intake in the rat: evidence for a dopaminergic involvement in thirst. <i>Irish Journal of Medical Science</i> , 1978 , 147 Suppl 1, 19-23	1.9	6
1	Experimental Psychopharmacology		5