

Pinar Bor

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/402708/publications.pdf>

Version: 2024-02-01

26
papers

415
citations

840776

11
h-index

752698

20
g-index

34
all docs

34
docs citations

34
times ranked

588
citing authors

#	ARTICLE	IF	CITATIONS
1	A force-measuring device combined with ultrasound-based elastography for assessment of the uterine cervix. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2022, 101, 241-247.	2.8	5
2	Cervical intraepithelial neoplasia in women with transformation zone type 3: cervical biopsy versus large loop excision. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2022, 129, 2132-2140.	2.3	10
3	Vitamin D insufficiency among Danish pregnant women—Prevalence and association with adverse obstetric outcomes and placental vitamin D metabolism. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2021, 100, 480-488.	2.8	9
4	Continued versus discontinued oxytocin stimulation in the active phase of labour (CONDISOX): double blind randomised controlled trial. <i>BMJ, The</i> , 2021, 373, n716.	6.0	10
5	Cervical cancer prevention among older women—challenges in screening, diagnostic workup and treatment. <i>Acta Obstetrica Et Gynecologica Scandinavica</i> , 2021, 100, 1364-1368.	2.8	8
6	The Childbirth Experience Questionnaire (CEQ)—Validation of its use in a Danish-speaking population of new mothers stimulated with oxytocin during labour. <i>PLoS ONE</i> , 2020, 15, e0233122.	2.5	17
7	Use of stimulants, over-the-counter and prescription drugs among Danish pregnant women. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2020, 127, 205-210.	2.5	6
8	Validity and reliability of State-Trait Anxiety Inventory in Danish women aged 45 years and older with abnormal cervical screening results. <i>BMC Medical Research Methodology</i> , 2020, 20, 89.	3.1	32
9	CONDISOX- continued versus discontinued oxytocin stimulation of induced labour in a double-blind randomised controlled trial. <i>BMC Pregnancy and Childbirth</i> , 2019, 19, 320.	2.4	1
10	Use of alternative medicine, ginger and licorice among Danish pregnant women—a prospective cohort study. <i>BMC Complementary and Alternative Medicine</i> , 2019, 19, 5.	3.7	16
11	Prevalence of xenobiotic substances in first-trimester blood samples from Danish pregnant women: a cross-sectional study. <i>BMJ Open</i> , 2018, 8, e018390.	1.9	9
12	Discontinuation of intravenous oxytocin in the active phase of induced labour. <i>The Cochrane Library</i> , 2018, 2018, CD012274.	2.8	20
13	Discontinuing Oxytocin Infusion in the Active Phase of Labor. <i>Obstetrics and Gynecology</i> , 2017, 130, 1090-1096.	2.4	50
14	Continuation versus discontinuation of oxytocin infusion during the active phase of labour: a randomised controlled trial. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , 2016, 123, 129-135.	2.3	38
15	Discontinuation of intravenous oxytocin in the active phase of induced labour. <i>The Cochrane Library</i> , 2016, , .	2.8	8
16	Life-threatening necrotizing myometritis, due to Group A streptococcus—still a life-threatening condition. <i>Clinical Case Reports (discontinued)</i> , 2015, 3, 291-293.	0.5	4
17	Comparison of two different methods for measuring anti-mullerian hormone in a clinical series. <i>Reproductive Biology and Endocrinology</i> , 2015, 13, 107.	3.3	26
18	Teenage Pregnancies: Obstetric and Neonatal Outcomes at a Danish Regional Hospital. <i>Journal of Women's Health Care</i> , 2015, 04, .	0.2	1

#	ARTICLE	IF	CITATIONS
19	A rare case with multiple true knots together with single artery and four umbilical cord nuchal loops. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2013, 168, 117-118.	1.1	1
20	Predictive value of serum human chorionic gonadotropin ratio, progesterone and inhibin A for expectant management of early pregnancies of unknown location. <i>European Journal of Obstetrics, Gynecology and Reproductive Biology</i> , 2012, 165, 66-69.	1.1	9
21	Collagen concentration and biomechanical properties of samples from the lower uterine cervix in relation to age and parity in non-pregnant women. <i>Reproductive Biology and Endocrinology</i> , 2010, 8, 82.	3.3	48
22	Screening for Y microdeletions in men with testicular cancer and undescended testis. <i>Journal of Assisted Reproduction and Genetics</i> , 2006, 23, 41-45.	2.5	11
23	A new approach for screening for Y microdeletions: capillary electrophoresis combined with fluorescent multiplex PCR. <i>Journal of Assisted Reproduction and Genetics</i> , 2003, 20, 46-51.	2.5	10
24	Y-chromosome microdeletions and cytogenetic findings in unselected ICSI candidates at a Danish fertility clinic. <i>Journal of Assisted Reproduction and Genetics</i> , 2002, 19, 224-231.	2.5	25
25	Multiplex PCR for screening of microdeletions on the Y chromosome. <i>Journal of Assisted Reproduction and Genetics</i> , 2001, 18, 291-298.	2.5	22
26	Serum fructosamine and fructosamine - albumin ratio as screening tests for gestational diabetes mellitus. <i>Archives of Gynecology and Obstetrics</i> , 1999, 262, 105-111.	1.7	16