
Joep J Van Oosterhout

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4043263/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Antifungal Combinations for Treatment of Cryptococcal Meningitis in Africa. New England Journal of Medicine, 2018, 378, 1004-1017.	27.0	296
2	Adjunctive Dexamethasone in HIV-Associated Cryptococcal Meningitis. New England Journal of Medicine, 2016, 374, 542-554.	27.0	257
3	Effect of Optional Home Initiation of HIV Care Following HIV Self-testing on Antiretroviral Therapy Initiation Among Adults in Malawi. JAMA - Journal of the American Medical Association, 2014, 312, 372.	7.4	164
4	Rapid urine-based screening for tuberculosis in HIV-positive patients admitted to hospital in Africa (STAMP): a pragmatic, multicentre, parallel-group, double-blind, randomised controlled trial. Lancet, The, 2018, 392, 292-301.	13.7	156
5	Retention in care during the first 3 years of antiretroviral therapy for women in Malawi's option B+ programme: an observational cohort study. Lancet HIV,the, 2016, 3, e175-e182.	4.7	154
6	Evaluation of antiretroviral therapy results in a resource-poor setting in Blantyre, Malawi. Tropical Medicine and International Health, 2005, 10, 464-470.	2.3	125
7	Adherence to Antiretroviral Therapy During and After Pregnancy: Cohort Study on Women Receiving Care in Malawi's Option B+ Program. Clinical Infectious Diseases, 2016, 63, ciw500.	5.8	123
8	Leave no one behind: response to new evidence and guidelines for the management of cryptococcal meningitis in low-income and middle-income countries. Lancet Infectious Diseases, The, 2019, 19, e143-e147.	9.1	63
9	A Prospective Longitudinal Study of the Clinical Outcomes from Cryptococcal Meningitis following Treatment Induction with 800 mg Oral Fluconazole in Blantyre, Malawi. PLoS ONE, 2013, 8, e67311.	2.5	62
10	HIV Associated Neurocognitive Disorders (HAND) in Malawian Adults and Effect on Adherence to Combination Anti-Retroviral Therapy: A Cross Sectional Study. PLoS ONE, 2014, 9, e98962.	2.5	52
11	Virological failure, HIV-1 drug resistance, and early mortality in adults admitted to hospital in Malawi: an observational cohort study. Lancet HIV,the, 2020, 7, e620-e628.	4.7	46
12	Genome-wide association study of nevirapine hypersensitivity in a sub-Saharan African HIV-infected population. Journal of Antimicrobial Chemotherapy, 2017, 72, dkw545.	3.0	42
13	Clinical Characteristics and Lung Function in Older Children Vertically Infected With Human Immunodeficiency Virus in Malawi. Journal of the Pediatric Infectious Diseases Society, 2016, 5, 161-169.	1.3	41
14	National estimates and risk factors associated with early mother-to-child transmission of HIV after implementation of option B+: a cross-sectional analysis. Lancet HIV,the, 2018, 5, e688-e695.	4.7	40
15	CYP2B6 c.983T>C polymorphism is associated with nevirapine hypersensitivity in Malawian and Ugandan HIV populations. Journal of Antimicrobial Chemotherapy, 2014, 69, 3329-3334.	3.0	36
16	The benefit of supplementary feeding for wasted Malawian adults initiating ART. AIDS Care - Psychological and Socio-Medical Aspects of AIDS/HIV, 2010, 22, 737-742.	1.2	33
17	Lopinavir plus nucleoside reverse-transcriptase inhibitors, lopinavir plus raltegravir, or lopinavir monotherapy for second-line treatment of HIV (EARNEST): 144-week follow-up results from a randomised controlled trial. Lancet Infectious Diseases, The, 2018, 18, 47-57.	9.1	32
18	Cost-effectiveness of urine-based tuberculosis screening in hospitalised patients with HIV in Africa: a microsimulation modelling study. The Lancet Global Health, 2019, 7, e200-e208.	6.3	32

#	Article	IF	CITATIONS
19	A Reduction in Adult Blood Stream Infection and Case Fatality at a Large African Hospital following Antiretroviral Therapy Roll-Out. PLoS ONE, 2014, 9, e92226.	2.5	30
20	Hepatitis B Virus Sub-genotype A1 Infection Is Characterized by High Replication Levels and Rapid Emergence of Drug Resistance in HIV-Positive Adults Receiving First-line Antiretroviral Therapy in Malawi. Clinical Infectious Diseases, 2014, 59, 1618-1626.	5.8	29
21	Systematic Review of Hypertension and Diabetes Burden, Risk Factors, and Interventions for Prevention and Control in Malawi: The NCD BRITE Consortium. Global Heart, 2019, 14, 109.	2.3	23
22	One-year Mortality Outcomes From the Advancing Cryptococcal Meningitis Treatment for Africa Trial of Cryptococcal Meningitis Treatment in Malawi. Clinical Infectious Diseases, 2020, 70, 521-524.	5.8	13
23	A targeted approach for routine viral load monitoring in Malawian adults on antiretroviral therapy. Tropical Medicine and International Health, 2018, 23, 526-532.	2.3	12
24	Impact of inter-partner HIV disclosure patterns in Malawi's PMTCT program: A mixed-method study. PLoS ONE, 2019, 14, e0219967.	2.5	10
25	Short-term Mortality Outcomes of HIV-Associated Cryptococcal Meningitis in Antiretroviral Therapy–NaÃ⁻ve and –Experienced Patients in Sub-Saharan Africa. Open Forum Infectious Diseases, 2021, 8, ofab397.	0.9	9
26	Challenges with targeted viral load testing for medical inpatients at Queen Elizabeth Central Hospital in Blantyre, Malawi. Malawi Medical Journal, 2017, 28, 179.	0.6	8
27	The Malawi NCD BRITE Consortium: Building Research Capacity, Implementation, and Translation Expertise for Noncommunicable Diseases. Global Heart, 2020, 14, 149.	2.3	7
28	Pregnancy intention and contraceptive use among HIV-positive Malawian women at 4-26 weeks post-partum: A nested cross-sectional study. PLoS ONE, 2019, 14, e0215947.	2.5	5
29	High mortality in subjects with both diabetes and HIV in sub-Saharan Africa. Aids, 2018, 32, 2083-2084.	2.2	4