

Young-Kwon Park

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4058776/publications.pdf>

Version: 2024-02-01

15
papers

2,014
citations

687363

13
h-index

1058476

14
g-index

19
all docs

19
docs citations

19
times ranked

5269
citing authors

#	ARTICLE	IF	CITATIONS
1	Interplay of BAF and MLL4 promotes cell type-specific enhancer activation. <i>Nature Communications</i> , 2021, 12, 1630.	12.8	38
2	MED1 is a lipogenesis coactivator required for postnatal adipose expansion. <i>Genes and Development</i> , 2021, 35, 713-728.	5.9	9
3	H3.3K4M destabilizes enhancer H3K4 methyltransferases MLL3/MLL4 and impairs adipose tissue development. <i>Nucleic Acids Research</i> , 2019, 47, 607-620.	14.5	1,326
4	Depletion of Nsd2-mediated histone H3K36 methylation impairs adipose tissue development and function. <i>Nature Communications</i> , 2018, 9, 1796.	12.8	58
5	Glucocorticoid Receptor Accelerates, but Is Dispensable for, Adipogenesis. <i>Molecular and Cellular Biology</i> , 2017, 37, .	2.3	69
6	Distinct Roles of Transcription Factors KLF4, Krox20, and Peroxisome Proliferator-Activated Receptor α in Adipogenesis. <i>Molecular and Cellular Biology</i> , 2017, 37, .	2.3	44
7	Brd4 binds to active enhancers to control cell identity gene induction in adipogenesis and myogenesis. <i>Nature Communications</i> , 2017, 8, 2217.	12.8	161
8	HIF-1-Dependent Induction of Jumonji Domain-Containing Protein (JMJD) 3 under Hypoxic Conditions. <i>Molecules and Cells</i> , 2014, 37, 43-50.	2.6	74
9	Hypoxia-inducible Factor-2 α -dependent Hypoxic Induction of Wnt10b Expression in Adipogenic Cells. <i>Journal of Biological Chemistry</i> , 2013, 288, 26311-26322.	3.4	33
10	Differentiated Embryo Chondrocyte 1 (DEC1) Represses PPAR α Gene through Interacting with CCAAT/Enhancer Binding Protein β (C/EBP β). <i>Molecules and Cells</i> , 2012, 33, 575-582.	2.6	29
11	Estrogen receptor beta inhibits transcriptional activity of hypoxia inducible factor-1 through the downregulation of arylhydrocarbon receptor nuclear translocator. <i>Breast Cancer Research</i> , 2011, 13, R32.	5.0	34
12	Prevention of CCAAT/enhancer-binding protein β DNA binding by hypoxia during adipogenesis.. <i>Journal of Biological Chemistry</i> , 2011, 286, 41904.	3.4	0
13	Prevention of CCAAT/Enhancer-binding Protein β DNA Binding by Hypoxia during Adipogenesis. <i>Journal of Biological Chemistry</i> , 2010, 285, 3289-3299.	3.4	27
14	Nitric Oxide Donor, (A \pm)-S-Nitroso-N-acetylpenicillamine, Stabilizes Transactive Hypoxia-Inducible Factor-1 α by Inhibiting von Hippel-Lindau Recruitment and Asparagine Hydroxylation. <i>Molecular Pharmacology</i> , 2008, 74, 236-245.	2.3	48
15	Clioquinol, a Cu(II)/Zn(II) Chelator, Inhibits Both Ubiquitination and Asparagine Hydroxylation of Hypoxia-inducible Factor-1 α , Leading to Expression of Vascular Endothelial Growth Factor and Erythropoietin in Normoxic Cells. <i>Journal of Biological Chemistry</i> , 2006, 281, 34056-34063.	3.4	58