

Clemens C C Bauer

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4062457/publications.pdf>

Version: 2024-02-01

16
papers

626
citations

840119

11
h-index

940134

16
g-index

19
all docs

19
docs citations

19
times ranked

1202
citing authors

#	ARTICLE	IF	CITATIONS
1	Optimizing real time fMRI neurofeedback for therapeutic discovery and development. <i>NeuroImage: Clinical</i> , 2014, 5, 245-255.	1.4	179
2	Child overweight and obesity are associated with reduced executive cognitive performance and brain alterations: a magnetic resonance imaging study in Mexican children. <i>Pediatric Obesity</i> , 2015, 10, 196-204.	1.4	78
3	Source-space EEG neurofeedback links subjective experience with brain activity during effortless awareness meditation. <i>NeuroImage</i> , 2017, 151, 117-127.	2.1	57
4	Mindfulness training reduces stress and amygdala reactivity to fearful faces in middle-school children.. <i>Behavioral Neuroscience</i> , 2019, 133, 569-585.	0.6	55
5	Mindfulness training preserves sustained attention and resting state anticorrelation between default mode network and dorsolateral prefrontal cortex: A randomized controlled trial. <i>Human Brain Mapping</i> , 2020, 41, 5356-5369.	1.9	43
6	From State-to-Trait Meditation: Reconfiguration of Central Executive and Default Mode Networks. <i>ENeuro</i> , 2019, 6, ENEURO.0335-18.2019.	0.9	42
7	Sustained attention to spontaneous thumb sensations activates brain somatosensory and other proprioceptive areas. <i>Brain and Cognition</i> , 2014, 87, 86-96.	0.8	38
8	The real-time fMRI neurofeedback based stratification of Default Network Regulation Neuroimaging data repository. <i>NeuroImage</i> , 2017, 146, 157-170.	2.1	37
9	Real-time fMRI neurofeedback reduces auditory hallucinations and modulates resting state connectivity of involved brain regions: Part 2: Default mode network -preliminary evidence. <i>Psychiatry Research</i> , 2020, 284, 112770.	1.7	34
10	Subjective Somatosensory Experiences Disclosed by Focused Attention: Cortical-Hippocampal-Insular and Amygdala Contributions. <i>PLoS ONE</i> , 2014, 9, e104721.	1.1	21
11	Image acquisition and quality assurance in the Boston Adolescent Neuroimaging of Depression and Anxiety study. <i>NeuroImage: Clinical</i> , 2020, 26, 102242.	1.4	13
12	Cortical plasticity in phantom limb pain: A fMRI study on the neural correlates of behavioral clinical manifestations.. <i>Psychiatry Research - Neuroimaging</i> , 2020, 304, 111151.	0.9	11
13	Abnormal Functioning of the Thalamocortical System Underlies the Conscious Awareness of the Phantom Limb Phenomenon. <i>Neuroradiology Journal</i> , 2010, 23, 671-679.	0.6	9
14	The Time Varying Networks of the Interoceptive Attention and Rest. <i>ENeuro</i> , 2021, 8, ENEURO.0341-20.2021.	0.9	3
15	Altered working memory-related brain activity in children at familial high risk for psychosis: A preliminary study. <i>Schizophrenia Research</i> , 2022, 240, 186-192.	1.1	3
16	Hyperactivation of Posterior Default Mode Network During Self-Referential Processing in Children at Familial High-Risk for Psychosis. <i>Frontiers in Psychiatry</i> , 2021, 12, 613142.	1.3	2