

# Hui-Hui Lin

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4078033/publications.pdf>

Version: 2024-02-01

18  
papers

277  
citations

1039880

9  
h-index

887953

17  
g-index

19  
all docs

19  
docs citations

19  
times ranked

489  
citing authors

#	ARTICLE	IF	CITATIONS
1	Increase sensitivity in detecting superficial, low grade bladder cancer by combination analysis of hypermethylation of E-cadherin, p16, p14, RASSF1A genes in urine. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2010, 28, 597-602.	0.8	61
2	MiR-193b Mediates CEBPD-Induced Cisplatin Sensitization Through Targeting ETS1 and Cyclin D1 in Human Urothelial Carcinoma Cells. <i>Journal of Cellular Biochemistry</i> , 2017, 118, 1563-1573.	1.2	44
3	Genetic Variations in Glutathione Pathway Genes Predict Cancer Recurrence in Patients Treated with Transurethral Resection and Bacillus Calmette-Guérin Instillation for Non-muscle Invasive Bladder Cancer. <i>Annals of Surgical Oncology</i> , 2015, 22, 4104-4110.	0.7	31
4	Hypermethylation of E-cadherin, p16, p14, and RASSF1A genes in pathologically normal urothelium predict bladder recurrence of bladder cancer after transurethral resection. <i>Urologic Oncology: Seminars and Original Investigations</i> , 2012, 30, 177-181.	0.8	25
5	Hypoxia-regulated MicroRNA-210 Overexpression is Associated with Tumor Development and Progression in Upper Tract Urothelial Carcinoma. <i>International Journal of Medical Sciences</i> , 2017, 14, 578-584.	1.1	22
6	MicroRNA-145 suppresses cell migration and invasion in upper tract urothelial carcinoma by targeting ARF6. <i>FASEB Journal</i> , 2020, 34, 5975-5992.	0.2	21
7	Potential Role of CCND1 G870A Genotype as a Predictor for Urothelial Carcinoma Susceptibility and Muscle-Invasiveness in Taiwan. <i>Chinese Journal of Physiology</i> , 2011, 54, 196-202.	0.4	14
8	Prognostic Value of Leptin Receptor Overexpression in Upper Tract Urothelial Carcinomas in Taiwan. <i>Clinical Genitourinary Cancer</i> , 2017, 15, e653-e659.	0.9	11
9	CSF-1 Overexpression Predicts Poor Prognosis in Upper Tract Urothelial Carcinomas. <i>Disease Markers</i> , 2019, 2019, 1-9.	0.6	9
10	High Transaldolase 1 expression predicts poor survival of patients with upper tract urothelial carcinoma. <i>Pathology International</i> , 2021, 71, 463-470.	0.6	8
11	Over-expression of Activated Signal Transducer and Activator of Transcription 3 Predicts Poor Prognosis in Upper Tract Urothelial Carcinoma. <i>International Journal of Medical Sciences</i> , 2017, 14, 1360-1367.	1.1	7
12	Overexpression of PTP4A3 is associated with metastasis and unfavorable prognosis in bladder cancer. <i>World Journal of Urology</i> , 2016, 34, 835-846.	1.2	6
13	High visfatin expression predicts poor prognosis of upper tract urothelial carcinoma patients. <i>American Journal of Cancer Research</i> , 2015, 5, 2447-54.	1.4	6
14	CCND1 1722 polymorphism and potential relevance to upper tract urothelial cancer. <i>Anticancer Research</i> , 2011, 31, 1043-7.	0.5	5
15	The prognostic value of CSN6 expression in upper tract urothelial carcinomas. <i>Kaohsiung Journal of Medical Sciences</i> , 2019, 35, 559-565.	0.8	2
16	MicroRNA-375-3p Suppresses Upper Tract Urothelial Carcinoma Cell Migration and Invasion via Targeting Derlin-1. <i>Cancers</i> , 2022, 14, 880.	1.7	2
17	MicroRNA-145-5p modulates Krüppel-like factor 5 and inhibits cell proliferation, migration, and invasion in nasopharyngeal carcinoma. <i>BMC Molecular and Cell Biology</i> , 2022, 23, .	1.0	2
18	High Ubiquitin-Specific Protease 2a Expression Level Predicts Poor Prognosis in Upper Tract Urothelial Carcinoma. <i>Applied Immunohistochemistry and Molecular Morphology</i> , 2022, 30, 304-310.	0.6	1