

István Kovanecz

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4078128/publications.pdf>

Version: 2024-02-01

13
papers

648
citations

840119

11
h-index

1125271

13
g-index

13
all docs

13
docs citations

13
times ranked

558
citing authors

#	ARTICLE	IF	CITATIONS
1	Vardenafil prevents fibrosis and loss of corporal smooth muscle that occurs after bilateral cavernosal nerve resection in the rat. <i>Urology</i> , 2006, 68, 429-435.	0.5	198
2	Chronic daily tadalafil prevents the corporal fibrosis and veno-occlusive dysfunction that occurs after cavernosal nerve resection. <i>BJU International</i> , 2007, 101, 070921231855003-???	1.3	137
3	Fibrosis and Loss of Smooth Muscle in the Corpora Cavernosa Precede Corporal Veno-Occlusive Dysfunction (CVOD) Induced by Experimental Cavernosal Nerve Damage in the Rat. <i>Journal of Sexual Medicine</i> , 2009, 6, 415-428.	0.3	100
4	Early onset of fibrosis within the arterial media in a rat model of type 2 diabetes mellitus with erectile dysfunction. <i>BJU International</i> , 2009, 103, 1396-1404.	1.3	46
5	Sildenafil Promotes Smooth Muscle Preservation and Ameliorates Fibrosis Through Modulation of Extracellular Matrix and Tissue Growth Factor Gene Expression After Bilateral Cavernosal Nerve Resection in the Rat. <i>Journal of Sexual Medicine</i> , 2011, 8, 1048-1060.	0.3	40
6	Sildenafil Attenuates Inflammation and Oxidative Stress in Pelvic Ganglia Neurons after Bilateral Cavernosal Nerve Damage. <i>International Journal of Molecular Sciences</i> , 2014, 15, 17204-17220.	1.8	34
7	Sildenafil promotes neuroprotection of the pelvic ganglia neurones after bilateral cavernosal nerve resection in the rat. <i>BJU International</i> , 2013, 111, 159-170.	1.3	22
8	Myostatin genetic inactivation inhibits myogenesis by muscle-derived stem cells in vitro but not when implanted in the mdx mouse muscle. <i>Stem Cell Research and Therapy</i> , 2013, 4, 4.	2.4	21
9	Implanted Muscle-Derived Stem Cells Ameliorate Erectile Dysfunction in a Rat Model of Type 2 Diabetes, but Their Repair Capacity Is Impaired by Their Prior Exposure to the Diabetic Milieu. <i>Journal of Sexual Medicine</i> , 2016, 13, 786-797.	0.3	20
10	Chronic High Dose Intraperitoneal Bisphenol A (BPA) Induces Substantial Histological and Gene Expression Alterations in Rat Penile Tissue Without Impairing Erectile Function. <i>Journal of Sexual Medicine</i> , 2013, 10, 2952-2966.	0.3	12
11	Dyslipidemia Is a Major Factor in Stem Cell Damage Induced by Uncontrolled Long-Term Type 2 Diabetes and Obesity in the Rat, as Suggested by the Effects on Stem Cell Culture. <i>Journal of Sexual Medicine</i> , 2018, 15, 1678-1697.	0.3	12
12	Stem Cells from a Female Rat Model of Type 2 Diabetes/Obesity and Stress Urinary Incontinence Are Damaged by In Vitro Exposure to its Dyslipidemic Serum, Predicting Inadequate Repair Capacity In Vivo. <i>International Journal of Molecular Sciences</i> , 2019, 20, 4044.	1.8	5
13	Evaluation of the In Vitro Damage Caused by Lipid Factors on Stem Cells from a Female Rat Model of Type 2 Diabetes/Obesity and Stress Urinary Incontinence. <i>International Journal of Molecular Sciences</i> , 2020, 21, 5045.	1.8	1