

Fumitaka Tanaka

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4081044/publications.pdf>

Version: 2024-02-01

24
papers

247
citations

840776

11
h-index

1058476

14
g-index

24
all docs

24
docs citations

24
times ranked

476
citing authors

#	ARTICLE	IF	CITATIONS
1	Prehypertension Subtype With Elevated C-Reactive Protein: Risk of Ischemic Stroke in a General Japanese Population. <i>American Journal of Hypertension</i> , 2010, 23, 1108-1113.	2.0	24
2	Relationship between the seismic scale of the 2011 northeast Japan earthquake and the incidence of acute myocardial infarction: A population-based study. <i>American Heart Journal</i> , 2015, 169, 861-869.	2.7	21
3	Long-Term Effects of the 2011 Japan Earthquake and Tsunami on Incidence of Fatal and Nonfatal Myocardial Infarction. <i>American Journal of Cardiology</i> , 2017, 120, 352-358.	1.6	18
4	Impact of the Japan earthquake disaster with massive Tsunami on emergency coronary intervention and in-hospital mortality in patients with acute ST-elevation myocardial infarction. <i>European Heart Journal: Acute Cardiovascular Care</i> , 2014, 3, 195-203.	1.0	17
5	Predictive Value of Lipoprotein Indices for Residual Risk of Acute Myocardial Infarction and Sudden Death in Men With Low-Density Lipoprotein Cholesterol Levels <120 mg/dl. <i>American Journal of Cardiology</i> , 2013, 112, 1063-1068.	1.6	13
6	Relative and absolute risks of all-cause and cause-specific deaths attributable to atrial fibrillation in middle-aged and elderly community dwellers. <i>International Journal of Cardiology</i> , 2015, 184, 692-698.	1.7	13
7	Ability of B-Type Natriuretic Peptide Testing to Predict Cardioembolic Stroke in the General Population—Comparisons With C-Reactive Protein and Urinary Albumin. <i>Circulation Journal</i> , 2018, 82, 1017-1025.	1.6	13
8	Sustained Increase in the Incidence of Acute Decompensated Heart Failure After the 2011 Japan Earthquake and Tsunami. <i>American Journal of Cardiology</i> , 2016, 118, 1374-1379.	1.6	12
9	Prognostic Value of Electrocardiographic Left Ventricular Hypertrophy on Cardiovascular Risk in a Non-Hypertensive Community-Based Population. <i>American Journal of Hypertension</i> , 2018, 31, 895-901.	2.0	12
10	Poor self-rated health predicts the incidence of functional disability in elderly community dwellers in Japan: a prospective cohort study. <i>BMC Geriatrics</i> , 2020, 20, 328.	2.7	12
11	Comparison between urine albumin-to-creatinine ratio and urine protein dipstick testing for prevalence and ability to predict the risk for chronic kidney disease in the general population (Iwate-KENCO study): a prospective community-based cohort study. <i>BMC Nephrology</i> , 2016, 17, 46.	1.8	11
12	Risk of stroke and heart failure attributable to atrial fibrillation in middle-aged and elderly people: Results from a five-year prospective cohort study of Japanese community dwellers. <i>Journal of Epidemiology</i> , 2017, 27, 360-367.	2.4	11
13	Temporal Trends in the Incidence and Clinical Features of Acute Myocardial Infarction in a Japanese Rural Area From 2006 to 2014. <i>Circulation Journal</i> , 2017, 81, 1854-1861.	1.6	11
14	Association between high-sensitivity cardiac troponin T and future cardiovascular incidence in a general Japanese population: results from the Tohoku medical megabank project. <i>Biomarkers</i> , 2019, 24, 566-573.	1.9	10
15	Bradycardia is associated with future cardiovascular diseases and death in men from the general population. <i>Atherosclerosis</i> , 2014, 236, 116-120.	0.8	9
16	Burden of high blood pressure as a contributing factor to stroke in the Japanese community-based diabetic population. <i>Hypertension Research</i> , 2018, 41, 531-538.	2.7	9
17	Plasma Xanthine Oxidoreductase Activity Is Associated with a High Risk of Cardiovascular Disease in a General Japanese Population. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 1894.	2.6	9
18	The urine albumin-creatinine ratio is a predictor for incident long-term care in a general population. <i>PLoS ONE</i> , 2018, 13, e0195013.	2.5	8

#	ARTICLE	IF	CITATIONS
19	Cardiovascular Risk Stratification With Plasma B-Type Natriuretic Peptide Levels in a Community-Based Hypertensive Cohort. <i>American Journal of Cardiology</i> , 2014, 113, 682-686.	1.6	5
20	The Value of a Cystatin C-based Estimated Glomerular Filtration Rate for Cardiovascular Assessment in a General Japanese Population: Results From the Iwate Tohoku Medical Megabank Project. <i>Journal of Epidemiology</i> , 2020, 30, 260-267.	2.4	5
21	Sex-specific temporal trends in the incidence and prevalence of hospitalized patients with preserved versus reduced left ventricular ejection fraction heart failure: A Japanese community-wide study. <i>IJC Heart and Vasculature</i> , 2015, 9, 15-21.	1.1	2
22	Association between Milk Intake and Incident Stroke among Japanese Community Dwellers: The Iwate-KENCO Study. <i>Nutrients</i> , 2021, 13, 3781.	4.1	2
23	Usefulness of risk grading system using albuminuria for predicting cardiovascular events and all-cause death in chronic kidney disease: A population-based prospective cohort study in Japan. <i>International Journal of Cardiology</i> , 2014, 175, 576-577.	1.7	0
24	Low educational level increases functional disability risk subsequent to heart failure in Japan: On behalf of the Iwate KENCO study group. <i>PLoS ONE</i> , 2021, 16, e0253017.	2.5	0