## Mark Van Der Gaag

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4083646/publications.pdf

Version: 2024-02-01

233 papers

10,178 citations

41258 49 h-index 43802

g-index

276 all docs

276 docs citations

times ranked

276

7968 citing authors

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | The five-factor model of the Positive and Negative Syndrome Scale II: A ten-fold cross-validation of a revised model. Schizophrenia Research, 2006, 85, 280-287.  | 1.1 | 466       |
| 2  | EPA guidance on the early intervention in clinical high risk states of psychoses. European Psychiatry, 2015, 30, 388-404.   | 0.1 | 390       |
| 3  | Preventing a first episode of psychosis: Meta-analysis of randomized controlled prevention trials of 12month and longer-term follow-ups. Schizophrenia Research, 2013, 149, 56-62.  | 1.1 | 333       |
| 4  | EPA guidance on the early detection of clinical high risk states of psychoses. European Psychiatry, 2015, 30, 405-416.  | 0.1 | 318       |
| 5  | The Validity of the 16-Item Version of the Prodromal Questionnaire (PQ-16) to Screen for Ultra High Risk of Developing Psychosis in the General Help-Seeking Population. Schizophrenia Bulletin, 2012, 38, 1288-1296.   | 2.3 | 304       |
| 6  | Psychological Interventions for Psychosis: A Meta-Analysis of Comparative Outcome Studies. American Journal of Psychiatry, 2014, 171, 523-538.  | 4.0 | 277       |
| 7  | Prolonged Exposure vs Eye Movement Desensitization and Reprocessing vs Waiting List for Posttraumatic Stress Disorder in Patients With a Psychotic Disorder. JAMA Psychiatry, 2015, 72, 259.  | 6.0 | 267       |
| 8  | Identifying Gene-Environment Interactions in Schizophrenia: Contemporary Challenges for Integrated, Large-scale Investigations. Schizophrenia Bulletin, 2014, 40, 729-736.  | 2.3 | 229       |
| 9  | Genetic Risk and Outcome of Psychosis (GROUP), a multi site longitudinal cohort study focused on gene–environment interaction: objectives, sample characteristics, recruitment and assessment methods. International Journal of Methods in Psychiatric Research, 2012, 21, 205-221. | 1.1 | 224       |
| 10 | The effects of individually tailored formulation-based cognitive behavioural therapy in auditory hallucinations and delusions: A meta-analysis. Schizophrenia Research, 2014, 156, 30-37.   | 1.1 | 221       |
| 11 | CRITICAL ANALYSIS OF THE CURRENT TREATMENT GUIDELINES FOR COMPLEX PTSD IN ADULTS. Depression and Anxiety, 2016, 33, 359-369.  | 2.0 | 208       |
| 12 | Stigma moderates the associations of insight with depressed mood, low self-esteem, and low quality of life in patients with schizophrenia spectrum disorders. Schizophrenia Research, 2009, 115, 363-369.   | 1.1 | 176       |
| 13 | Two subdomains of negative symptoms in psychotic disorders: Established and confirmed in two large cohorts. Journal of Psychiatric Research, 2013, 47, 718-725.   | 1.5 | 169       |
| 14 | Cognitive Behavioral Therapy for Subjects at Ultrahigh Risk for Developing Psychosis: A Randomized Controlled Clinical Trial. Schizophrenia Bulletin, 2012, 38, 1180-1188.  | 2.3 | 168       |
| 15 | The five-factor model of the Positive and Negative Syndrome Scale I: Confirmatory factor analysis fails to confirm 25 published five-factor solutions. Schizophrenia Research, 2006, 85, 273-279.   | 1.1 | 157       |
| 16 | Psychological Therapies for Auditory Hallucinations (Voices): Current Status and Key Directions for Future Research. Schizophrenia Bulletin, 2014, 40, S202-S212.   | 2.3 | 153       |
| 17 | The Treatment of Hallucinations in Schizophrenia Spectrum Disorders. Schizophrenia Bulletin, 2012, 38, 704-714.   | 2.3 | 150       |
| 18 | Trauma and recent life events in individuals at ultra high risk for psychosis: Review and meta-analysis. Schizophrenia Research, 2015, 161, 143-149.  | 1.1 | 147       |

| #  | Article   | IF  | Citations |
|----|---|-----|-----------|
| 19 | Virtual-reality-based cognitive behavioural therapy versus waiting list control for paranoid ideation and social avoidance in patients with psychotic disorders: a single-blind randomised controlled trial. Lancet Psychiatry,the, 2018, 5, 217-226. | 3.7 | 144       |
| 20 | Cognitive–behavioural therapy for refractory psychotic symptoms of schizophrenia resistant to atypical antipsychotic medication. British Journal of Psychiatry, 2005, 186, 324-330.   | 1.7 | 141       |
| 21 | A Meta-Analysis of Social Skills Training and Related Interventions for Psychosis. Schizophrenia<br>Bulletin, 2018, 44, 475-491.  | 2.3 | 141       |
| 22 | The effectiveness of peer support groups in psychosis: a randomized controlled trial. Acta Psychiatrica Scandinavica, 2008, 118, 64-72.   | 2.2 | 139       |
| 23 | Adapted cognitive–behavioural therapy required for targeting negative symptoms in schizophrenia: meta-analysis and meta-regression. Psychological Medicine, 2015, 45, 453-465.  | 2.7 | 126       |
| 24 | IS IT TIME TO DEVELOP A NEW COGNITIVE THERAPY FOR PSYCHOSIS—COGNITIVE REMEDIATION THERAPY (CRT)?. Clinical Psychology Review, 2001, 21, 1227-1256.  | 6.0 | 117       |
| 25 | Treating trauma in psychosis with EMDR: A pilot study. Journal of Behavior Therapy and Experimental Psychiatry, 2012, 43, 664-671.  | 0.6 | 116       |
| 26 | Environmental Social Stress, Paranoia and Psychosis Liability: A Virtual Reality Study. Schizophrenia Bulletin, 2016, 42, 1363-1371.  | 2.3 | 106       |
| 27 | A Controlled Trial of Cognitive Remediation in Schizophrenia. Schizophrenia Bulletin, 2002, 28, 167-176.  | 2.3 | 104       |
| 28 | Trauma-Focused Treatment in PTSD Patients With Psychosis: Symptom Exacerbation, Adverse Events, and Revictimization. Schizophrenia Bulletin, 2016, 42, 693-702.   | 2.3 | 104       |
| 29 | A Neuropsychiatric Model of Biological and Psychological Processes in the Remission of Delusions and Auditory Hallucinations. Schizophrenia Bulletin, 2006, 32, S113-S122.  | 2.3 | 103       |
| 30 | Cannabis use and transition to psychosis in individuals at ultra-high risk: review and meta-analysis. Psychological Medicine, 2016, 46, 673-681.  | 2.7 | 103       |
| 31 | Metacognitive reflection and insight therapy (MERIT) for patients with schizophrenia. Psychological Medicine, 2019, 49, 303-313.  | 2.7 | 102       |
| 32 | Development of the Davos Assessment of Cognitive Biases Scale (DACOBS). Schizophrenia Research, 2013, 144, 63-71.   | 1.1 | 93        |
| 33 | Prolonged exposure and EMDR for PTSD <i>v</i> . a PTSD waiting-list condition: effects on symptoms of psychosis, depression and social functioning in patients with chronic psychotic disorders. Psychological Medicine, 2016, 46, 2411-2421.         | 2.7 | 93        |
| 34 | Treatment adherence therapy in people with psychotic disorders: randomised controlled trial. British Journal of Psychiatry, 2010, 197, 448-455.   | 1.7 | 89        |
| 35 | Whither the Attenuated Psychosis Syndrome?. Schizophrenia Bulletin, 2012, 38, 1130-1134.  | 2.3 | 85        |
| 36 | Predictive validity of the Trauma Screening Questionnaire in detecting post-traumatic stress disorder in patients with psychotic disorders. British Journal of Psychiatry, 2015, 206, 408-416.  | 1.7 | 78        |

| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 37 | Criminal Victimisation in People with Severe Mental Illness: A Multi-Site Prevalence and Incidence Survey in the Netherlands. PLoS ONE, 2014, 9, e91029.   | 1.1 | 77        |
| 38 | Cognitive Behavioral Therapy for negative symptoms (CBT-n) in psychotic disorders: A pilot study. Journal of Behavior Therapy and Experimental Psychiatry, 2013, 44, 300-306.  | 0.6 | 74        |
| 39 | Detection of people at risk of developing a first psychosis: comparison of two recruitment strategies. Acta Psychiatrica Scandinavica, 2012, 126, 21-30.   | 2.2 | 71        |
| 40 | Early intervention in patients at ultra high risk of psychosis: benefits and risks. Acta Psychiatrica Scandinavica, 2009, 119, 426-442.  | 2.2 | 68        |
| 41 | Efficacy of Acceptance and Commitment Therapy in Daily Life (ACT-DL) in early psychosis: study protocol for a multi-centre randomized controlled trial. Trials, 2019, 20, 769.   | 0.7 | 65        |
| 42 | Competitive Memory Training for treating depression and rumination in depressed older adults: A randomized controlled trial. Behaviour Research and Therapy, 2011, 49, 588-596.  | 1.6 | 63        |
| 43 | Metacognitive training for schizophrenia spectrum patients: a meta-analysis on outcome studies.<br>Psychological Medicine, 2016, 46, 47-57.  | 2.7 | 63        |
| 44 | Evidence for a Cognitive Model of Auditory Hallucinations. Journal of Nervous and Mental Disease, 2003, 191, 542-545.  | 0.5 | 62        |
| 45 | Metacognitive group training for schizophrenia spectrum patients with delusions: a randomized controlled trial. Psychological Medicine, 2014, 44, 3025-3035.   | 2.7 | 58        |
| 46 | Initial evaluation of the effects of competitive memory training (COMET) on depression in schizophreniaâ€spectrum patients with persistent auditory verbal hallucinations: A randomized controlled trial. British Journal of Clinical Psychology, 2012, 51, 158-171. | 1.7 | 57        |
| 47 | Four-Year Follow-up of Cognitive Behavioral Therapy in Persons at Ultra-High Risk for Developing<br>Psychosis: The Dutch Early Detection Intervention Evaluation (EDIE-NL) Trial. Schizophrenia Bulletin,<br>2016, 42, 1243-1252.                                    | 2.3 | 57        |
| 48 | Development of Proteomic Prediction Models for Transition to Psychotic Disorder in the Clinical High-Risk State and Psychotic Experiences in Adolescence. JAMA Psychiatry, 2021, 78, 77.   | 6.0 | 57        |
| 49 | Depression and social anxiety in help-seeking patients with an ultra-high risk for developing psychosis.<br>Psychiatry Research, 2013, 209, 309-313.   | 1.7 | 55        |
| 50 | Pathways to psychosis: Help-seeking behavior in the prodromal phase. Schizophrenia Research, 2011, 132, 213-219.   | 1.1 | 54        |
| 51 | Brave New WorldsReview and Update on Virtual Reality Assessment and Treatment in Psychosis.<br>Schizophrenia Bulletin, 2014, 40, 1194-1197.  | 2.3 | 54        |
| 52 | Hallucinations in borderline personality disorder: Prevalence, characteristics and associations with comorbid symptoms and disorders. Scientific Reports, 2017, 7, 13920.  | 1.6 | 52        |
| 53 | Preventive interventions for individuals at ultra high risk for psychosis: An updated and extended meta-analysis. Clinical Psychology Review, 2021, 86, 102005.  | 6.0 | 52        |
| 54 | A manual-based individual therapy to improve metacognition in schizophrenia: protocol of a multi-center RCT. BMC Psychiatry, 2014, 14, 27.   | 1.1 | 49        |

| #  | Article   | IF  | Citations |
|----|---|-----|-----------|
| 55 | Cost-effectiveness of preventing first-episode psychosis in ultra-high-risk subjects: multi-centre randomized controlled trial. Psychological Medicine, 2015, 45, 1435-1446.  | 2.7 | 49        |
| 56 | Cognitive disorders and negative symptoms as correlates of motivational deficits in psychotic patients. Psychological Medicine, 1994, 24, 869-884.  | 2.7 | 48        |
| 57 | A single blind randomized controlled trial of cognitive behavioural therapy in a help-seeking population with an At Risk Mental State for psychosis: the Dutch Early Detection and Intervention Evaluation (EDIE-NL) trial. Trials, 2010, 11, 30. | 0.7 | 46        |
| 58 | Creating a Supportive Environment: Peer Support Groups for Psychotic Disorders. Schizophrenia Bulletin, 2015, 41, 1211-1213.  | 2.3 | 46        |
| 59 | Psychosocial functioning in firstâ€episode psychosis and associations with neurocognition, social cognition, psychotic and affective symptoms. Microbial Biotechnology, 2017, 11, 23-36.  | 0.9 | 46        |
| 60 | Childhood trauma, psychosis liability and social stress reactivity: a virtual reality study. Psychological Medicine, 2016, 46, 3339-3348.   | 2.7 | 45        |
| 61 | Long-term outcomes of trauma-focused treatment in psychosis. British Journal of Psychiatry, 2018, 212, 180-182.   | 1.7 | 45        |
| 62 | Measuring Empowerment Among People With Psychotic Disorders: A Comparison of Three Instruments. Psychiatric Services, 2008, 59, 1338-1342.  | 1.1 | 44        |
| 63 | Anxiety Partially Mediates Cybersickness Symptoms in Immersive Virtual Reality Environments. Cyberpsychology, Behavior, and Social Networking, 2018, 21, 187-193.   | 2.1 | 44        |
| 64 | Dysregulated Lipid Metabolism Precedes Onset of Psychosis. Biological Psychiatry, 2021, 89, 288-297.  | 0.7 | 42        |
| 65 | Fewer symptoms vs. more side-effects in schizophrenia? Opposing pathways between antipsychotic medication compliance and quality of life. Schizophrenia Research, 2009, 113, 27-33.   | 1.1 | 41        |
| 66 | Virtual Reality Experiments Linking Social Environment and Psychosis: A Pilot Study. Cyberpsychology, Behavior, and Social Networking, 2014, 17, 191-195.   | 2.1 | 41        |
| 67 | Practical Implications of Metacognitively Oriented Psychotherapy in Psychosis. Journal of Nervous and Mental Disease, 2016, 204, 713-716.   | 0.5 | 41        |
| 68 | The effect of mind-body and aerobic exercise on negative symptoms in schizophrenia: A meta-analysis. Psychiatry Research, 2019, 279, 295-305.   | 1.7 | 41        |
| 69 | To continue or not to continue? Antipsychotic medication maintenance versus dose-reduction/discontinuation in first episode psychosis: HAMLETT, a pragmatic multicenter single-blind randomized controlled trial. Trials, 2020, 21, 147.          | 0.7 | 41        |
| 70 | Effect of Crisis Plans on Admissions and Emergency Visits: A Randomized Controlled Trial. PLoS ONE, 2014, 9, e91882.  | 1.1 | 40        |
| 71 | Childhood trauma and clinical outcome in patients at ultra-high risk of transition to psychosis.<br>Schizophrenia Research, 2015, 169, 193-198.   | 1.1 | 40        |
| 72 | Suicidality and hospitalisation in patients with borderline personality disorder who experience auditory verbal hallucinations. European Psychiatry, 2017, 41, 47-52.   | 0.1 | 40        |

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 73 | A multi-site single blind clinical study to compare the effects of prolonged exposure, eye movement desensitization and reprocessing and waiting list on patients with a current diagnosis of psychosis and co morbid post traumatic stress disorder: study protocol for the randomized controlled trial Treating Trauma in Psychosis. Trials, 2013, 14, 151. | 0.7 | 39        |
| 74 | What Constitutes Sufficient Evidence for Case Formulation–Driven CBT for Psychosis? Cumulative Meta-analysis of the Effect on Hallucinations and Delusions. Schizophrenia Bulletin, 2020, 46, 1072-1085.  | 2.3 | 39        |
| 75 | Child Maltreatment and Clinical Outcome in Individuals at Ultra-High Risk for Psychosis in the EU-GEI<br>High Risk Study. Schizophrenia Bulletin, 2018, 44, 584-592.  | 2.3 | 38        |
| 76 | Psychotic symptoms, cognition and affect as predictors of psychosocial problems and functional change in first-episode psychosis. Schizophrenia Research, 2014, 158, 113-119.   | 1.1 | 37        |
| 77 | The dynamics of pAL2-1 homologous linear plasmids in Podospora anserina. Molecular Genetics and Genomics, 1998, 258, 521-529.   | 2.4 | 36        |
| 78 | Cognitive–behavioural therapy for persistent and recurrent psychosis in people with schizophrenia-spectrum disorder: cost-effectiveness analysis. British Journal of Psychiatry, 2011, 198, 59-65.  | 1.7 | 36        |
| 79 | Cost-Effectiveness of Virtual Reality Cognitive Behavioral Therapy for Psychosis: Health-Economic Evaluation Within a Randomized Controlled Trial. Journal of Medical Internet Research, 2020, 22, e17098.  | 2.1 | 35        |
| 80 | Duration of untreated psychosis and the long-term course of schizophrenia. European Psychiatry, 2000, 15, 264-267.  | 0.1 | 33        |
| 81 | The Validity of the DSM-IV Diagnostic Classification System of Non-Affective Psychoses. Australian and New Zealand Journal of Psychiatry, 2011, 45, 1061-1068.  | 1.3 | 33        |
| 82 | Development of a group and familyâ€based cognitive behavioural therapy program for youth at risk for psychosis. Microbial Biotechnology, 2016, 10, 511-521.   | 0.9 | 31        |
| 83 | CBT in the prevention of psychosis and other severe mental disorders in patients with an at risk mental state: A review and proposed next steps. Schizophrenia Research, 2019, 203, 88-93.  | 1.1 | 31        |
| 84 | Byssal detachment underestimates tolerance of mussels to toxic compounds. Marine Pollution Bulletin, 2005, 50, 20-29.   | 2.3 | 29        |
| 85 | REFLEX, a social-cognitive group treatment to improve insight in schizophrenia: study protocol of a multi-center RCT. BMC Psychiatry, 2011, 11, 161.  | 1.1 | 29        |
| 86 | Four-Year Cost-effectiveness of Cognitive Behavior Therapy for Preventing First-episode Psychosis: The Dutch Early Detection Intervention Evaluation (EDIE-NL) Trial. Schizophrenia Bulletin, 2017, 43, sbw084.   | 2.3 | 29        |
| 87 | Individualized Prediction of Transition to Psychosis in 1,676 Individuals at Clinical High Risk:<br>Development and Validation of a Multivariable Prediction Model Based on Individual Patient Data<br>Meta-Analysis. Frontiers in Psychiatry, 2019, 10, 345.   | 1.3 | 29        |
| 88 | Effectiveness of trauma-focused treatment for patients with psychosis with and without the dissociative subtype of post-traumatic stress disorder. British Journal of Psychiatry, 2016, 209, 347-348.   | 1.7 | 28        |
| 89 | Screening for psychosis risk among adolescents in Child and Adolescent Mental Health Services: a description of the first step with the 16â€item version of the Prodromal Questionnaire (PQâ€16). Microbial Biotechnology, 2018, 12, 669-676.   | 0.9 | 26        |
| 90 | Social environments and interpersonal distance regulation in psychosis: A virtual reality study. Schizophrenia Research, 2018, 192, 96-101.   | 1,1 | 26        |

| #   | Article  | IF  | Citations |
|-----|--|-----|-----------|
| 91  | A network analysis of post-traumatic stress and psychosis symptoms. Psychological Medicine, 2021, 51, 2485-2492.   | 2.7 | 26        |
| 92  | CBT for Those at Risk of a First Episode Psychosis. , 0, , .   |     | 26        |
| 93  | A randomised controlled trial of positive memory training for the treatment of depression within schizophrenia. BMC Psychiatry, 2015, 15, 85.  | 1.1 | 24        |
| 94  | Crisis Plans Facilitated by Patient Advocates are Better than those Drawn up by Clinicians: Results from an RCT. Administration and Policy in Mental Health and Mental Health Services Research, 2014, 41, 220-227.                    | 1.2 | 23        |
| 95  | The effect of childhood adversity on 4-year outcome in individuals at ultra high risk for psychosis in the Dutch Early Detection Intervention Evaluation (EDIE-NL) Trial. Psychiatry Research, 2017, 247, 55-62.                       | 1.7 | 23        |
| 96  | Association of Adverse Outcomes With Emotion Processing and Its Neural Substrate in Individuals at Clinical High Risk for Psychosis. JAMA Psychiatry, 2020, 77, 190.   | 6.0 | 23        |
| 97  | Cognitive functioning throughout adulthood and illness stages in individuals with psychotic disorders and their unaffected siblings. Molecular Psychiatry, 2021, 26, 4529-4543.  | 4.1 | 23        |
| 98  | Prescription of antipsychotic medication to patients at ultra high risk of developing psychosis. International Clinical Psychopharmacology, 2009, 24, 223-228.   | 0.9 | 22        |
| 99  | Economic Aspects of Peer Support Groups for Psychosis. Community Mental Health Journal, 2011, 47, 99-105.  | 1.1 | 22        |
| 100 | The Cognitive Biases Questionnaire for Psychosis (CBQ-P) and the Davos Assessment of Cognitive Biases (DACOBS): Validation in a Flemish sample of psychotic patients and healthy controls. Schizophrenia Research, 2013, 147, 310-314. | 1.1 | 22        |
| 101 | Metacognitive beliefs, beliefs about voices and affective symptoms in patients with severe auditory verbal hallucinations. British Journal of Clinical Psychology, 2013, 52, 235-248.  | 1.7 | 22        |
| 102 | COMT Val 158 Met genotype and cannabis use in people with an At Risk Mental State for psychosis: Exploring Gene x Environment interactions. Schizophrenia Research, 2016, 174, 24-28.  | 1.1 | 22        |
| 103 | Reducing distress and improving social functioning in daily life in people with auditory verbal hallucinations: study protocol for the †Temstem' randomised controlled trial. BMJ Open, 2018, 8, e020537.                              | 0.8 | 22        |
| 104 | Symptom changes in five dimensions of the Positive and Negative Syndrome Scale in refractory psychosis. European Archives of Psychiatry and Clinical Neuroscience, 2014, 264, 673-682.   | 1.8 | 21        |
| 105 | Factor Analysis of the Scale of Prodromal Symptoms: Differentiating between Negative and Depression Symptoms. Psychopathology, 2011, 44, 379-385.  | 1.1 | 20        |
| 106 | Working Alliance in Patients with Severe Mental Illness Who Need a Crisis Intervention Plan. Community Mental Health Journal, 2016, 52, 102-108.   | 1.1 | 20        |
| 107 | The effect of dialectical behaviour therapy in autism spectrum patients with suicidality and/ or self-destructive behaviour (DIASS): study protocol for a multicentre randomised controlled trial. BMC Psychiatry, 2020, 20, 127.      | 1.1 | 20        |
| 108 | Recognition of early warning signs in patients with schizophrenia: A review of the literature. International Journal of Mental Health Nursing, 2004, 13, 107-116.  | 2.1 | 19        |

| #   | Article  | IF  | CITATIONS |
|-----|--|-----|-----------|
| 109 | Understanding and Improving Treatment Adherence in Patients with Psychotic Disorders: A Review and a Proposed Intervention. Current Psychiatry Reviews, 2006, 2, 487-494.  | 0.9 | 19        |
| 110 | Bridging the science-to-service gap in schizophrenia care in the Netherlands: the Schizophrenia Quality Improvement Collaborative. International Journal for Quality in Health Care, 2013, 25, 626-632.                                    | 0.9 | 19        |
| 111 | High psychosis liability is associated with altered autonomic balance during exposure to Virtual Reality social stressors. Schizophrenia Research, 2017, 184, 14-20.   | 1.1 | 19        |
| 112 | Self-esteem moderates affective and psychotic responses to social stress in psychosis: A virtual reality study. Schizophrenia Research, 2018, 202, 80-85.  | 1.1 | 19        |
| 113 | Gender differences of patients at-risk for psychosis regarding symptomatology, drug use, comorbidity and functioning – Results from the EU-GEI study. European Psychiatry, 2019, 59, 52-59.  | 0.1 | 19        |
| 114 | Health–economic benefits of treating trauma in psychosis. Högre Utbildning, 2019, 10, 1565032.   | 1.4 | 19        |
| 115 | The practice of early recognition and early intervention to prevent psychotic relapse in patients with schizophrenia: an exploratory study. Part 1. Journal of Psychiatric and Mental Health Nursing, 2002, 9, 347-355.                    | 1.2 | 18        |
| 116 | Ethnicity and baseline symptomatology in patients with an At Risk Mental State for psychosis. Psychological Medicine, 2012, 42, 247-256.   | 2.7 | 18        |
| 117 | Social-cognitive risk factors for violence in psychosis: A discriminant function analysis. Psychiatry Research, 2018, 265, 93-99.  | 1.7 | 18        |
| 118 | Are social phobia and paranoia related, and which comes first?. Psychosis, 2009, 1, 29-38.   | 0.4 | 17        |
| 119 | The efficacy of CBT for severe mental illness and the challenge of dissemination in routine care.<br>World Psychiatry, 2014, 13, 257-258.  | 4.8 | 17        |
| 120 | A time-series network approach to auditory verbal hallucinations: Examining dynamic interactions using experience sampling methodology. Schizophrenia Research, 2020, 215, 148-156.  | 1.1 | 17        |
| 121 | Efficacy and Moderators of Cognitive Behavioural Therapy for Psychosis Versus Other Psychological Interventions: An Individual-Participant Data Meta-Analysis. Frontiers in Psychiatry, 2020, 11, 402.                                     | 1.3 | 17        |
| 122 | Stress reactivity as a putative mechanism linking childhood trauma with clinical outcomes in individuals at ultra-high-risk for psychosis: Findings from the EU-GEI High Risk Study. Epidemiology and Psychiatric Sciences, 2021, 30, e40. | 1.8 | 17        |
| 123 | The practice of early recognition and early intervention to prevent psychotic relapse in patients with schizophrenia: an exploratory study. Part 2. Journal of Psychiatric and Mental Health Nursing, 2002, 9, 357-363.                    | 1.2 | 16        |
| 124 | An Intervention Study to Prevent Relapse in Patients With Schizophrenia. Journal of Nursing Scholarship, 2006, 38, 42-49.  | 1.1 | 16        |
| 125 | Depressive symptoms are associated with (sub)clinical psychotic symptoms in patients with non-affective psychotic disorder, siblings and healthy controls. Psychological Medicine, 2013, 43, 747-756.                                      | 2.7 | 16        |
| 126 | The distribution of self-reported psychotic-like experiences in non-psychotic help-seeking mental health patients in the general population; a factor mixture analysis. Social Psychiatry and Psychiatric Epidemiology, 2014, 49, 349-358. | 1.6 | 16        |

| #   | Article  | IF  | Citations |
|-----|--|-----|-----------|
| 127 | Social cognitive group treatment for impaired insight in psychosis: A multicenter randomized controlled trial. Schizophrenia Research, 2019, 206, 362-369.   | 1.1 | 16        |
| 128 | Clinical, cognitive and neuroanatomical associations of serum NMDAR autoantibodies in people at clinical high risk for psychosis. Molecular Psychiatry, 2021, 26, 2590-2604.                                 | 4.1 | 16        |
| 129 | Measuring personal recovery in people with a psychotic disorder based on CHIME: A comparison of three validated measures. International Journal of Mental Health Nursing, 2020, 29, 808-819.                 | 2.1 | 16        |
| 130 | Cognitive behavioral therapy for social activation in recent-onset psychosis: Randomized controlled trial Journal of Consulting and Clinical Psychology, 2019, 87, 151-160.                                  | 1.6 | 15        |
| 131 | The effect of thiopental on glutamate metabolism in mousecerebellar astrocytes in vitro.<br>Neuroscience Letters, 2001, 304, 141-144.  | 1.0 | 14        |
| 132 | Relapse prevention in patients with schizophrenia. Archives of Psychiatric Nursing, 2003, 17, 117-125.   | 0.7 | 14        |
| 133 | Recovery Style Predicts Remission at One-Year Follow-Up in Outpatients With Schizophrenia Spectrum Disorders. Journal of Nervous and Mental Disease, 2011, 199, 295-300.                                     | 0.5 | 14        |
| 134 | Why Do Patients with Schizophrenia Who Have Poor Insight Still Take Antipsychotics? Memory Deficits as Moderators Between Adherence Belief and Behavior. Journal of Psychiatric Practice, 2011, 17, 320-329. | 0.3 | 14        |
| 135 | Determinants of Completion and Use of Psychiatric Advance Statements in Mental Health Care in the Netherlands. Psychiatric Services, 2016, 67, 858-863.  | 1.1 | 14        |
| 136 | Sex differences in cognitive functioning of patients at-risk for psychosis and healthy controls: Results from the European Gene–Environment Interactions study. European Psychiatry, 2020, 63, e25.          | 0.1 | 14        |
| 137 | Emotion Recognition and Adverse Childhood Experiences in Individuals at Clinical High Risk of Psychosis. Schizophrenia Bulletin, 2020, 46, 823-833.  | 2.3 | 14        |
| 138 | Negative Beliefs about Voices in Patients with Borderline Personality Disorder Are Associated with Distress: A Plea for Cognitive-Behavioural Therapy?. Psychopathology, 2017, 50, 255-261.                  | 1.1 | 13        |
| 139 | Virtual reality based cognitive behavioral therapy for paranoia: Effects on mental states and the dynamics among them. Schizophrenia Research, 2020, 222, 227-234.   | 1.1 | 13        |
| 140 | Clinical, societal and personal recovery in schizophrenia spectrum disorders across time: states and annual transitions. British Journal of Psychiatry, 2021, 219, 401-408.                                  | 1.7 | 13        |
| 141 | Development of a stage-dependent prognostic model to predict psychosis in ultra-high-risk patients seeking treatment for co-morbid psychiatric disorders. Psychological Medicine, 2016, 46, 1839-1851.       | 2.7 | 12        |
| 142 | Effect of virtual reality exposure therapy on social participation in people with a psychotic disorder (VRETp): study protocol for a randomized controlled trial. Trials, 2016, 17, 25.                      | 0.7 | 12        |
| 143 | Positive memory training for the treatment of depression in schizophrenia: A randomised controlled trial. Behaviour Research and Therapy, 2020, 135, 103734.   | 1.6 | 12        |
| 144 | Efficacy of Acceptance and Commitment Therapy in Daily Life in Early Psychosis: Results from the Multi-Center INTERACT Randomized Controlled Trial. Psychotherapy and Psychosomatics, 2022, 91, 411-423.     | 4.0 | 12        |

| #   | Article   | IF  | Citations |
|-----|---|-----|-----------|
| 145 | Symptomatic and functional remission and its associations with quality of life in patients with psychotic disorder in Assertive Community Treatment teams. Comprehensive Psychiatry, 2012, 53, 1174-1180.                         | 1.5 | 11        |
| 146 | Temperature dependent larval occurrence and spat settlement of the invasive brackish water bivalve Mytilopsis leucophaeata (Conrad, 1831) (Dreissenidae). Journal of Sea Research, 2014, 87, 30-34.                               | 0.6 | 11        |
| 147 | The influence of adjunctive treatment and metacognitive deficits in schizophrenia on the experience of work. Schizophrenia Research, 2014, 157, 107-111.  | 1.1 | 11        |
| 148 | Predicting trauma-focused treatment outcome in psychosis. Schizophrenia Research, 2016, 176, 239-244.   | 1.1 | 11        |
| 149 | Self-reported Cognitive Biases Moderate the Associations Between Social Stress and Paranoid Ideation in a Virtual Reality Experimental Study. Schizophrenia Bulletin, 2017, 44, 749-756.  | 2.3 | 11        |
| 150 | Peer support and skills training through an eating club for people with psychotic disorders: A feasibility study. Journal of Behavior Therapy and Experimental Psychiatry, 2019, 64, 80-86.                                       | 0.6 | 11        |
| 151 | Designing and Testing of a Health-Economic Markov Model for Prevention and Treatment of Early Psychosis. Expert Review of Pharmacoeconomics and Outcomes Research, 2020, 20, 269-279.   | 0.7 | 11        |
| 152 | Relationship between jumping to conclusions and clinical outcomes in people at clinical high-risk for psychosis. Psychological Medicine, 2022, 52, 1569-1577.   | 2.7 | 11        |
| 153 | Exposing therapists to trauma-focused treatment in psychosis: effects on credibility, expected burden, and harm expectancies. HÃ $\P$ gre Utbildning, 2016, 7, 31712.   | 1.4 | 10        |
| 154 | The Prodromal Questionnaire: a case for IRT-based adaptive testing of psychotic experiences?. International Journal of Methods in Psychiatric Research, 2017, 26, e1518.  | 1.1 | 10        |
| 155 | Obsessive-Compulsive Symptoms and Other Symptoms of the At-risk Mental State for Psychosis: A Network Perspective. Schizophrenia Bulletin, 2021, 47, 1018-1028.   | 2.3 | 10        |
| 156 | Verbal memory performance predicts remission and functional outcome in people at clinical high-risk for psychosis. Schizophrenia Research: Cognition, 2022, 28, 100222.   | 0.7 | 10        |
| 157 | Does Treatment Adherence Therapy reduce expense of healthcare use in patients with psychotic disorders? Cost-minimization analysis in a randomized controlled trial. Schizophrenia Research, 2011, 133, 47-53.                    | 1.1 | 9         |
| 158 | Schizophrenia and antipsychotic medication $\hat{a} \in$ Better adherence, better outcomes? Schizophrenia Research, 2013, 151, 296-297.   | 1.1 | 9         |
| 159 | Cognitive deficits and ethnicity: a cohort study of early psychosis patients in The Netherlands. Social Psychiatry and Psychiatric Epidemiology, 2013, 48, 37-47.   | 1.6 | 9         |
| 160 | Evidence for preventive treatments in young patients at clinical high risk of psychosis: the need for context. Lancet Psychiatry, the, 2020, 7, 378-380.  | 3.7 | 9         |
| 161 | Impact of Comorbid Affective Disorders on Longitudinal Clinical Outcomes in Individuals at Ultra-high Risk for Psychosis. Schizophrenia Bulletin, 2022, 48, 100-110.  | 2.3 | 9         |
| 162 | Efficacy of a transdiagnostic ecological momentary intervention for improving self-esteem (SELFIE) in youth exposed to childhood adversity: study protocol for a multi-center randomized controlled trial. Trials, 2021, 22, 641. | 0.7 | 9         |

| #   | Article   | IF  | Citations |
|-----|---|-----|-----------|
| 163 | Relapse prevention in patients with schizophrenia: the application of an intervention protocol in nursing practice. Archives of Psychiatric Nursing, 2003, 17, 165-172.   | 0.7 | 8         |
| 164 | Letter to the Editor: Should we focus on quality or quantity in meta-analyses?. Psychological Medicine, 2016, 46, 2003-2005.  | 2.7 | 8         |
| 165 | Economic evaluation of brief cognitive behavioural therapy for social activation in recent-onset psychosis. PLoS ONE, 2018, 13, e0206236.   | 1.1 | 8         |
| 166 | The associations between migrant status and ethnicity and the identification of individuals at ultra-high risk for psychosis and transition to psychosis: a systematic review. Social Psychiatry and Psychiatric Epidemiology, 2021, 56, 1923-1941. | 1.6 | 8         |
| 167 | Psychopathology, cognition and outcome in Dutch and immigrant firstâ€episode psychosis patients. Microbial Biotechnology, 2019, 13, 646-656.  | 0.9 | 7         |
| 168 | Cross Validation of the Prodromal Questionnaire 16-Item Version in an Adolescent Help-Seeking Population. Schizophrenia Bulletin Open, 2020, $1$ , .  | 0.9 | 7         |
| 169 | Functional recovery of individuals with serious mental illnesses: Development and testing of a new short instrument for routine outcome monitoring Psychiatric Rehabilitation Journal, 2018, 41, 341-350.   | 0.8 | 7         |
| 170 | The Effectiveness of Transdiagnostic Applications of Competitive Memory Training (COMET) on Low Self-Esteem and Comorbid Depression: A Meta-analysis of Randomized Controlled Trials. Cognitive Therapy and Research, 2022, 46, 532-543.            | 1.2 | 7         |
| 171 | Voice phenomenology as a mirror of the past. Psychological Medicine, 2023, 53, 2954-2962.   | 2.7 | 7         |
| 172 | Psychosocial Treatments for Clinical High Risk Individuals. Schizophrenia Bulletin, 2015, 41, 22-22.  | 2.3 | 6         |
| 173 | Predicting therapy success from the outset: The moderating effect of insight into the illness on metacognitive psychotherapy outcome among persons with schizophrenia. Clinical Psychology and Psychotherapy, 2019, 26, 650-660.                    | 1.4 | 6         |
| 174 | Pre-training inter-rater reliability of clinical instruments in an international psychosis research project. Schizophrenia Research, 2020, 230, 104-107.  | 1.1 | 6         |
| 175 | Association between tobacco use and symptomatology in individuals at ultra-high risk to develop a psychosis: A longitudinal study. Schizophrenia Research, 2021, 236, 48-53.  | 1.1 | 6         |
| 176 | Characteristics and stability of hallucinations and delusions in patients with borderline personality disorder. Comprehensive Psychiatry, 2022, 113, 152290.  | 1.5 | 6         |
| 177 | Treating prolactinoma and psychosis: medication and cognitive behavioural therapy. BMJ Case Reports, 2011, 2011, bcr0720103185-bcr0720103185.   | 0.2 | 5         |
| 178 | The association between migrant status and transition in an ultra-high risk for psychosis population. Social Psychiatry and Psychiatric Epidemiology, 2021, 56, 943-952.  | 1.6 | 5         |
| 179 | Ultrahigh risk for developing psychosis and psychotic personality organization. Microbial Biotechnology, 2019, 13, 673-676.   | 0.9 | 4         |
| 180 | Changes in Posttraumatic Cognitions Mediate the Effects of Trauma-Focused Therapy on Paranoia. Schizophrenia Bulletin Open, 2020, $1$ , .   | 0.9 | 4         |

| #   | Article   | IF  | Citations |
|-----|---|-----|-----------|
| 181 | UP'S: A Cohort Study on Recovery in Psychotic Disorder Patients: Design Protocol. Frontiers in Psychiatry, 2020, 11, 609530.  | 1.3 | 4         |
| 182 | Detection of Domestic Violence and Abuse by Community Mental Health Teams Using the BRAVE Intervention: A Multicenter, Cluster Randomized Controlled Trial. Journal of Interpersonal Violence, 2022, 37, NP14310-NP14336.   | 1.3 | 4         |
| 183 | Momentary Manifestations of Negative Symptoms as Predictors of Clinical Outcomes in People at High Risk for Psychosis: Experience Sampling Study. JMIR Mental Health, 2021, 8, e30309.  | 1.7 | 4         |
| 184 | Virtual reality to study responses to social environmental stressors in individuals with and without psychosis. Studies in Health Technology and Informatics, 2011, 167, 86-91.   | 0.2 | 4         |
| 185 | Sensory processing deficiencies in patients with borderline personality disorder who experience auditory verbal hallucinations. Psychiatry Research, 2019, 281, 112545.   | 1.7 | 3         |
| 186 | From Speech Illusions to Onset of Psychotic Disorder: Applying Network Analysis to an Experimental Measure of Aberrant Experiences. Schizophrenia Bulletin Open, 2020, 1, .   | 0.9 | 3         |
| 187 | Assessing the Acceptability, Feasibility and Sustainability of an Intervention to Increase Detection of Domestic Violence and Abuse in Patients Suffering From Severe Mental Illness: A Qualitative Study. Frontiers in Psychiatry, 2020, $11,581031$ .                                     | 1.3 | 3         |
| 188 | Improving screening methods for psychosis in an adolescent help-seeking population using the Child Behavior Checklist (CBCL) and the Youth Self Report (YSR) versus the Prodromal Questionnaire -16 items version (PQ-16). Child and Adolescent Psychiatry and Mental Health, 2022, 16, 25. | 1.2 | 3         |
| 189 | Rehabilitatie bij chronische schizofrenie: wat voegt cognitieve–functietraining toe?. Dth, 1997, 17, 160-168.   | 0.2 | 2         |
| 190 | Schizofrenie en posttraumatische stress-stoornis. Dth, 2002, 22, 7-13.  | 0.2 | 2         |
| 191 | Field Testing Attenuated Psychosis Syndrome Criteria. American Journal of Psychiatry, 2011, 168, 1221-1221.   | 4.0 | 2         |
| 192 | VIRTUAL REALITY IN PSYCHOSIS: EXPERIMENTS WITH SOCIAL RISK ENVIRONMENTS. Schizophrenia Research, 2012, 136, S20.  | 1.1 | 2         |
| 193 | Family-focused treatment improves attenuated psychotic symptoms, but does not differ from brief treatment in negative symptoms and social functioning in ultra high risk patients aged 12–35 years. Evidence-Based Mental Health, 2015, 18, 59-59.  | 2.2 | 2         |
| 194 | Effects of Aerobic Exercise On Negative Symptoms in Schizophrenia: a Meta-analysis. European Psychiatry, 2015, 30, 927.   | 0.1 | 2         |
| 195 | Response to â€The factor structure of complex posttraumatic stress disorder in traumatized refugees'.<br>Högre Utbildning, 2017, 8, 1307078.  | 1.4 | 2         |
| 196 | Linguistic dual tasking reduces emotionality, vividness and credibility of voice memories in voice-hearing individuals: Results from a controlled trial. Schizophrenia Research, 2020, 216, 249-254.  | 1.1 | 2         |
| 197 | Voice-hearers' beliefs about the causes of their voices. Psychiatry Research, 2021, 302, 113997.  | 1.7 | 2         |
| 198 | Impact of Smoking Behavior on Cognitive Functioning in Persons at Risk for Psychosis and Healthy Controls: A Longitudinal Study. European Psychiatry, 2021, 64, 1-20.   | 0.1 | 2         |

| #   | Article   | IF           | CITATIONS |
|-----|---|--------------|-----------|
| 199 | Satisfaction with social connectedness as a predictor for positive and negative symptoms of psychosis: A PHAMOUS study. Schizophrenia Research, 2021, 238, 121-127.   | 1.1          | 2         |
| 200 | Psychodramatechnieken voor de behandeling van angst bij stemmen. Dth, 2006, 26, 2-9.  | 0.2          | 1         |
| 201 | PREVENTIVE PSYCHOTHERAPY. Schizophrenia Research, 2014, 153, S42-S43.   | 1.1          | 1         |
| 202 | F50. METACOGNITIVE REFLECTION AND INSIGHT THERAPY: A MULTICENTER RANDOMIZED CONTROLLED TRIAL. Schizophrenia Bulletin, 2018, 44, S238-S239.  | 2.3          | 1         |
| 203 | S52. WORKING MECHANISMS OF VIRTUAL REALITY BASED CBT FOR PARANOIA: A RANDOMIZED CONTROLLED TRIAL EXAMINING COGNITIVE BIASES, SCHEMATIC BELIEFS AND SAFETY BEHAVIOR. Schizophrenia Bulletin, 2018, 44, S344-S344.      | 2.3          | 1         |
| 204 | 5.2 COGNITIVE BEHAVIORAL THERAPY FOR SOCIAL ACTIVATION IN RECENT-ONSET PSYCHOSIS: RANDOMIZED CONTROLLED TRIAL. Schizophrenia Bulletin, 2019, 45, S93-S94.   | 2.3          | 1         |
| 205 | Competitive Memory Training. , 2010, , 143-162.   |              | 1         |
| 206 | The results of training of perceptual and cognitive functions with chronic schizophrenic patients. Schizophrenia Research, 1990, 3, 66.   | 1.1          | 0         |
| 207 | S50.01 Assessment of disablement in schizophrenia. European Psychiatry, 2000, 15, 310s-310s.  | 0.1          | О         |
| 208 | Placebo in een leertheoretisch perspectief. Dth, 2001, 21, 156-159.   | 0.2          | 0         |
| 209 | School performance in secondary education shows no decline before the onset of a first episode of psychosis in schizophrenia. Journal of Mental Health, 2003, 12, 585-593.  | 1.0          | O         |
| 210 | TC13E THE EFFECTIVENESS OF â€~PROFESSIONALLY GUIDED' SELF-HELP GROUPS: A MULTI-CENTER RANDON CONTROLLED TRIAL Schizophrenia Research, 2006, 86, S56.  | NIZED<br>I.I | 0         |
| 211 | ARE SOCIAL ANXIETY AND PARANOIA ASSOCIATED, AND WHICH COMES FIRST?. Schizophrenia Research, 2010, 117, 317-318.   | 1.1          | 0         |
| 212 | Predictors of persistence of ultrahigh risk symptoms and predictors of transition into psychosis. European Psychiatry, 2011, 26, 2086-2086.   | 0.1          | 0         |
| 213 | Poster #194 DEPRESSIVE SYMPTOMS ARE ASSOCIATED WITH (SUB) CLINICAL PSYCHOSIS VULNERABILITY IN PATIENTS WITH NON-AFFECTIVE PSYCHOTIC DISORDER, SIBLINGS AND HEALTHY CONTROLS. Schizophrenia Research, 2012, 136, S255. | 1.1          | 0         |
| 214 | Poster #81 SUBDOMAINS OF NEGATIVE SYMPTOMS IN SCHIZOPHRENIA: A TWO-FACTOR MODEL CONFIRMED BY CONFIRMATORY FACTOR ANALYSIS. Schizophrenia Research, 2012, 136, S309.   | 1.1          | 0         |
| 215 | Positive symptoms in atâ€risk mental state: the importance of differentiating within the scope. Microbial Biotechnology, 2013, 7, 100-101.  | 0.9          | O         |
| 216 | Reply. Acta Psychiatrica Scandinavica, 2013, 127, 332-332.  | 2.2          | 0         |

| #   | Article   | IF  | CITATIONS |
|-----|---|-----|-----------|
| 217 | Response to the letters of Dr Amos and Dr Preti and colleagues. Schizophrenia Research, 2014, 153, 237-239.   | 1.1 | O         |
| 218 | THE RESULTS OF EYE MOVEMENT DESENSITISATION AND REPROCESSING AND PROLONGED EXPOSURE IN PATIENTS WITH POSTTRAUMATIC STRESS DISORDER AND CHRONIC PSYCHOTIC DISORDER. Schizophrenia Research, 2014, 153, S74-S75.                                    | 1.1 | O         |
| 219 | O6.5. INVESTIGATING VARIABLES FROM THE NAPLS RISK CALCULATOR FOR PSYCHOSIS IN THE EU-GEI HIGH RISK STUDY. Schizophrenia Bulletin, 2019, 45, S177-S178.  | 2.3 | 0         |
| 220 | T21. DEVELOPMENT OF PROTEOMIC PREDICTION MODELS FOR OUTCOMES IN THE CLINICAL HIGH RISK STATE AND PSYCHOTIC EXPERIENCES IN ADOLESCENCE: MACHINE LEARNING ANALYSES IN TWO NESTED CASE-CONTROL STUDIES. Schizophrenia Bulletin, 2020, 46, S238-S239. | 2.3 | 0         |
| 221 | T234. MEASURING PERSONAL RECOVERY IN PEOPLE WITH A PSYCHOTIC DISORDER BASED ON CHIME: A COMPARISON OF RELIABILITY AND VALIDITY PROPERTIES ON THREE VALIDATED MEASURES. Schizophrenia Bulletin, 2020, 46, S322-S322.                               | 2.3 | O         |
| 222 | M36. VIRTUAL REALITY COGNITIVE BEHAVIORAL THERAPY FOR PARANOID DELUSIONS. Schizophrenia Bulletin, 2020, 46, S148-S148.  | 2.3 | 0         |
| 223 | S175. CLINICAL OUTCOMES IN PEOPLE AT HIGH RISK FOR PSYCHOSIS RELATED TO INTERACTIONS BETWEEN POLYGENIC RISK SCORES AND CHILDHOOD ADVERSITY. Schizophrenia Bulletin, 2020, 46, S104-S104.  | 2.3 | O         |
| 224 | T103. THE EFFECT OF VIRTUAL REALITY COGNITIVE BEHAVIORAL THERAPY ON PARANOIA AND MOOD STATES. Schizophrenia Bulletin, 2020, 46, S270-S270.  | 2.3 | 0         |
| 225 | Letter to the Editor: In reply to the correspondence of Johannes Rentzsch, Kristin Koller and Golo Kronenberg. Psychological Medicine, 2016, 46, 3063-3064.   | 2.7 | 0         |
| 226 | Traumabehandlung bei Psychose. Trauma Und Gewalt, 2017, 11, 216-233.  | 0.1 | 0         |
| 227 | Positive Memory Training for the Treatment of Depression in Schizophrenia: A Randomised Controlled Trial. SSRN Electronic Journal, 0, , .   | 0.4 | 0         |
| 228 | Virtual Reality and Psychotic Disorders. Virtual Reality Technologies for Health and Clinical Applications, 2019, , 289-305.  | 0.8 | 0         |
| 229 | Cognitieve therapie bij auditief verbale hallucinaties en wanen. , 2019, , 335-356.   |     | 0         |
| 230 | 6.2 Cognitieve gedragstherapie., 2018,, 209-219.  |     | 0         |
| 231 | Controlling a stream of paranoia evoking events in a virtual reality environment. Studies in Health Technology and Informatics, 2014, 199, 55-60.   | 0.2 | 0         |
| 232 | Impact of smoking behavior on cognitive functioning in persons at risk for psychosis and healthy controls: A longitudinal study $\hat{a} \in CORRIGENDUM$ . European Psychiatry, 2022, 65, e17.   | 0.1 | 0         |
| 233 | The relationship between grey matter volume and clinical and functional outcomes in people at clinical high risk for psychosis. Schizophrenia Bulletin Open, 0, , .   | 0.9 | 0         |