Mark Van Der Gaag

List of Publications by Citations

Source: https://exaly.com/author-pdf/4083646/mark-van-der-gaag-publications-by-citations.pdf

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

234 7,928 49 82 g-index

274 9,525 4.4 ext. papers ext. citations avg, IF 6.14 L-index

#	Paper	IF	Citations
234	The five-factor model of the Positive and Negative Syndrome Scale II: a ten-fold cross-validation of a revised model. <i>Schizophrenia Research</i> , 2006 , 85, 280-7	3.6	387
233	Preventing a first episode of psychosis: meta-analysis of randomized controlled prevention trials of 12 month and longer-term follow-ups. <i>Schizophrenia Research</i> , 2013 , 149, 56-62	3.6	270
232	EPA guidance on the early detection of clinical high risk states of psychoses. <i>European Psychiatry</i> , 2015 , 30, 405-16	6	235
231	EPA guidance on the early intervention in clinical high risk states of psychoses. <i>European Psychiatry</i> , 2015 , 30, 388-404	6	233
230	Psychological interventions for psychosis: a meta-analysis of comparative outcome studies. <i>American Journal of Psychiatry</i> , 2014 , 171, 523-38	11.9	213
229	Prolonged exposure vs eye movement desensitization and reprocessing vs waiting list for posttraumatic stress disorder in patients with a psychotic disorder: a randomized clinical trial. <i>JAMA Psychiatry</i> , 2015 , 72, 259-67	14.5	198
228	The validity of the 16-item version of the Prodromal Questionnaire (PQ-16) to screen for ultra high risk of developing psychosis in the general help-seeking population. <i>Schizophrenia Bulletin</i> , 2012 , 38, 1288-96	1.3	191
227	Identifying gene-environment interactions in schizophrenia: contemporary challenges for integrated, large-scale investigations. <i>Schizophrenia Bulletin</i> , 2014 , 40, 729-36	1.3	186
226	Genetic Risk and Outcome of Psychosis (GROUP), a multi-site longitudinal cohort study focused on gene-environment interaction: objectives, sample characteristics, recruitment and assessment methods. <i>International Journal of Methods in Psychiatric Research</i> , 2012 , 21, 205-21	4.3	179
225	The effects of individually tailored formulation-based cognitive behavioural therapy in auditory hallucinations and delusions: a meta-analysis. <i>Schizophrenia Research</i> , 2014 , 156, 30-7	3.6	175
224	CRITICAL ANALYSIS OF THE CURRENT TREATMENT GUIDELINES FOR COMPLEX PTSD IN ADULTS. Depression and Anxiety, 2016 , 33, 359-69	8.4	153
223	Stigma moderates the associations of insight with depressed mood, low self-esteem, and low quality of life in patients with schizophrenia spectrum disorders. <i>Schizophrenia Research</i> , 2009 , 115, 363	3- 3 .6	144
222	The five-factor model of the Positive and Negative Syndrome Scale I: confirmatory factor analysis fails to confirm 25 published five-factor solutions. <i>Schizophrenia Research</i> , 2006 , 85, 273-9	3.6	140
221	Two subdomains of negative symptoms in psychotic disorders: established and confirmed in two large cohorts. <i>Journal of Psychiatric Research</i> , 2013 , 47, 718-25	5.2	135
220	Cognitive behavioral therapy for subjects at ultrahigh risk for developing psychosis: a randomized controlled clinical trial. <i>Schizophrenia Bulletin</i> , 2012 , 38, 1180-8	1.3	135
219	Trauma and recent life events in individuals at ultra high risk for psychosis: review and meta-analysis. <i>Schizophrenia Research</i> , 2015 , 161, 143-9	3.6	119
218	Cognitive-behavioural therapy for refractory psychotic symptoms of schizophrenia resistant to atypical antipsychotic medication. Randomised controlled trial. <i>British Journal of Psychiatry</i> , 2005 , 186, 324-30	5.4	119

217	Psychological therapies for auditory hallucinations (voices): current status and key directions for future research. <i>Schizophrenia Bulletin</i> , 2014 , 40 Suppl 4, S202-12	1.3	117
216	The treatment of hallucinations in schizophrenia spectrum disorders. <i>Schizophrenia Bulletin</i> , 2012 , 38, 704-14	1.3	117
215	The effectiveness of peer support groups in psychosis: a randomized controlled trial. <i>Acta Psychiatrica Scandinavica</i> , 2008 , 118, 64-72	6.5	111
214	Adapted cognitive-behavioural therapy required for targeting negative symptoms in schizophrenia: meta-analysis and meta-regression. <i>Psychological Medicine</i> , 2015 , 45, 453-65	6.9	96
213	Virtual-reality-based cognitive behavioural therapy versus waiting list control for paranoid ideation and social avoidance in patients with psychotic disorders: a single-blind randomised controlled trial. <i>Lancet Psychiatry,the</i> , 2018 , 5, 217-226	23.3	93
212	A neuropsychiatric model of biological and psychological processes in the remission of delusions and auditory hallucinations. <i>Schizophrenia Bulletin</i> , 2006 , 32 Suppl 1, S113-22	1.3	93
211	Is it time to develop a new cognitive therapy for psychosiscognitive remediation therapy (CRT)?. <i>Clinical Psychology Review</i> , 2001 , 21, 1227-56	10.8	93
21 0	A controlled trial of cognitive remediation in schizophrenia. Schizophrenia Bulletin, 2002, 28, 167-76	1.3	91
209	Treating trauma in psychosis with EMDR: a pilot study. <i>Journal of Behavior Therapy and Experimental Psychiatry</i> , 2012 , 43, 664-71	2.6	82
208	O6.5. INVESTIGATING VARIABLES FROM THE NAPLS RISK CALCULATOR FOR PSYCHOSIS IN THE EU-GEI HIGH RISK STUDY. <i>Schizophrenia Bulletin</i> , 2019 , 45, S177-S178	1.3	78
207	5.2 COGNITIVE BEHAVIORAL THERAPY FOR SOCIAL ACTIVATION IN RECENT-ONSET PSYCHOSIS: RANDOMIZED CONTROLLED TRIAL. <i>Schizophrenia Bulletin</i> , 2019 , 45, S93-S94	1.3	78
206	S175. CLINICAL OUTCOMES IN PEOPLE AT HIGH RISK FOR PSYCHOSIS RELATED TO INTERACTIONS BETWEEN POLYGENIC RISK SCORES AND CHILDHOOD ADVERSITY. <i>Schizophrenia Bulletin</i> , 2020 , 46, S1	04:310	14 ⁷⁸
205	T103. THE EFFECT OF VIRTUAL REALITY COGNITIVE BEHAVIORAL THERAPY ON PARANOIA AND MOOD STATES. <i>Schizophrenia Bulletin</i> , 2020 , 46, S270-S270	1.3	78
204	T21. DEVELOPMENT OF PROTEOMIC PREDICTION MODELS FOR OUTCOMES IN THE CLINICAL HIGH RISK STATE AND PSYCHOTIC EXPERIENCES IN ADOLESCENCE: MACHINE LEARNING ANALYSES IN TWO NESTED CASE-CONTROL STUDIES. <i>Schizophrenia Bulletin</i> , 2020 , 46, S238-S239	1.3	78
203	T234. MEASURING PERSONAL RECOVERY IN PEOPLE WITH A PSYCHOTIC DISORDER BASED ON CHIME: A COMPARISON OF RELIABILITY AND VALIDITY PROPERTIES ON THREE VALIDATED MEASURES. <i>Schizophrenia Bulletin</i> , 2020 , 46, S322-S322	1.3	78
202	M36. VIRTUAL REALITY COGNITIVE BEHAVIORAL THERAPY FOR PARANOID DELUSIONS. <i>Schizophrenia Bulletin</i> , 2020 , 46, S148-S148	1.3	78
201	A Meta-Analysis of Social Skills Training and Related Interventions for Psychosis. <i>Schizophrenia Bulletin</i> , 2018 , 44, 475-491	1.3	76
200	Whither the attenuated psychosis syndrome?. <i>Schizophrenia Bulletin</i> , 2012 , 38, 1130-4	1.3	74

199	Metacognitive reflection and insight therapy (MERIT) for patients with schizophrenia. <i>Psychological Medicine</i> , 2019 , 49, 303-313	6.9	74
198	Cannabis use and transition to psychosis in individuals at ultra-high risk: review and meta-analysis. <i>Psychological Medicine</i> , 2016 , 46, 673-81	6.9	73
197	Trauma-Focused Treatment in PTSD Patients With Psychosis: Symptom Exacerbation, Adverse Events, and Revictimization. <i>Schizophrenia Bulletin</i> , 2016 , 42, 693-702	1.3	70
196	Development of the Davos assessment of cognitive biases scale (DACOBS). <i>Schizophrenia Research</i> , 2013 , 144, 63-71	3.6	69
195	Treatment adherence therapy in people with psychotic disorders: randomised controlled trial. British Journal of Psychiatry, 2010 , 197, 448-55	5.4	69
194	Environmental Social Stress, Paranoia and Psychosis Liability: A Virtual Reality Study. <i>Schizophrenia Bulletin</i> , 2016 , 42, 1363-1371	1.3	69
193	Detection of people at risk of developing a first psychosis: comparison of two recruitment strategies. <i>Acta Psychiatrica Scandinavica</i> , 2012 , 126, 21-30	6.5	60
192	Early intervention in patients at ultra high risk of psychosis: benefits and risks. <i>Acta Psychiatrica Scandinavica</i> , 2009 , 119, 426-42	6.5	60
191	Prolonged exposure and EMDR for PTSD v. a PTSD waiting-list condition: effects on symptoms of psychosis, depression and social functioning in patients with chronic psychotic disorders. <i>Psychological Medicine</i> , 2016 , 46, 2411-21	6.9	54
190	Criminal victimisation in people with severe mental illness: a multi-site prevalence and incidence survey in the Netherlands. <i>PLoS ONE</i> , 2014 , 9, e91029	3.7	54
189	Cognitive Behavioral Therapy for negative symptoms (CBT-n) in psychotic disorders: a pilot study. Journal of Behavior Therapy and Experimental Psychiatry, 2013 , 44, 300-6	2.6	52
188	Competitive Memory Training for treating depression and rumination in depressed older adults: a randomized controlled trial. <i>Behaviour Research and Therapy</i> , 2011 , 49, 588-96	5.2	51
187	Predictive validity of the Trauma Screening Questionnaire in detecting post-traumatic stress disorder in patients with psychotic disorders. <i>British Journal of Psychiatry</i> , 2015 , 206, 408-16	5.4	50
186	Initial evaluation of the effects of competitive memory training (COMET) on depression in schizophrenia-spectrum patients with persistent auditory verbal hallucinations: a randomized controlled trial. <i>British Journal of Clinical Psychology</i> , 2012 , 51, 158-71	3.6	49
185	Metacognitive group training for schizophrenia spectrum patients with delusions: a randomized controlled trial. <i>Psychological Medicine</i> , 2014 , 44, 3025-35	6.9	49
184	Four-Year Follow-up of Cognitive Behavioral Therapy in Persons at Ultra-High Risk for Developing Psychosis: The Dutch Early Detection Intervention Evaluation (EDIE-NL) Trial. <i>Schizophrenia Bulletin</i> , 2016 , 42, 1243-52	1.3	48
183	Metacognitive training for schizophrenia spectrum patients: a meta-analysis on outcome studies. <i>Psychological Medicine</i> , 2016 , 46, 47-57	6.9	48
182	Evidence for a cognitive model of auditory hallucinations. <i>Journal of Nervous and Mental Disease</i> , 2003 , 191, 542-5	1.8	47

(2015-2014)

181	A manual-based individual therapy to improve metacognition in schizophrenia: protocol of a multi-center RCT. <i>BMC Psychiatry</i> , 2014 , 14, 27	4.2	44	
180	Depression and social anxiety in help-seeking patients with an ultra-high risk for developing psychosis. <i>Psychiatry Research</i> , 2013 , 209, 309-13	9.9	44	
179	Pathways to psychosis: help-seeking behavior in the prodromal phase. <i>Schizophrenia Research</i> , 2011 , 132, 213-9	3.6	44	
178	A single blind randomized controlled trial of cognitive behavioural therapy in a help-seeking population with an At Risk Mental State for psychosis: the Dutch Early Detection and Intervention Evaluation (EDIE-NL) trial. <i>Trials</i> , 2010 , 11, 30	2.8	43	
177	Cost-effectiveness of preventing first-episode psychosis in ultra-high-risk subjects: multi-centre randomized controlled trial. <i>Psychological Medicine</i> , 2015 , 45, 1435-46	6.9	41	
176	Cognitive disorders and negative symptoms as correlates of motivational deficits in psychotic patients. <i>Psychological Medicine</i> , 1994 , 24, 869-84	6.9	41	
175	Brave new worldsreview and update on virtual reality assessment and treatment in psychosis. <i>Schizophrenia Bulletin</i> , 2014 , 40, 1194-7	1.3	39	
174	Psychosocial functioning in first-episode psychosis and associations with neurocognition, social cognition, psychotic and affective symptoms. <i>Microbial Biotechnology</i> , 2017 , 11, 23-36	3.3	37	
173	Measuring empowerment among people with psychotic disorders: a comparison of three instruments. <i>Psychiatric Services</i> , 2008 , 59, 1338-42	3.3	37	
172	Fewer symptoms vs. more side-effects in schizophrenia? Opposing pathways between antipsychotic medication compliance and quality of life. <i>Schizophrenia Research</i> , 2009 , 113, 27-33	3.6	36	
171	Practical Implications of Metacognitively Oriented Psychotherapy in Psychosis: Findings From a Pilot Study. <i>Journal of Nervous and Mental Disease</i> , 2016 , 204, 713-6	1.8	34	
170	Virtual reality experiments linking social environment and psychosis: a pilot study. <i>Cyberpsychology, Behavior, and Social Networking</i> , 2014 , 17, 191-5	4.4	33	
169	A multi-site single blind clinical study to compare the effects of prolonged exposure, eye movement desensitization and reprocessing and waiting list on patients with a current diagnosis of psychosis and co morbid post traumatic stress disorder: study protocol for the randomized	2.8	33	
168	controlled trial Treating Trauma in Psychosis. <i>Trials</i> , 2013 , 14, 151 Creating a Supportive Environment: Peer Support Groups for Psychotic Disorders. <i>Schizophrenia Bulletin</i> , 2015 , 41, 1211-3	1.3	32	
167	Suicidality and hospitalisation in patients with borderline personality disorder who experience auditory verbal hallucinations. <i>European Psychiatry</i> , 2017 , 41, 47-52	6	31	
166	Hallucinations in borderline personality disorder: Prevalence, characteristics and associations with comorbid symptoms and disorders. <i>Scientific Reports</i> , 2017 , 7, 13920	4.9	30	
165	Psychotic symptoms, cognition and affect as predictors of psychosocial problems and functional change in first-episode psychosis. <i>Schizophrenia Research</i> , 2014 , 158, 113-9	3.6	30	
164	Childhood trauma and clinical outcome in patients at ultra-high risk of transition to psychosis. <i>Schizophrenia Research</i> , 2015 , 169, 193-198	3.6	28	

163	Anxiety Partially Mediates Cybersickness Symptoms in Immersive Virtual Reality Environments. <i>Cyberpsychology, Behavior, and Social Networking</i> , 2018 , 21, 187-193	4.4	28	
162	Effect of crisis plans on admissions and emergency visits: a randomized controlled trial. <i>PLoS ONE</i> , 2014 , 9, e91882	3.7	28	
161	Cognitive-behavioural therapy for persistent and recurrent psychosis in people with schizophrenia-spectrum disorder: cost-effectiveness analysis. <i>British Journal of Psychiatry</i> , 2011 , 198, 59-65, sup 1	5.4	28	
160	Long-term outcomes of trauma-focused treatment in psychosis. <i>British Journal of Psychiatry</i> , 2018 , 212, 180-182	5.4	26	
159	The dynamics of pAL2-1 homologous linear plasmids in Podospora anserina. <i>Molecular Genetics and Genomics</i> , 1998 , 258, 521-9		26	
158	Childhood trauma, psychosis liability and social stress reactivity: a virtual reality study. <i>Psychological Medicine</i> , 2016 , 46, 3339-3348	6.9	26	
157	Child Maltreatment and Clinical Outcome in Individuals at Ultra-High Risk for Psychosis in the EU-GEI High Risk Study. <i>Schizophrenia Bulletin</i> , 2018 , 44, 584-592	1.3	25	
156	Development of a group and family-based cognitive behavioural therapy program for youth at risk for psychosis. <i>Microbial Biotechnology</i> , 2016 , 10, 511-521	3.3	25	
155	REFLEX, a social-cognitive group treatment to improve insight in schizophrenia: study protocol of a multi-center RCT. <i>BMC Psychiatry</i> , 2011 , 11, 161	4.2	25	
154	The validity of the DSM-IV diagnostic classification system of non-affective psychoses. <i>Australian and New Zealand Journal of Psychiatry</i> , 2011 , 45, 1061-8	2.6	25	
153	Byssal detachment underestimates tolerance of mussels to toxic compounds. <i>Marine Pollution Bulletin</i> , 2005 , 50, 20-9	6.7	24	
152	Duration of untreated psychosis and the long-term course of schizophrenia. <i>European Psychiatry</i> , 2000 , 15, 264-7	6	23	
151	A randomised controlled trial of positive memory training for the treatment of depression within schizophrenia. <i>BMC Psychiatry</i> , 2015 , 15, 85	4.2	22	
150	Four-Year Cost-effectiveness of Cognitive Behavior Therapy for Preventing First-episode Psychosis: The Dutch Early Detection Intervention Evaluation (EDIE-NL) Trial. <i>Schizophrenia Bulletin</i> , 2017 , 43, 365	-3 ¹ 7 ² 4	22	
149	CBT for Those at Risk of a First Episode Psychosis		22	
148	Prescription of antipsychotic medication to patients at ultra high risk of developing psychosis. <i>International Clinical Psychopharmacology</i> , 2009 , 24, 223-8	2.2	20	
147	Working Alliance in Patients with Severe Mental Illness Who Need a Crisis Intervention Plan. <i>Community Mental Health Journal</i> , 2016 , 52, 102-8	2.1	19	
146	Cost-Effectiveness of Virtual Reality Cognitive Behavioral Therapy for Psychosis: Health-Economic Evaluation Within a Randomized Controlled Trial. <i>Journal of Medical Internet Research</i> , 2020 , 22, e17098	37.6	19	

145	Development of Proteomic Prediction Models for Transition to Psychotic Disorder in the Clinical High-Risk State and Psychotic Experiences in Adolescence. <i>JAMA Psychiatry</i> , 2021 , 78, 77-90	14.5	19
144	Individualized Prediction of Transition to Psychosis in 1,676 Individuals at Clinical High Risk: Development and Validation of a Multivariable Prediction Model Based on Individual Patient Data Meta-Analysis. <i>Frontiers in Psychiatry</i> , 2019 , 10, 345	5	18
143	Effectiveness of trauma-focused treatment for patients with psychosis with and without the dissociative subtype of post-traumatic stress disorder. <i>British Journal of Psychiatry</i> , 2016 , 209, 347-348	5.4	18
142	Economic aspects of peer support groups for psychosis. <i>Community Mental Health Journal</i> , 2011 , 47, 99-105	2.1	18
141	Factor analysis of the scale of prodromal symptoms: differentiating between negative and depression symptoms. <i>Psychopathology</i> , 2011 , 44, 379-85	3.4	18
140	Efficacy of Acceptance and Commitment Therapy in Daily Life (ACT-DL) in early psychosis: study protocol for a multi-centre randomized controlled trial. <i>Trials</i> , 2019 , 20, 769	2.8	18
139	The effect of mind-body and aerobic exercise on negative symptoms in schizophrenia: A meta-analysis. <i>Psychiatry Research</i> , 2019 , 279, 295-305	9.9	17
138	What Constitutes Sufficient Evidence for Case Formulation-Driven CBT for Psychosis? Cumulative Meta-analysis of the Effect on Hallucinations and Delusions. <i>Schizophrenia Bulletin</i> , 2020 ,	1.3	17
137	Social environments and interpersonal distance regulation in psychosis: A virtual reality study. <i>Schizophrenia Research</i> , 2018 , 192, 96-101	3.6	17
136	Metacognitive beliefs, beliefs about voices and affective symptoms in patients with severe auditory verbal hallucinations. <i>British Journal of Clinical Psychology</i> , 2013 , 52, 235-48	3.6	17
135	COMT Val(158)Met genotype and cannabis use in people with an At Risk Mental State for psychosis: Exploring Gene x Environment interactions. <i>Schizophrenia Research</i> , 2016 , 174, 24-28	3.6	17
134	CBT in the prevention of psychosis and other severe mental disorders in patients with an at risk mental state: A review and proposed next steps. <i>Schizophrenia Research</i> , 2019 , 203, 88-93	3.6	17
133	Symptom changes in five dimensions of the Positive and Negative Syndrome Scale in refractory psychosis. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2014 , 264, 673-82	5.1	16
132	The effect of childhood adversity on 4-year outcome in individuals at ultra high risk for psychosis in the Dutch Early Detection Intervention Evaluation (EDIE-NL) Trial. <i>Psychiatry Research</i> , 2017 , 247, 55-62	9.9	16
131	Understanding and Improving Treatment Adherence in Patients with Psychotic Disorders: A Review and a Proposed Intervention. <i>Current Psychiatry Reviews</i> , 2006 , 2, 487-494	0.9	16
130	Gender differences of patients at-risk for psychosis regarding symptomatology, drug use, comorbidity and functioning - Results from the EU-GEI study. <i>European Psychiatry</i> , 2019 , 59, 52-59	6	15
129	Screening for psychosis risk among adolescents in Child and Adolescent Mental Health Services: a description of the first step with the 16-item version of the Prodromal Questionnaire (PQ-16). <i>Microbial Biotechnology</i> , 2018 , 12, 669-676	3.3	15
128	The distribution of self-reported psychotic-like experiences in non-psychotic help-seeking mental health patients in the general population; a factor mixture analysis. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2014 , 49, 349-58	4.5	15

127	The Cognitive Biases Questionnaire for Psychosis (CBQ-P) and the Davos Assessment of Cognitive Biases (DACOBS): validation in a Flemish sample of psychotic patients and healthy controls. <i>Schizophrenia Research</i> , 2013 , 147, 310-4	3.6	15
126	The efficacy of CBT for severe mental illness and the challenge of dissemination in routine care. <i>World Psychiatry</i> , 2014 , 13, 257-8	14.4	15
125	Bridging the science-to-service gap in schizophrenia care in the Netherlands: the Schizophrenia Quality Improvement Collaborative. <i>International Journal for Quality in Health Care</i> , 2013 , 25, 626-32	1.9	15
124	Are social phobia and paranoia related, and which comes first?. <i>Psychosis</i> , 2009 , 1, 29-38	1.2	15
123	Recognition of early warning signs in patients with schizophrenia: a review of the literature. <i>International Journal of Mental Health Nursing</i> , 2004 , 13, 107-16	3.8	15
122	To continue or not to continue? Antipsychotic medication maintenance versus dose-reduction/discontinuation in first episode psychosis: HAMLETT, a pragmatic multicenter single-blind randomized controlled trial. <i>Trials</i> , 2020 , 21, 147	2.8	14
121	Self-esteem moderates affective and psychotic responses to social stress in psychosis: A virtual reality study. <i>Schizophrenia Research</i> , 2018 , 202, 80-85	3.6	14
120	High psychosis liability is associated with altered autonomic balance during exposure to Virtual Reality social stressors. <i>Schizophrenia Research</i> , 2017 , 184, 14-20	3.6	14
119	Depressive symptoms are associated with (sub)clinical psychotic symptoms in patients with non-affective psychotic disorder, siblings and healthy controls. <i>Psychological Medicine</i> , 2013 , 43, 747-56	6.9	14
118	Ethnicity and baseline symptomatology in patients with an At Risk Mental State for psychosis. <i>Psychological Medicine</i> , 2012 , 42, 247-56	6.9	14
117	The practice of early recognition and early intervention to prevent psychotic relapse in patients with schizophrenia: an exploratory study. Part 2. <i>Journal of Psychiatric and Mental Health Nursing</i> , 2002 , 9, 357-63	2.4	14
116	The practice of early recognition and early intervention to prevent psychotic relapse in patients with schizophrenia: an exploratory study. Part 1. <i>Journal of Psychiatric and Mental Health Nursing</i> , 2002 , 9, 347-55	2.4	14
115	Association of Adverse Outcomes With Emotion Processing and Its Neural Substrate in Individuals at Clinical High Risk for Psychosis. <i>JAMA Psychiatry</i> , 2020 , 77, 190-200	14.5	14
114	Reducing distress and improving social functioning in daily life in people with auditory verbal hallucinations: study protocol for the RemstemPrandomised controlled trial. <i>BMJ Open</i> , 2018 , 8, e0205.	3-7	13
113	Crisis plans facilitated by patient advocates are better than those drawn up by clinicians: results from an RCT. <i>Administration and Policy in Mental Health and Mental Health Services Research</i> , 2014 , 41, 220-7	3.1	13
112	An intervention study to prevent relapse in patients with schizophrenia. <i>Journal of Nursing Scholarship</i> , 2006 , 38, 42-9	3.6	13
111	The effect of thiopental on glutamate metabolism in mouse cerebellar astrocytes in vitro. <i>Neuroscience Letters</i> , 2001 , 304, 141-4	3.3	13
110	A network analysis of post-traumatic stress and psychosis symptoms. <i>Psychological Medicine</i> , 2021 , 51, 2485-2492	6.9	12

109	Recovery style predicts remission at one-year follow-up in outpatients with schizophrenia spectrum disorders. <i>Journal of Nervous and Mental Disease</i> , 2011 , 199, 295-300	1.8	12
108	Relapse prevention in patients with schizophrenia. Archives of Psychiatric Nursing, 2003, 17, 117-25	2.1	12
107	Development of a stage-dependent prognostic model to predict psychosis in ultra-high-risk patients seeking treatment for co-morbid psychiatric disorders. <i>Psychological Medicine</i> , 2016 , 46, 1839-5	5 ^{6.9}	12
106	Social-cognitive risk factors for violence in psychosis: A discriminant function analysis. <i>Psychiatry Research</i> , 2018 , 265, 93-99	9.9	11
105	Social cognitive group treatment for impaired insight in psychosis: A multicenter randomized controlled trial. <i>Schizophrenia Research</i> , 2019 , 206, 362-369	3.6	11
104	Health-economic benefits of treating trauma in psychosis. <i>Hgre Utbildning</i> , 2019 , 10, 1565032	5	10
103	Temperature dependent larval occurrence and spat settlement of the invasive brackish water bivalve Mytilopsis leucophaeata (Conrad, 1831) (Dreissenidae). <i>Journal of Sea Research</i> , 2014 , 87, 30-34	1.9	10
102	Why do patients with schizophrenia who have poor insight still take antipsychotics? Memory deficits as moderators between adherence belief and behavior. <i>Journal of Psychiatric Practice</i> , 2011 , 17, 320-9	1.3	10
101	Determinants of Completion and Use of Psychiatric Advance Statements in Mental Health Care in the Netherlands. <i>Psychiatric Services</i> , 2016 , 67, 858-63	3.3	10
100	Dysregulated Lipid Metabolism Precedes Onset of Psychosis. <i>Biological Psychiatry</i> , 2021 , 89, 288-297	7.9	10
99	Schizophrenia and antipsychotic medicationbetter adherence, better outcomes?. <i>Schizophrenia Research</i> , 2013 , 151, 296-7	3.6	9
98	Symptomatic and functional remission and its associations with quality of life in patients with psychotic disorder in Assertive Community Treatment teams. <i>Comprehensive Psychiatry</i> , 2012 , 53, 1174-	·8⁄0 ³	9
97	Metacognitieve training voor patiliten met een psychotische kwetsbaarheid. <i>Dth</i> , 2010 , 30, 263-279		9
96	Preventive interventions for individuals at ultra high risk for psychosis: An updated and extended meta-analysis. <i>Clinical Psychology Review</i> , 2021 , 86, 102005	10.8	9
95	The Prodromal Questionnaire: a case for IRT-based adaptive testing of psychotic experiences?. <i>International Journal of Methods in Psychiatric Research</i> , 2017 , 26,	4.3	8
94	Efficacy and Moderators of Cognitive Behavioural Therapy for Psychosis Versus Other Psychological Interventions: An Individual-Participant Data Meta-Analysis. <i>Frontiers in Psychiatry</i> , 2020 , 11, 402	5	8
93	Virtual reality based cognitive behavioral therapy for paranoia: Effects on mental states and the dynamics among them. <i>Schizophrenia Research</i> , 2020 , 222, 227-234	3.6	8
92	Sex differences in cognitive functioning of patients at-risk for psychosis and healthy controls: Results from the European Gene-Environment Interactions study. <i>European Psychiatry</i> , 2020 , 63, e25	6	8

91	Effect of virtual reality exposure therapy on social participation in people with a psychotic disorder (VRETp): study protocol for a randomized controlled trial. <i>Trials</i> , 2016 , 17, 25	2.8	8
90	Cognitive deficits and ethnicity: a cohort study of early psychosis patients in The Netherlands. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2013 , 48, 37-47	4.5	8
89	Cognitive behavioral therapy for social activation in recent-onset psychosis: Randomized controlled trial. <i>Journal of Consulting and Clinical Psychology</i> , 2019 , 87, 151-160	6.5	8
88	Evidence for preventive treatments in young patients at clinical high risk of psychosis: the need for context. <i>Lancet Psychiatry,the</i> , 2020 , 7, 378-380	23.3	8
87	A time-series network approach to auditory verbal hallucinations: Examining dynamic interactions using experience sampling methodology. <i>Schizophrenia Research</i> , 2020 , 215, 148-156	3.6	8
86	Exposing therapists to trauma-focused treatment in psychosis: effects on credibility, expected burden, and harm expectancies. <i>Hgre Utbildning</i> , 2016 , 7, 31712	5	8
85	The effect of dialectical behaviour therapy in autism spectrum patients with suicidality and/or self-destructive behaviour (DIASS): study protocol for a multicentre randomised controlled trial. <i>BMC Psychiatry</i> , 2020 , 20, 127	4.2	7
84	Measuring personal recovery in people with a psychotic disorder based on CHIME: A comparison of three validated measures. <i>International Journal of Mental Health Nursing</i> , 2020 , 29, 808-819	3.8	7
83	Emotion Recognition and Adverse Childhood Experiences in Individuals at Clinical High Risk of Psychosis. <i>Schizophrenia Bulletin</i> , 2020 , 46, 823-833	1.3	7
82	The influence of adjunctive treatment and metacognitive deficits in schizophrenia on the experience of work. <i>Schizophrenia Research</i> , 2014 , 157, 107-11	3.6	7
81	Negative Beliefs about Voices in Patients with Borderline Personality Disorder Are Associated with Distress: A Plea for Cognitive-Behavioural Therapy?. <i>Psychopathology</i> , 2017 , 50, 255-261	3.4	7
80	Self-reported Cognitive Biases Moderate the Associations Between Social Stress and Paranoid Ideation in a Virtual Reality Experimental Study. <i>Schizophrenia Bulletin</i> , 2018 , 44, 749-756	1.3	7
79	Does Treatment Adherence Therapy reduce expense of healthcare use in patients with psychotic disorders? Cost-minimization analysis in a randomized controlled trial. <i>Schizophrenia Research</i> , 2011 , 133, 47-53	3.6	7
78	Relapse prevention in patients with schizophrenia: the application of an intervention protocol in nursing practice. <i>Archives of Psychiatric Nursing</i> , 2003 , 17, 165-72	2.1	7
77	Clinical, cognitive and neuroanatomical associations of serum NMDAR autoantibodies in people at clinical high risk for psychosis. <i>Molecular Psychiatry</i> , 2021 , 26, 2590-2604	15.1	7
76	Cognitive functioning throughout adulthood and illness stages in individuals with psychotic disorders and their unaffected siblings. <i>Molecular Psychiatry</i> , 2021 , 26, 4529-4543	15.1	7
75	Letter to the Editor: Should we focus on quality or quantity in meta-analyses?. <i>Psychological Medicine</i> , 2016 , 46, 2003-5	6.9	6
74	Designing and Testing of a Health-Economic Markov Model for Prevention and Treatment of Early Psychosis. <i>Expert Review of Pharmacoeconomics and Outcomes Research</i> , 2020 , 20, 269-279	2.2	6

73	Peer support and skills training through an eating club for people with psychotic disorders: A feasibility study. <i>Journal of Behavior Therapy and Experimental Psychiatry</i> , 2019 , 64, 80-86	2.6	5
72	Psychosocial treatments for clinical high risk individuals. <i>Schizophrenia Bulletin</i> , 2015 , 41, 22	1.3	5
71	Predicting trauma-focused treatment outcome in psychosis. <i>Schizophrenia Research</i> , 2016 , 176, 239-244	1 3.6	5
70	The Effectiveness of Transdiagnostic Applications of Competitive Memory Training (COMET) on Low Self-Esteem and Comorbid Depression: A Meta-analysis of Randomized Controlled Trials. <i>Cognitive Therapy and Research</i> ,1	2.7	5
69	Functional recovery of individuals with serious mental illnesses: Development and testing of a new short instrument for routine outcome monitoring. <i>Psychiatric Rehabilitation Journal</i> , 2018 , 41, 341-350	2.4	5
68	Positive memory training for the treatment of depression in schizophrenia: A randomised controlled trial. <i>Behaviour Research and Therapy</i> , 2020 , 135, 103734	5.2	5
67	Pre-training inter-rater reliability of clinical instruments in an international psychosis research project. <i>Schizophrenia Research</i> , 2021 , 230, 104-107	3.6	5
66	Psychopathology, cognition and outcome in Dutch and immigrant first-episode psychosis patients. <i>Microbial Biotechnology</i> , 2019 , 13, 646-656	3.3	5
65	Economic evaluation of brief cognitive behavioural therapy for social activation in recent-onset psychosis. <i>PLoS ONE</i> , 2018 , 13, e0206236	3.7	5
64	Predicting therapy success from the outset: The moderating effect of insight into the illness on metacognitive psychotherapy outcome among persons with schizophrenia. <i>Clinical Psychology and Psychotherapy</i> , 2019 , 26, 650-660	2.9	4
63	Relationship between jumping to conclusions and clinical outcomes in people at clinical high-risk for psychosis. <i>Psychological Medicine</i> , 2020 , 1-9	6.9	4
62	Virtual reality to study responses to social environmental stressors in individuals with and without psychosis. <i>Studies in Health Technology and Informatics</i> , 2011 , 167, 86-91	0.5	4
61	Sensory processing deficiencies in patients with borderline personality disorder who experience auditory verbal hallucinations. <i>Psychiatry Research</i> , 2019 , 281, 112545	9.9	3
60	Efficacy of a transdiagnostic ecological momentary intervention for improving self-esteem (SELFIE) in youth exposed to childhood adversity: study protocol for a multi-centre randomized controlled trial		3
59	Detection of Domestic Violence and Abuse by Community Mental Health Teams Using the BRAVE Intervention: A Multicenter, Cluster Randomized Controlled Trial. <i>Journal of Interpersonal Violence</i> , 2021 , 8862605211004177	2.2	3
58	Clinical, societal and personal recovery in schizophrenia spectrum disorders across time: states and annual transitions <i>British Journal of Psychiatry</i> , 2021 , 219, 401-408	5.4	3
57	Stress reactivity as a putative mechanism linking childhood trauma with clinical outcomes in individuals at ultra-high-risk for psychosis: Findings from the EU-GEI High Risk Study. <i>Epidemiology and Psychiatric Sciences</i> , 2021 , 30, e40	5.1	3
56	The associations between migrant status and ethnicity and the identification of individuals at ultra-high risk for psychosis and transition to psychosis: a systematic review. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2021 , 56, 1923-1941	4.5	3

55	Response to The factor structure of complex posttraumatic stress disorder in traumatized refugees Higre Utbildning, 2017, 8, 1307078	5	2
54	Treating prolactinoma and psychosis: medication and cognitive behavioural therapy. <i>BMJ Case Reports</i> , 2011 , 2011,	0.9	2
53	Cognitieve gedragstherapie bij het voorkomen van ernstige psychiatrische klachten. <i>Psychopraktijk</i> , 2010 , 2, 7-10		2
52	Field testing attenuated psychosis syndrome criteria. <i>American Journal of Psychiatry</i> , 2011 , 168, 1221; author reply 1221	11.9	2
51	Rehabilitatie bij chronische schizofrenie: wat voegt cognitieveßunctietraining toe?. <i>Dth</i> , 1997 , 17, 160-1	68	2
50	Schizofrenie en posttraumatische stress-stoornis. <i>Dth</i> , 2002 , 22, 7-13		2
49	From Speech Illusions to Onset of Psychotic Disorder: Applying Network Analysis to an Experimental Measure of Aberrant Experiences. <i>Schizophrenia Bulletin Open</i> , 2020 , 1,	2.2	2
48	Obsessive-Compulsive Symptoms and Other Symptoms of the At-risk Mental State for Psychosis: A Network Perspective. <i>Schizophrenia Bulletin</i> , 2021 , 47, 1018-1028	1.3	2
47	Impact of Comorbid Affective Disorders on Longitudinal Clinical Outcomes in Individuals at Ultra-high Risk for Psychosis. <i>Schizophrenia Bulletin</i> , 2021 ,	1.3	2
46	Family-focused treatment improves attenuated psychotic symptoms, but does not differ from brief treatment in negative symptoms and social functioning in ultra high risk patients aged 12-35 years. <i>Evidence-Based Mental Health</i> , 2015 , 18, 59	11.1	1
45	S52. WORKING MECHANISMS OF VIRTUAL REALITY BASED CBT FOR PARANOIA: A RANDOMIZED CONTROLLED TRIAL EXAMINING COGNITIVE BIASES, SCHEMATIC BELIEFS AND SAFETY BEHAVIOR. <i>Schizophrenia Bulletin</i> , 2018 , 44, S344-S344	1.3	1
44	Ultrahigh risk for developing psychosis and psychotic personality organization. <i>Microbial Biotechnology</i> , 2019 , 13, 673-676	3.3	1
43	PREVENTIVE PSYCHOTHERAPY. Schizophrenia Research, 2014, 153, S42-S43	3.6	1
42	Preventie van psychose door cognitieve gedragstherapie van invaliderende intrusies. <i>Dth</i> , 2010 , 30, 247	7-262	1
41	Competitive Memory Training (COMET) bij een psychotische vrouw met auditieve hallucinaties. <i>Dth</i> , 2007 , 27, 76-83		1
40	De ontwikkeling van een biopsychisch model van psychose. <i>Tijdschrift Voor Psychotherapie</i> , 2008 , 34, 260-274		1
39	Psychodramatechnieken voor de behandeling van angst bij stemmen. <i>Dth</i> , 2006 , 26, 2-9		1
38	Assessing the Acceptability, Feasibility and Sustainability of an Intervention to Increase Detection of Domestic Violence and Abuse in Patients Suffering From Severe Mental Illness: A Qualitative Study. <i>Frontiers in Psychiatry</i> , 2020 , 11, 581031	5	1

37	Momentary Manifestations of Negative Symptoms as Predictors of Clinical Outcomes in People at High Risk for Psychosis: Experience Sampling Study. <i>JMIR Mental Health</i> , 2021 , 8, e30309	6	1
36	Dysregulated lipid metabolism precedes onset of psychosis in people at clinical high risk		1
35	Cognitive-Behavioral Therapy 2012 , 361-374		1
34	Changes in Posttraumatic Cognitions Mediate the Effects of Trauma-Focused Therapy on Paranoia. <i>Schizophrenia Bulletin Open</i> , 2020 , 1,	2.2	1
33	F50. METACOGNITIVE REFLECTION AND INSIGHT THERAPY: A MULTICENTER RANDOMIZED CONTROLLED TRIAL. <i>Schizophrenia Bulletin</i> , 2018 , 44, S238-S239	1.3	1
32	Voice-hearersRbeliefs about the causes of their voices. <i>Psychiatry Research</i> , 2021 , 302, 113997	9.9	1
31	Impact of smoking Behavior on cognitive functioning in persons at risk for psychosis and healthy controls: A longitudinal study. <i>European Psychiatry</i> , 2021 , 64, e60	6	1
30	Efficacy of a transdiagnostic ecological momentary intervention for improving self-esteem (SELFIE) in youth exposed to childhood adversity: study protocol for a multi-center randomized controlled trial. <i>Trials</i> , 2021 , 22, 641	2.8	1
29	Association between tobacco use and symptomatology in individuals at ultra-high risk to develop a psychosis: A longitudinal study. <i>Schizophrenia Research</i> , 2021 , 236, 48-53	3.6	1
28	Efficacy of Acceptance and Commitment Therapy in Daily Life in Early Psychosis: Results from the Multi-Center INTERACT Randomized Controlled Trial <i>Psychotherapy and Psychosomatics</i> , 2022 , 1-13	9.4	1
27	Characteristics and stability of hallucinations and delusions in patients with borderline personality disorder <i>Comprehensive Psychiatry</i> , 2021 , 113, 152290	7.3	О
26	Verbal memory performance predicts remission and functional outcome in people at clinical high-risk for psychosis <i>Schizophrenia Research: Cognition</i> , 2022 , 28, 100222	2.8	O
25	Linguistic dual tasking reduces emotionality, vividness and credibility of voice memories in voice-hearing individuals: Results from a controlled trial. <i>Schizophrenia Research</i> , 2020 , 216, 249-254	3.6	О
24	UP 店 : A Cohort Study on Recovery in Psychotic Disorder Patients: Design Protocol. <i>Frontiers in Psychiatry</i> , 2020 , 11, 609530	5	O
23	The association between migrant status and transition in an ultra-high risk for psychosis population. <i>Social Psychiatry and Psychiatric Epidemiology</i> , 2021 , 56, 943-952	4.5	О
22	Improving screening methods for psychosis in an adolescent help-seeking population using the Child Behavior Checklist (CBCL) and the Youth Self Report (YSR) versus the Prodromal Questionnaire -16 items version (PQ-16) Child and Adolescent Psychiatry and Mental Health, 2022,	6.8	O
21	Response to the letters of Dr Amos and Dr Preti and colleagues. <i>Schizophrenia Research</i> , 2014 , 153, 23	7-9 .6	
20	Positive symptoms in at-risk mental state: the importance of differentiating within the scope. <i>Microbial Biotechnology</i> , 2013 , 7, 100-1	3.3	

Fasen van psychose en het neurocognitief model. *Psychopraktijk*, **2012**, 4, 15-19

18	Reply: To PMID 22335365. <i>Acta Psychiatrica Scandinavica</i> , 2013 , 127, 332	6.5
17	Voorkomen is beter dan genezen. <i>GZ - Psychologie</i> , 2011 , 3, 12-17	O
16	ARE SOCIAL ANXIETY AND PARANOIA ASSOCIATED, AND WHICH COMES FIRST?. <i>Schizophrenia Research</i> , 2010 , 117, 317-318	3.6
15	Predictors of persistence of ultrahigh risk symptoms and predictors of transition into psychosis. European Psychiatry, 2011 , 26, 2086-2086	6
14	School performance in secondary education shows no decline before the onset of a first episode of psychosis in schizophrenia. <i>Journal of Mental Health</i> , 2003 , 12, 585-593	2.7
13	Zorgelijke en zorgeloze wanen. <i>Dth</i> , 2005 , 25, 90-94	
12	Placebo in een leertheoretisch perspectief. <i>Dth</i> , 2001 , 21, 156-159	
11	S50.01 Assessment of disablement in schizophrenia. <i>European Psychiatry</i> , 2000 , 15, 310s-310s	6
10	Voice phenomenology as a mirror of the past <i>Psychological Medicine</i> , 2022 , 1-9	6.9
9	6.2 Cognitieve gedragstherapie 2018 , 209-219	
8	Satisfaction with social connectedness as a predictor for positive and negative symptoms of psychosis: A PHAMOUS study. <i>Schizophrenia Research</i> , 2021 , 238, 121-127	3.6
7	Virtual Reality and Psychotic Disorders. Virtual Reality Technologies for Health and Clinical Applications, 2019 , 289-305	1.7
6	Cognitieve therapie bij auditief verbale hallucinaties en wanen 2019 , 335-356	
5	Letter to the Editor: In reply to the correspondence of Johannes Rentzsch, Kristin Koller and Golo Kronenberg. <i>Psychological Medicine</i> , 2016 , 46, 3063-3064	6.9
4	Traumabehandlung bei Psychose. <i>Trauma Und Gewalt</i> , 2017 , 11, 216-233	0.2
3	15 Cognitieve therapie bij psychose 2011 , 425-449	
2	Controlling a stream of paranoia evoking events in a virtual reality environment. <i>Studies in Health Technology and Informatics</i> , 2014 , 199, 55-60	0.5

Impact of smoking behavior on cognitive functioning in persons at risk for psychosis and healthy controls: A longitudinal study - CORRIGENDUM.. *European Psychiatry*, **2022**, 65, e17

6