Maria P Lemos

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4086182/publications.pdf

Version: 2024-02-01

12	1,002	5	8
papers	citations	h-index	g-index
16	16	16	2519
all docs	docs citations	times ranked	citing authors

#	Article	IF	CITATIONS
1	mRNA vaccination boosts cross-variant neutralizing antibodies elicited by SARS-CoV-2 infection. Science, 2021, 372, 1413-1418.	12.6	468
2	Longitudinal analysis shows durable and broad immune memory after SARS-CoV-2 infection with persisting antibody responses and memory B and Tâcells. Cell Reports Medicine, 2021, 2, 100354.	6.5	316
3	The Inner Foreskin of Healthy Males at Risk of HIV Infection Harbors Epithelial CD4+ CCR5+ Cells and Has Features of an Inflamed Epidermal Barrier. PLoS ONE, 2014, 9, e108954.	2.5	27
4	Rank-based two-sample tests for paired data with missing values. Biostatistics, 2018, 19, 281-294.	1.5	19
5	Rectal tissue and vaginal tissue from intravenous VRC01 recipients show protection against ex vivo HIV-1 challenge. Journal of Clinical Investigation, 2021, 131, .	8.2	17
6	Transient Peripheral Immune Activation follows Elective Sigmoidoscopy or Circumcision in a Cohort Study of MSM at Risk of HIV Infection. PLoS ONE, 2016, 11, e0160487.	2.5	6
7	Daily Vaginal Swabs and Mobile Phone Sex Report for Assessing HIV Virion Exposure Prospectively Among a Cohort of Young Sexually Active Women in South Africa (HVTN 915). Journal of Acquired Immune Deficiency Syndromes (1999), 2019, 81, e39-e48.	2.1	5
8	"You Are on the Right Track With the App:―Qualitative Analysis of Mobile Phone Use and User Feedback Regarding Mobile Phone Sexual Risk Assessments for HIV Prevention Research. Frontiers in Digital Health, 2021, 3, 576514.	2.8	3
9	Circulating CD4+ T-cells Are Transiently Activated Following Sigmoidoscopy or Circumcision in MSM at High Risk of HIV Infection. AIDS Research and Human Retroviruses, 2014, 30, A176-A176.	1.1	O
10	Response to Guo et al.'s Letter to the Editor. Biostatistics, 2019, 20, 363-365.	1.5	0
11	Rapid Collection of Human Rectal Secretions Using OriCol Devices Is Suitable for Measurement of Mucosal Ig without Blood Contamination. Journal of Immunology, 2020, 205, 2312-2320.	0.8	O
12	HIV-1 Nucleic Acids Identify Rectal HIV Exposures in Self-Collected Rectal Swabs, Whereas Y-Chromosome Single Tandem Repeat Mixtures Are Not Reliable Biomarkers of Condomless Receptive Anal Intercourse. Journal of Acquired Immune Deficiency Syndromes (1999), 2021, 88, 138-148.	2.1	0