

Olof Beck

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4093862/olof-beck-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

170
papers

5,384
citations

44
h-index

63
g-index

176
ext. papers

6,023
ext. citations

3.8
avg, IF

5.96
L-index

#	Paper	IF	Citations
170	Severe acute respiratory syndrome coronavirus 2 can be detected in exhaled aerosol sampled during a few minutes of breathing or coughing.. <i>Influenza and Other Respiratory Viruses</i> , 2022 ,	5.6	4
169	Multianalyte serology in home-sampled blood enables an unbiased assessment of the immune response against SARS-CoV-2. <i>Nature Communications</i> , 2021 , 12, 3695	17.4	11
168	Peanuts in the air - clinical and experimental studies. <i>Clinical and Experimental Allergy</i> , 2021 , 51, 585-593	4.1	6
167	Measurement of the alcohol biomarker phosphatidylethanol (PEth) in dried blood spots and venous blood-importance of inhibition of post-sampling formation from ethanol. <i>Analytical and Bioanalytical Chemistry</i> , 2021 , 413, 5601-5606	4.4	5
166	Hydromorphone and codeine concentrations in oral fluid specimens from patients receiving substitution therapy with Substitol (morphine sulfate). <i>Drug Testing and Analysis</i> , 2021 , 13, 1743-1748	3.5	
165	Impact of morphine dose on ticagrelor uptake and platelet inhibition in patients with ST-segment elevation myocardial infarction - A substudy from the prospective randomized MOVEMENT trial. <i>Thrombosis Update</i> , 2021 , 5, 100071	0.9	
164	Drug trends and harm related to new psychoactive substances (NPS) in Sweden from 2010 to 2016: Experiences from the STRIDA project. <i>PLoS ONE</i> , 2020 , 15, e0232038	3.7	9
163	Drugs in breath 2020 , 493-507		1
162	Pharmacokinetics of methylphenidate and ritalinic acid in plasma correlations with exhaled breath and oral fluid in healthy volunteers. <i>European Journal of Clinical Pharmacology</i> , 2020 , 76, 229-237	2.8	3
161	Drug trends and harm related to new psychoactive substances (NPS) in Sweden from 2010 to 2016: Experiences from the STRIDA project 2020 , 15, e0232038		
160	Drug trends and harm related to new psychoactive substances (NPS) in Sweden from 2010 to 2016: Experiences from the STRIDA project 2020 , 15, e0232038		
159	Drug trends and harm related to new psychoactive substances (NPS) in Sweden from 2010 to 2016: Experiences from the STRIDA project 2020 , 15, e0232038		
158	Drug trends and harm related to new psychoactive substances (NPS) in Sweden from 2010 to 2016: Experiences from the STRIDA project 2020 , 15, e0232038		
157	Drug trends and harm related to new psychoactive substances (NPS) in Sweden from 2010 to 2016: Experiences from the STRIDA project 2020 , 15, e0232038		
156	Drug trends and harm related to new psychoactive substances (NPS) in Sweden from 2010 to 2016: Experiences from the STRIDA project 2020 , 15, e0232038		
155	Dose-Response Characteristics of the Alcohol Biomarker Phosphatidylethanol (PEth)-A Study of Outpatients in Treatment for Reduced Drinking. <i>Alcohol and Alcoholism</i> , 2019 , 54, 567-573	3.5	15
154	First evaluation of the possibility of testing for drugged driving using exhaled breath sampling. <i>Traffic Injury Prevention</i> , 2019 , 20, 238-243	1.8	4

153	An Autonomous Microfluidic Device for Generating Volume-Defined Dried Plasma Spots. <i>Analytical Chemistry</i> , 2019 , 91, 7125-7130	7.8	9
152	Elimination Characteristics of the Alcohol Biomarker Phosphatidylethanol (PEth) in Blood during Alcohol Detoxification. <i>Alcohol and Alcoholism</i> , 2019 , 54, 251-257	3.5	22
151	Detection of heroin intake in patients in substitution treatment using oral fluid as specimen for drug testing. <i>Drug and Alcohol Dependence</i> , 2019 , 198, 136-139	4.9	3
150	Evaluation of a Volumetric Dried Blood Spot Card Using a Gravimetric Method and a Bioanalytical Method with Capillary Blood from 44 Volunteers. <i>Analytical Chemistry</i> , 2019 , 91, 5558-5565	7.8	11
149	Application of Liquid Chromatography Combined With High Resolution Mass Spectrometry for Urine Drug Testing 2019 , 321-332		2
148	Evaluation of an on-site test device for the heroin metabolite 6-acetylmorphine in urine. <i>Drug Testing and Analysis</i> , 2019 , 11, 536-540	3.5	2
147	The MOVEMENT Trial. <i>Journal of the American Heart Association</i> , 2019 , 8, e010152	6	11
146	Occurrence and time course of NPS benzodiazepines in Sweden - results from intoxication cases in the STRIDA project. <i>Clinical Toxicology</i> , 2019 , 57, 203-212	2.9	27
145	An analytically confirmed non-fatal intoxication by carfentanil in Sweden. <i>Clinical Toxicology</i> , 2019 , 57, 372-374	2.9	4
144	Drug abuse screening with exhaled breath and oral fluid in adults with substance use disorder. <i>Drug Testing and Analysis</i> , 2019 , 11, 27-32	3.5	11
143	Chemical synthesis, characterisation and in vitro and in vivo metabolism of the synthetic opioid MT-45 and its newly identified fluorinated analogue 2F-MT-45 with metabolite confirmation in urine samples from known drug users. <i>Forensic Toxicology</i> , 2018 , 36, 359-374	2.6	22
142	Evaluation of a new simple collection device for sampling of microparticles in exhaled breath. <i>Journal of Breath Research</i> , 2018 , 12, 036005	3.1	8
141	Study of measurement of the alcohol biomarker phosphatidylethanol (PEth) in dried blood spot (DBS) samples and application of a volumetric DBS device. <i>Clinica Chimica Acta</i> , 2018 , 479, 38-42	6.2	26
140	Detectability of fentanyl and designer fentanyls in urine by 3 commercial fentanyl immunoassays. <i>Drug Testing and Analysis</i> , 2018 , 10, 1297	3.5	24
139	Investigation of drug products received for analysis in the Swedish STRIDA project on new psychoactive substances. <i>Drug Testing and Analysis</i> , 2018 , 10, 340-349	3.5	17
138	European guidelines for workplace drug testing in oral fluid. <i>Drug Testing and Analysis</i> , 2018 , 10, 402-415	3.5	20
137	Human urinary metabolic patterns of the designer benzodiazepines flubromazolam and pyrazolam studied by liquid chromatography-high resolution mass spectrometry. <i>Drug Testing and Analysis</i> , 2018 , 10, 496-506	3.5	21
136	Intoxications in the STRIDA project involving a panorama of psychostimulant pyrovalerone derivatives, MDPV copycats. <i>Clinical Toxicology</i> , 2018 , 56, 256-263	2.9	32

135	A liquid chromatography and tandem mass spectrometry method to determine 28 non-volatile drugs of abuse in exhaled breath. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2018 , 148, 251-258	3.5	13
134	Acute Intoxications Involving Pyrrolidinobutiophenone (PBPP): Results from the Swedish STRIDA Project. <i>Journal of Medical Toxicology</i> , 2018 , 14, 265-271	2.6	12
133	Urine analysis of 28 designer benzodiazepines by liquid chromatography-high-resolution mass spectrometry. <i>Clinical Mass Spectrometry</i> , 2018 , 10, 25-32	1.9	8
132	Sevoflurane metabolite cross-react in the ethyl glucuronide DRUG-Immunoassay. <i>Drug Testing and Analysis</i> , 2018 , 10, 379-380	3.5	1
131	High-Yield Passive Plasma Filtration from Human Finger Prick Blood. <i>Analytical Chemistry</i> , 2018 , 90, 13392-13399	3.5	20
130	Comparison between dried blood spot and plasma sampling for therapeutic drug monitoring of antiepileptic drugs in children with epilepsy: A step towards home sampling. <i>Clinical Biochemistry</i> , 2017 , 50, 418-424	3.5	24
129	European guidelines for workplace drug testing in urine. <i>Drug Testing and Analysis</i> , 2017 , 9, 853-865	3.5	16
128	Endogenous and xenobiotic metabolic stability of primary human hepatocytes in long-term 3D spheroid cultures revealed by a combination of targeted and untargeted metabolomics. <i>FASEB Journal</i> , 2017 , 31, 2696-2708	0.9	90
127	Use of LC-HRMS in full scan-XIC mode for multi-analyte urine drug testing - a step towards a black-box solution?. <i>Journal of Mass Spectrometry</i> , 2017 , 52, 497-506	2.2	17
126	Oil-Fortified Maize Porridge Increases Absorption of Lumefantrine in Children with Uncomplicated Falciparum Malaria. <i>Basic and Clinical Pharmacology and Toxicology</i> , 2017 , 120, 457-465	3.1	4
125	Intoxications involving acrylfentanyl and other novel designer fentanyls - results from the Swedish STRIDA project. <i>Clinical Toxicology</i> , 2017 , 55, 589-599	2.9	77
124	Detectability of designer benzodiazepines in CEDIA, EMIT II Plus, HEIA, and KIMS II immunochemical screening assays. <i>Drug Testing and Analysis</i> , 2017 , 9, 640-645	3.5	55
123	Analytically Confirmed Intoxications Involving MDMB-CHMICA from the STRIDA Project. <i>Journal of Medical Toxicology</i> , 2017 , 13, 52-60	2.6	54
122	Cannabinoids in Exhaled Breath 2017 , 1018-1024		
121	Two techniques to sample non-volatiles in breath-exemplified by methadone. <i>Journal of Breath Research</i> , 2017 , 12, 016011	3.1	14
120	Experiences from using LC-MS/MS for analysis of immunosuppressive drugs in a TDM service. <i>Clinical Biochemistry</i> , 2016 , 49, 1024-31	3.5	12
119	Potential of Mass Spectrometry in Developing Clinical Laboratory Biomarkers of Nonvolatiles in Exhaled Breath. <i>Clinical Chemistry</i> , 2016 , 62, 84-91	5.5	35
118	Intoxications involving the fentanyl analogs acetylfentanyl, 4-methoxybutyrfentanyl and furanylfentanyl: results from the Swedish STRIDA project. <i>Clinical Toxicology</i> , 2016 , 54, 324-32	2.9	115

117	Identification of main human urinary metabolites of the designer nitrobenzodiazepines clonazolam, meclonazepam, and nifoxipam by nano-liquid chromatography-high-resolution mass spectrometry for drug testing purposes. <i>Analytical and Bioanalytical Chemistry</i> , 2016 , 408, 3571-91	4.4	50
116	Characterization of exhaled breath particles collected by an electret filter technique. <i>Journal of Breath Research</i> , 2016 , 10, 026001	3.1	23
115	Toxicity evaluation of Epyrrolidinovalerophenone (EPVP): results from intoxication cases within the STRIDA project. <i>Clinical Toxicology</i> , 2016 , 54, 568-75	2.9	41
114	Development and application of a multi-component LC-MS/MS method for determination of designer benzodiazepines in urine. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2016 , 1035, 104-110	3.2	42
113	Adverse events related to the new psychoactive substance 3-fluorophenmetrazine - results from the Swedish STRIDA project. <i>Clinical Toxicology</i> , 2016 , 54, 819-825	2.9	19
112	(9)-Tetrahydrocannabinol concentrations in exhaled breath and physiological effects following cannabis intake - A pilot study using illicit cannabis. <i>Clinical Biochemistry</i> , 2016 , 49, 1072-7	3.5	27
111	Opioid intoxications involving butyrfentanyl, 4-fluorobutyrfentanyl, and fentanyl from the Swedish STRIDA project. <i>Clinical Toxicology</i> , 2015 , 53, 609-17	2.9	120
110	Narcolepsy Treated with Racemic Amphetamine during Pregnancy and Breastfeeding. <i>Journal of Human Lactation</i> , 2015 , 31, 374-6	2.6	9
109	Intoxications by the dissociative new psychoactive substances diphenidine and methoxphenidine. <i>Clinical Toxicology</i> , 2015 , 53, 446-53	2.9	58
108	First report on the pharmacokinetics of tramadol and O-desmethyltramadol in exhaled breath compared to plasma and oral fluid after a single oral dose. <i>Biochemical Pharmacology</i> , 2015 , 98, 502-10	6	26
107	A disposable sampling device to collect volume-measured DBS directly from a fingerprick onto DBS paper. <i>Bioanalysis</i> , 2015 , 7, 2085-94	2.1	47
106	The effect of drying on the homogeneity of DBS. <i>Bioanalysis</i> , 2015 , 7, 1977-85	2.1	22
105	A LC-MS/MS method for therapeutic drug monitoring of carbamazepine, lamotrigine and valproic acid in DBS. <i>Bioanalysis</i> , 2015 , 7, 2031-9	2.1	25
104	Phencyclidine analog use in Sweden--intoxication cases involving 3-MeO-PCP and 4-MeO-PCP from the STRIDA project. <i>Clinical Toxicology</i> , 2015 , 53, 856-64	2.9	53
103	Intoxications involving MDPV in Sweden during 2010-2014: Results from the STRIDA project. <i>Clinical Toxicology</i> , 2015 , 53, 865-73	2.9	41
102	Measurement of Lung Phosphatidylcholines in Exhaled Breath Particles by a Convenient Collection Procedure. <i>Analytical Chemistry</i> , 2015 , 87, 11553-60	7.8	16
101	Methodologies for assessment of limit of detection and limit of identification using surface-enhanced Raman spectroscopy. <i>Sensors and Actuators B: Chemical</i> , 2015 , 207, 437-446	8.5	48
100	Clinical trial of a new technique for drugs of abuse testing: a new possible sampling technique. <i>Journal of Substance Abuse Treatment</i> , 2015 , 48, 132-6	4.2	19

99	The effects of the monoamine stabilizer (-)-OSU6162 on craving in alcohol dependent individuals: A human laboratory study. <i>European Neuropsychopharmacology</i> , 2015 , 25, 2240-51	1.2	20
98	Possible mechanism for inhibition of morphine formation from 6-acetylmorphine after intake of street heroin. <i>Forensic Science International</i> , 2015 , 252, 150-6	2.6	3
97	Phosphatidylethanol in breath: a possible noninvasive screening test for heavy alcohol consumption. <i>Clinical Chemistry</i> , 2015 , 61, 991-3	5.5	12
96	Characteristics of analytically confirmed 3-MMC-related intoxications from the Swedish STRIDA project. <i>Clinical Toxicology</i> , 2015 , 53, 46-53	2.9	60
95	Method validation and application of a liquid chromatography-tandem mass spectrometry method for drugs of abuse testing in exhaled breath. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2015 , 985, 189-96	3.2	25
94	Application of drug testing using exhaled breath for compliance monitoring of drug addicts in treatment. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2015 , 75, 156-61	2	15
93	Detectability of new psychoactive substances, Regal highs, in CEDIA, EMIT, and KIMS immunochemical screening assays for drugs of abuse. <i>Drug Testing and Analysis</i> , 2014 , 6, 492-9	3.5	31
92	Detection of new psychoactive substance use among emergency room patients: results from the Swedish STRIDA project. <i>Forensic Science International</i> , 2014 , 243, 23-9	2.6	95
91	Direct and efficient liquid chromatographic-tandem mass spectrometric method for opiates in urine drug testing - importance of 6-acetylmorphine and reduction of analytes. <i>Drug Testing and Analysis</i> , 2014 , 6, 317-24	3.5	16
90	Methylphenidate for attention deficit hyperactivity disorder and drug relapse in criminal offenders with substance dependence: a 24-week randomized placebo-controlled trial. <i>Addiction</i> , 2014 , 109, 440-9	4.6	98
89	Phospholipid removal combined with a semi-automated 96-well SPE application for determination of budesonide in human plasma with LC-MS/MS. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2014 , 970, 31-5	3.2	16
88	MT-45, a new psychoactive substance associated with hearing loss and unconsciousness. <i>Clinical Toxicology</i> , 2014 , 52, 901-4	2.9	132
87	On the monitoring of dabigatran treatment in "real life" patients with atrial fibrillation. <i>Thrombosis Research</i> , 2014 , 134, 783-9	8.2	42
86	Exhaled breath for drugs of abuse testing - evaluation in criminal justice settings. <i>Science and Justice - Journal of the Forensic Science Society</i> , 2014 , 54, 57-60	2	25
85	Methods for urine drug testing using one-step dilution and direct injection in combination with LC-MS/MS and LC-HRMS. <i>Bioanalysis</i> , 2014 , 6, 2229-44	2.1	24
84	Quantification of cocaine and metabolites in exhaled breath by liquid chromatography-high-resolution mass spectrometry following controlled administration of intravenous cocaine. <i>Analytical and Bioanalytical Chemistry</i> , 2014 , 406, 6213-23	4.4	23
83	Intoxications of the new psychoactive substance 5-(2-aminopropyl)indole (5-IT): a case series from the Swedish STRIDA project. <i>Clinical Toxicology</i> , 2014 , 52, 618-24	2.9	37
82	Determination of amphetamine and methylphenidate in exhaled breath of patients undergoing attention-deficit/hyperactivity disorder treatment. <i>Therapeutic Drug Monitoring</i> , 2014 , 36, 528-34	3.2	16

81	Laboratory and clinical evaluation of on-site urine drug testing. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2014 , 74, 681-6	2	16
80	Identification of novel psychoactive drug use in Sweden based on laboratory analysis--initial experiences from the STRIDA project. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2013 , 73, 400-6	2	67
79	Detection of drugs of abuse in exhaled breath using a device for rapid collection: comparison with plasma, urine and self-reporting in 47 drug users. <i>Journal of Breath Research</i> , 2013 , 7, 026006	3.1	58
78	Multicomponent LC-MS/MS screening method for detection of new psychoactive drugs, legal highs, in urine-experience from the Swedish population. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2013 , 930, 112-20	3.2	98
77	Cannabinoids in exhaled breath following controlled administration of smoked cannabis. <i>Clinical Chemistry</i> , 2013 , 59, 1780-9	5.5	68
76	Synthesis and bioanalytical evaluation of morphine-3-O-sulfate and morphine-6-O-sulfate in human urine and plasma using LC-MS/MS. <i>Journal of Separation Science</i> , 2012 , 35, 367-75	3.4	4
75	Evaluation of a direct high-capacity target screening approach for urine drug testing using liquid chromatography-time-of-flight mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2012 , 909, 6-13	3.2	27
74	Detection of drugs of abuse in exhaled breath from users following recovery from intoxication. <i>Journal of Analytical Toxicology</i> , 2012 , 36, 638-46	2.9	32
73	191 Morphine Premedication for Intubation in Preterm Infants - A Pharmacokinetic and Pharmacogenetic Report. <i>Archives of Disease in Childhood</i> , 2012 , 97, A55-A55	2.2	
72	Demonstration that methadone is being present in the exhaled breath aerosol fraction. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 2011 , 56, 1024-8	3.5	25
71	Detection of Δ^9 -tetrahydrocannabinol in exhaled breath collected from cannabis users. <i>Journal of Analytical Toxicology</i> , 2011 , 35, 541-4	2.9	47
70	Determination of methadone in exhaled breath condensate by liquid chromatography-tandem mass spectrometry. <i>Journal of Analytical Toxicology</i> , 2011 , 35, 129-33	2.9	35
69	Study on the sampling of methadone from exhaled breath. <i>Journal of Analytical Toxicology</i> , 2011 , 35, 257-63	2.9	35
68	Amphetamines detected in exhaled breath from drug addicts: A new possible method for drugs-of-abuse testing. <i>Journal of Analytical Toxicology</i> , 2010 , 34, 233-7	2.9	45
67	Sustained release methylphenidate for the treatment of ADHD in amphetamine abusers: a pilot study. <i>Drug and Alcohol Dependence</i> , 2010 , 108, 130-3	4.9	54
66	Acamprosate determinations in plasma and cerebrospinal fluid after multiple dosing measured by liquid chromatography-mass spectroscopy: a pharmacokinetic study in healthy volunteers. <i>Therapeutic Drug Monitoring</i> , 2010 , 32, 489-96	3.2	10
65	Method for determination of methadone in exhaled breath collected from subjects undergoing methadone maintenance treatment. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2010 , 878, 2255-9	3.2	31
64	Comparison of analytical approaches for liquid chromatography/mass spectrometry determination of the alcohol biomarker ethyl glucuronide in urine. <i>Rapid Communications in Mass Spectrometry</i> , 2010 , 24, 1737-43	2.2	27

63	The effects of acamprosate on alcohol-cue reactivity and alcohol priming in dependent patients: a randomized controlled trial. <i>Psychopharmacology</i> , 2009 , 205, 53-62	4.7	45
62	Unreliable alcohol testing in a shipping safety programme. <i>Forensic Science International</i> , 2009 , 189, e45-76	20	
61	Precise determination of glucose-d(2)/glucose ratio in human serum and plasma by APCI LC-MS/MS. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2009 , 69, 837-42	2	5
60	Detection times for urinary ethyl glucuronide and ethyl sulfate in heavy drinkers during alcohol detoxification. <i>Alcohol and Alcoholism</i> , 2009 , 44, 55-61	3.5	142
59	Plasma concentrations of lamotrigine and its 2-N-glucuronide metabolite during pregnancy in women with epilepsy. <i>Epilepsia</i> , 2008 , 49, 1075-80	6.4	80
58	Chapter 17 Analytical markers of acute and chronic alcohol consumption. <i>Handbook of Analytical Separations</i> , 2008 , 6, 567-588	0.7	2
57	Naltrexone attenuates the subjective effects of amphetamine in patients with amphetamine dependence. <i>Neuropsychopharmacology</i> , 2008 , 33, 1856-63	8.7	82
56	Naltrexone for the treatment of amphetamine dependence: a randomized, placebo-controlled trial. <i>American Journal of Psychiatry</i> , 2008 , 165, 1442-8	11.9	132
55	Development and clinical application of an LC-MS-MS method for mescaline in urine. <i>Journal of Analytical Toxicology</i> , 2008 , 32, 227-31	2.9	13
54	Direct injection LC-MS/MS method for identification and quantification of amphetamine, methamphetamine, 3,4-methylenedioxyamphetamine and 3,4-methylenedioxymethamphetamine in urine drug testing. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2008 , 871, 82-8	3.2	43
53	Accurate identification and quantification of 11-nor-delta(9)-tetrahydrocannabinol-9-carboxylic acid in urine drug testing: evaluation of a direct high efficiency liquid chromatographic-mass spectrometric method. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2008 , 871, 101-8	3.2	25
52	Evaluation of a new immunoassay for urinary ethyl glucuronide testing. <i>Alcohol and Alcoholism</i> , 2008 , 43, 46-8	3.5	81
51	Validation of direct injection electrospray LC-MS/MS for confirmation of opiates in urine drug testing. <i>Journal of Mass Spectrometry</i> , 2007 , 42, 881-9	2.2	70
50	Alcohol biomarker analysis: simultaneous determination of 5-hydroxytryptophol glucuronide and 5-hydroxyindoleacetic acid by direct injection of urine using ultra-performance liquid chromatography-tandem mass spectrometry. <i>Journal of Mass Spectrometry</i> , 2007 , 42, 940-9	2.2	23
49	Determination of glucuronidated 5-hydroxytryptophol (GTOL), a marker of recent alcohol intake, by ELISA technique. <i>Clinical Biochemistry</i> , 2007 , 40, 128-31	3.5	6
48	Biomarkers to disclose recent intake of alcohol: potential of 5-hydroxytryptophol glucuronide testing using new direct UPLC-tandem MS and ELISA methods. <i>Alcohol and Alcoholism</i> , 2007 , 42, 321-5	3.5	15
47	Electrospray LC-MS method with solid-phase extraction for accurate determination of morphine-, codeine-, and ethylmorphine-glucuronides and 6-acetylmorphine in urine. <i>Journal of Analytical Toxicology</i> , 2007 , 31, 81-6	2.9	27
46	Paradoxical results in urine drug testing for 6-acetylmorphine and total opiates: implications for best analytical strategy. <i>Journal of Analytical Toxicology</i> , 2006 , 30, 73-9	2.9	17

45	Determination of lamotrigine and its metabolites in human plasma by liquid chromatography-mass spectrometry. <i>Therapeutic Drug Monitoring</i> , 2006 , 28, 603-7	3.2	43
44	Determination of urinary 5-hydroxytryptophol glucuronide by liquid chromatography-mass spectrometry. <i>Journal of Chromatography B: Analytical Technologies in the Biomedical and Life Sciences</i> , 2005 , 816, 107-12	3.2	12
43	Application of direct urine LC-MS-MS analysis for screening of novel substances in drug abusers. <i>Journal of Analytical Toxicology</i> , 2005 , 29, 234-9	2.9	48
42	Vibration-induced hearing loss: comparison of the protective efficacy of antioxidants and neurotrophins. <i>Audiological Medicine</i> , 2004 , 2, 169-173		
41	Multicomponent screening for drugs of abuse: direct analysis of urine by LC-MS-MS. <i>Therapeutic Drug Monitoring</i> , 2004 , 26, 90-7	3.2	51
40	A comparison of two intensities of psychosocial intervention for alcohol dependent patients treated with acamprosate. <i>Alcohol and Alcoholism</i> , 2004 , 39, 251-5	3.5	15
39	5-hydroxytryptophol as a marker for recent alcohol intake. <i>Addiction</i> , 2003 , 98 Suppl 2, 63-72	4.6	36
38	Direct screening of urine for MDMA and MDA by liquid chromatography-tandem mass spectrometry. <i>Journal of Analytical Toxicology</i> , 2003 , 27, 15-9	2.9	32
37	Comparison of urinary excretion characteristics of ethanol and ethyl glucuronide. <i>Journal of Analytical Toxicology</i> , 2002 , 26, 201-4	2.9	170
36	Direct quantification of ethyl glucuronide in clinical urine samples by liquid chromatography-mass spectrometry. <i>Therapeutic Drug Monitoring</i> , 2002 , 24, 645-51	3.2	103
35	Frontline immunochromatographic device for on-site urine testing of amphetamines: laboratory validation using authentic specimens. <i>Annals of Clinical Biochemistry</i> , 2000 , 37 (Pt 2), 199-204	2.2	21
34	Comparison of urinary 5-hydroxytryptophol, breath ethanol, and self-report for detection of recent alcohol use during outpatient treatment: a study on methadone patients. <i>Drug and Alcohol Dependence</i> , 1999 , 56, 33-8	4.9	41
33	Concentration-dependent stimulation of intestinal phase III of migrating motor complex by circulating serotonin in humans. <i>Clinical Science</i> , 1998 , 94, 663-70	6.5	20
32	Natural killer cell activity and csf monoamine metabolites in suicide attempters. <i>Archives of Suicide Research</i> , 1997 , 3, 153-169	2.3	
31	Natural killer cell activity and CSF monoamine metabolites in suicide attempters. <i>Archives of Suicide Research</i> , 1997 , 3, 153-169	2.3	
30	Urine and plasma pharmacokinetics of codeine in healthy volunteers: implications for drugs-of-abuse testing. <i>Journal of Analytical Toxicology</i> , 1996 , 20, 541-6	2.9	28
29	Evaluation of clinical assays for measuring high-dose methotrexate in plasma. <i>Clinical Chemistry</i> , 1996 , 42, 39-44	5.5	61
28	Laboratory testing for recent alcohol consumption: comparison of ethanol, methanol, and 5-hydroxytryptophol. <i>Clinical Chemistry</i> , 1996 , 42, 618-624	5.5	72

27	Liquid chromatographic determination of plasma lamotrigine in pediatric samples. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1996 , 14, 755-8	3.5	28
26	Determination of 5-hydroxytryptophol in urine by high-performance liquid chromatography: application of a new post-column derivatization method with fluorometric detection. <i>Journal of Pharmaceutical and Biomedical Analysis</i> , 1995 , 13, 651-4	3.5	7
25	5-Hydroxytryptophol conjugation in man: influence of alcohol consumption and altered serotonin turnover. <i>Life Sciences</i> , 1995 , 56, 1529-34	6.8	17
24	Changes in serotonin metabolism during treatment with the aldehyde dehydrogenase inhibitors disulfiram and cyanamide. <i>Basic and Clinical Pharmacology and Toxicology</i> , 1995 , 77, 323-6		24
23	Distinguishing Ingested Ethanol from Microbial Formation by Analysis of Urinary 5-Hydroxytryptophol and 5-Hydroxyindoleacetic Acid. <i>Journal of Forensic Sciences</i> , 1995 , 40, 13768J	1.8	32
22	Chloroquine reduces the bioavailability of methotrexate in patients with rheumatoid arthritis. A possible mechanism of reduced hepatotoxicity. <i>Arthritis and Rheumatism</i> , 1994 , 37, 830-3		34
21	Evaluation of analytical procedures for urinary codeine and morphine measurements. <i>Journal of Analytical Toxicology</i> , 1994 , 18, 129-33	2.9	27
20	The use of 5-hydroxytryptophol as an alcohol intake marker. <i>Alcohol and Alcoholism Supplement</i> , 1994 , 2, 497-502		7
19	On the accurate determination of serotonin in human plasma. <i>Biochemical and Biophysical Research Communications</i> , 1993 , 196, 260-6	3.4	54
18	Time course of ethanol-induced changes in serotonin metabolism. <i>Life Sciences</i> , 1993 , 53, 847-55	6.8	55
17	Detection of relapses in alcohol-dependent patients: comparison of carbohydrate-deficient transferrin in serum, 5-hydroxytryptophol in urine, and self-reports. <i>Alcoholism: Clinical and Experimental Research</i> , 1993 , 17, 703-8	3.7	44
16	Urinary 5-hydroxytryptophol: a possible marker of recent alcohol consumption. <i>Alcoholism: Clinical and Experimental Research</i> , 1992 , 16, 281-5	3.7	68
15	Urinary excretion of 5-hydroxyindole-3-acetic acid and 5-hydroxytryptophol after oral loading with serotonin. <i>Life Sciences</i> , 1992 , 50, 1207-13	6.8	46
14	Urinary 5HTOL/5HIAA as biochemical marker of postmortem ethanol synthesis. <i>Lancet, The</i> , 1992 , 340, 1159	4.0	19
13	Detection of Benzodiazepine Intake in Therapeutic Doses by Immunoanalysis of Urine: Two Techniques Evaluated and Modified for Improved Performance. <i>Clinical Chemistry</i> , 1992 , 38, 271-275	5.5	52
12	Determination of urinary 5-hydroxytryptophol by high-performance liquid chromatography with electrochemical detection. <i>Biomedical Applications</i> , 1992 , 579, 340-5		20
11	Detection of benzodiazepine intake in therapeutic doses by immunoanalysis of urine: two techniques evaluated and modified for improved performance. <i>Clinical Chemistry</i> , 1992 , 38, 271-5	5.5	4
10	Digoxin-verapamil interaction: reduction of biliary but not renal digoxin clearance in humans. <i>Clinical Pharmacology and Therapeutics</i> , 1991 , 49, 256-62	6.1	69

9	Determination of urinary 5-hydroxyindole-3-acetic acid by high-performance liquid chromatography with electrochemical detection and direct sample injection. <i>Analytical Biochemistry</i> , 1991 , 196, 170-3	3.1	52
8	Importance of creatinine analyses of urine when screening for abused drugs. <i>Clinical Chemistry</i> , 1991 , 37, 1927-1931	5.5	88
7	Monitoring of plasma methadone: intercorrelation between immunoassay and gas chromatography-mass spectrometry. <i>Therapeutic Drug Monitoring</i> , 1990 , 12, 473-7	3.2	1
6	5-hydroxytryptophol in the cerebrospinal fluid and urine of alcoholics and healthy subjects. <i>Naunyn-Schmiedeberg's Archives of Pharmacology</i> , 1982 , 321, 293-7	3.4	47
5	Synthesis of six specifically deuterated indoles of biological interest. <i>Journal of Labelled Compounds and Radiopharmaceuticals</i> , 1980 , 17, 411-419	1.9	15
4	5-methoxytryptamine in the human pineal gland: identification and quantitation by mass fragmentography. <i>Journal of Neurochemistry</i> , 1979 , 32, 1853-5	6	8
3	The synthesis of 5-hydroxyindole-3-(α -D2-acetic acid) and α -D4-5-hydroxytryptamine. <i>Journal of Labelled Compounds</i> , 1975 , 11, 57-61		21
2	Study of Innervation, Sensory Neuropeptides, and Serotonin in Murine Contact Allergic Skin		2
1	A translational multiplex serology approach to profile the prevalence of anti-SARS-CoV-2 antibodies in home-sampled blood		6