
Lin Jiangkai

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4108738/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Curcumin Improves Human Umbilical Cord-Derived Mesenchymal Stem Cell Survival via ERK1/2 Signaling and Promotes Motor Outcomes After Spinal Cord Injury. Cellular and Molecular Neurobiology, 2022, 42, 1241-1252.	3.3	8
2	Microsurgical sealing for symptomatic sacral Tarlov cysts: a series of 265 cases. Journal of Neurosurgery: Spine, 2022, 37, 905-913.	1.7	6
3	Treatment of symptomatic Chiari I malformation by "all-factors-surgery": a report of 194 cases. European Spine Journal, 2021, 30, 1615-1622.	2.2	3
4	Risk factor analysis for progressive spinal deformity after resection of intracanal tumors─ a retrospective study of 272 cases. BMC Neurology, 2020, 20, 34.	1.8	9
5	T lymphocytes infiltration promotes blood-brain barrier injury after experimental intracerebral hemorrhage. Brain Research, 2017, 1670, 96-105.	2.2	29
6	Curcumin attenuates blood-brain barrier disruption after subarachnoid hemorrhage in mice. Journal of Surgical Research, 2017, 207, 85-91.	1.6	36
7	Curcumin inhibits glial scar formation by suppressing astrocyte-induced inflammation and fibrosis in vitro and in vivo. Brain Research, 2017, 1655, 90-103.	2.2	56
8	M2 microglia promotes neurogenesis and oligodendrogenesis from neural stem/progenitor cells via the PPARÎ ³ signaling pathway. Oncotarget, 2017, 8, 19855-19865.	1.8	78
9	Curcumin reduces brain-infiltrating T lymphocytes after intracerebral hemorrhage in mice. Neuroscience Letters, 2016, 620, 74-82.	2.1	28
10	Poly-L-ornithine promotes preferred differentiation of neural stem/progenitor cells via ERK signalling pathway. Scientific Reports, 2015, 5, 15535.	3.3	65
11	Curcumin improves neural function after spinal cord injury by the joint inhibition of the intracellular and extracellular components of glial scar. Journal of Surgical Research, 2015, 195, 235-245.	1.6	52
12	Antisense vimentin cDNA combined with chondroitinase ABC promotes axon regeneration and functional recovery following spinal cord injury in rats. Neuroscience Letters, 2015, 590, 74-79.	2.1	11
13	Curcumin increased the differentiation rate of neurons in neural stem cells via wnt signaling inÂvitro study. Journal of Surgical Research, 2014, 192, 298-304.	1.6	42
14	Complement C5a is detrimental to histological and functional locomotor recovery after spinal cord injury in mice. Neurobiology of Disease, 2014, 66, 74-82.	4.4	39
15	Use of suction to treat intramedullary spinal cysticercosis. BMJ Case Reports, 2010, 2010, bcr0420091755-bcr0420091755.	0.5	4
16	Antisense vimentin cDNA combined with chondroitinase ABC reduces glial scar and cystic cavity formation following spinal cord injury in rats. Biochemical and Biophysical Research Communications, 2008, 377, 562-566.	2.1	25
17	Effect of Vimentin on Reactive Gliosis:In VitroandIn VivoAnalysis. Journal of Neurotrauma, 2004, 21, 1671-1682.	3.4	22
18	Intraparenchymal schwannoma of the medulla oblongata. Journal of Neurosurgery, 2003, 98, 621-624.	1.6	17