## Anna Prats

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4112889/publications.pdf Version: 2024-02-01



ΔΝΙΝΙΑ ΔΠΑΤΟ

#	Article	IF	CITATIONS
1	Effects of integrase inhibitor-based antiretroviral therapy on brain outcomes according to time since acquisition of HIV-1 infection. Scientific Reports, 2021, 11, 11289.	3.3	8
2	NEU Screen Shows High Accuracy in Detecting Cognitive Impairment in Older Persons Living With HIV. Journal of the Association of Nurses in AIDS Care, 2019, 30, 35-41.	1.0	4
3	Determining the optimal features in freezing of gait detection through a single waist accelerometer in home environments. Pattern Recognition Letters, 2018, 105, 135-143.	4.2	43
4	Deep learning for freezing of gait detection in Parkinson's disease patients in their homes using a waist-worn inertial measurement unit. Knowledge-Based Systems, 2018, 139, 119-131.	7.1	151
5	A "HOLTER―for Parkinson's disease: Validation of the ability to detect on-off states using the REMPARK system. Gait and Posture, 2018, 59, 1-6.	1.4	46
6	A Waist-Worn Inertial Measurement Unit for Long-Term Monitoring of Parkinson's Disease Patients. Sensors, 2017, 17, 827.	3.8	34
7	Transdermal rivastigmine for HIV-associated cognitive impairment: A randomized pilot study. PLoS ONE, 2017, 12, e0182547.	2.5	8
8	Deep Learning for Detecting Freezing of Gait Episodes in Parkinson's Disease Based on Accelerometers. Lecture Notes in Computer Science, 2017, , 344-355.	1.3	17
9	Home detection of freezing of gait using support vector machines through a single waist-worn triaxial accelerometer. PLoS ONE, 2017, 12, e0171764.	2.5	135
10	Classification Models for Neurocognitive Impairment in HIV Infection Based on Demographic and Clinical Variables. PLoS ONE, 2014, 9, e107625.	2.5	19
11	A Brief and Feasible Paper-Based Method to Screen for Neurocognitive Impairment in HIV-Infected Patients. Journal of Acquired Immune Deficiency Syndromes (1999), 2013, 63, 585-592.	2.1	26
12	Virological Efficacy in Cerebrospinal Fluid and Neurocognitive Status in Patients with Long-Term Monotherapy Based on Lopinavir/Ritonavir: An Exploratory Study. PLoS ONE, 2013, 8, e70201.	2.5	22
13	Interruptions of antiretroviral therapy in human immunodeficiency virus infection: are they detrimental to neurocognitive functioning?. Journal of NeuroVirology, 2010, 16, 208-218.	2.1	13
14	Nadir CD4 Cell Count Predicts Neurocognitive Impairment in HIV-Infected Patients. AIDS Research and Human Retroviruses, 2008, 24, 1301-1307.	1.1	87