

Giovanni Fanelli

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4116841/publications.pdf>

Version: 2024-02-01

28
papers

892
citations

566801

15
h-index

500791

28
g-index

28
all docs

28
docs citations

28
times ranked

1113
citing authors

#	ARTICLE	IF	CITATIONS
1	Effect of temperature and duration of immersion on the stability of prepared feeds in echinoculture. <i>Journal of Applied Aquaculture</i> , 2021, 33, 150-164.	0.7	4
2	Strategies for Successful Scallops Spat Collection on Artificial Collectors in the Taranto Gulf (Mediterranean Sea). <i>Water (Switzerland)</i> , 2021, 13, 462.	1.2	2
3	Estimation of Growth Parameters of the Black Scallop <i>Mimachlamys varia</i> in the Gulf of Taranto (Ionian Sea, Southern Italy). <i>Water (Switzerland)</i> , 2020, 12, 3342.	1.2	7
4	Comparative Characteristics of Percentage Edibility, Condition Index, Biochemical Constituents and Lipids Nutritional Quality Indices of Wild and Farmed Scallops (<i>Flexopecten glaber</i>). <i>Water (Switzerland)</i> , 2020, 12, 1777.	1.2	8
5	Bioactive fatty acids in seafood from Ionian Sea and relation to dietary recommendations. <i>International Journal of Food Sciences and Nutrition</i> , 2020, 71, 693-705.	1.3	17
6	Seasonal changes of commercial traits, proximate and fatty acid compositions of the scallop <i>Flexopecten glaber</i> from the Mediterranean Sea (Southern Italy). <i>PeerJ</i> , 2019, 7, e5810.	0.9	21
7	Can Different Body Tissues of Two Sea Cucumbers Supply a Fair Amount of Omega 3 for Health Benefit?. <i>Journal of Aquatic Food Product Technology</i> , 2019, 28, 821-836.	0.6	6
8	Effect of formulated diets on the proximate composition and fatty acid profiles of sea urchin <i>Paracentrotus lividus</i> gonad. <i>Aquaculture International</i> , 2018, 26, 185-202.	1.1	26
9	Bioactive fatty acids of three commercial scallop species. <i>International Journal of Food Properties</i> , 2018, 21, 519-532.	1.3	14
10	Influence of a prepared diet and a macroalga (<i>Ulva</i> sp.) on the growth, nutritional and sensory qualities of gonads of the sea urchin <i>Paracentrotus lividus</i> . <i>Aquaculture</i> , 2018, 493, 240-250.	1.7	41
11	The recruitment of scallops (and beyond) by two different artificial collectors (Gulf of Taranto, Italy). <i>Journal of Shellfish Research</i> , 2018, 37, 109-112.	0.9	12
12	New Mediterranean Biodiversity Records (April 2015). <i>Mediterranean Marine Science</i> , 2015, 16, 266.	0.6	25
13	Rearing experiences of the polychaete <i>Sabella spallanzanii</i> in the Gulf of Taranto (Mediterranean Sea, Italy). <i>Journal of Shellfish Research</i> , 2015, 34, 109-112.	1.1	9
14	Destructive standard squares or low-impact visually driven collection? A comparison of methods for quantitative samplings of benthic hydrozoans. <i>Italian Journal of Zoology</i> , 2013, 80, 424-436.	0.6	6
15	Hydrodynamism and its influence on the reproductive condition of the edible sea urchin <i>Paracentrotus lividus</i> . <i>Marine Environmental Research</i> , 2013, 85, 29-33.	1.1	23
16	Leading role of the sea urchin <i>Arbacia lixula</i> in maintaining the barren state in southwestern Mediterranean. <i>Marine Biology</i> , 2011, 158, 2505-2513.	0.7	77
17	<i>Sabella spallanzanii</i> filter-feeding on bacterial community: Ecological implications and applications. <i>Marine Environmental Research</i> , 2006, 61, 74-92.	1.1	59
18	Experimental co-culture of low food-chain organisms, <i>Sabella spallanzanii</i> (Polychaeta, Sabellidae) and <i>Cladophora prolifera</i> (Chlorophyta, Cladophorales), in Porto Cesareo area (Mediterranean Sea). <i>Aquaculture Research</i> , 2006, 37, 966-974.	0.9	17

#	ARTICLE	IF	CITATIONS
19	From biodiversity and ecosystem functioning to the roots of ecological complexity. <i>Ecological Complexity</i> , 2004, 1, 101-109.	1.4	26
20	First record of the alien gastropod <i>Melibe fimbriata</i> (Opisthobranchia: Tethyidae) in the Taranto seas (Mediterranean Sea). <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2004, 84, 1067-1068.	0.4	6
21	Plankton biodiversity around the Salento Peninsula (South East Italy): an integrated water/sediment approach. <i>Scientia Marina</i> , 2004, 68, 85-102.	0.3	34
22	Coastal fish indicate human-induced changes in the Mediterranean littoral. <i>Marine Environmental Research</i> , 2002, 53, 77-94.	1.1	76
23	Variability of species' roles in marine communities: change of paradigms for conservation priorities. <i>Marine Biology</i> , 2002, 140, 1067-1074.	0.7	112
24	Plankton-Derived Resting Stages in Marine Coastal Sediments along the Salento Peninsula (Apulia), Tj ETQq0 0 0 rgBT /Overlock 10 Tf 5	0.4	15
25	Bioturbation behaviour in two Mediterranean polychaetes. <i>Journal of the Marine Biological Association of the United Kingdom</i> , 2001, 81, 341-342.	0.4	3
26	Spatial variability and human disturbance in shallow subtidal hard substrate assemblages: a regional approach. <i>Marine Ecology - Progress Series</i> , 2001, 212, 1-12.	0.9	115
27	The continuity of living matter and the discontinuities of its constituents: do plankton and benthos really exist?. <i>Trends in Ecology and Evolution</i> , 1996, 11, 177-180.	4.2	129
28	Reply from F. Boero et al.. <i>Trends in Ecology and Evolution</i> , 1996, 11, 472.	4.2	2