
Hong Lin

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/411788/publications.pdf Version: 2024-02-01



HONCLIN

#	Article	IF	CITATIONS
1	High fatigue and its associations with health and work related factors among female medical personnel at 54 hospitals in Zhuhai, China. Psychology, Health and Medicine, 2018, 23, 304-316.	2.4	28
2	The association between dietary protein intake and colorectal cancer risk: a meta-analysis. World Journal of Surgical Oncology, 2017, 15, 169.	1.9	19
3	High Serum Phosphorus Level Is Associated with Left Ventricular Diastolic Dysfunction in Peritoneal Dialysis Patients. PLoS ONE, 2016, 11, e0163659.	2.5	8
4	Erectile dysfunction is associated with subclinical carotid vascular disease in young men lacking widely-known risk factors. Asian Journal of Andrology, 2018, 20, 400.	1.6	8
5	Left atrial strain associated with alterations in cardiac diastolic function in patients with end-stage renal disease. International Journal of Cardiovascular Imaging, 2019, 35, 1803-1810.	1.5	6
6	Myocardial contrast echocardiography in the diagnosis of postoperative takotsubo myocardiopathy: case report and literature review. BMC Cardiovascular Disorders, 2019, 19, 9.	1.7	6
7	Serum apolipoprotein B is inversely associated with eccentric left ventricular hypertrophy in peritoneal dialysis patients. International Urology and Nephrology, 2018, 50, 155-165.	1.4	5
8	Serum Prealbumin and Echocardiography Parameters Predict Mortality in Peritoneal Dialysis Patients. Kidney and Blood Pressure Research, 2020, 45, 671-685.	2.0	4
9	Assessment of Myocardial Dysfunction by Three-Dimensional Echocardiography Combined With Myocardial Contrast Echocardiography in Type 2 Diabetes Mellitus. Frontiers in Cardiovascular Medicine, 2021, 8, 677990.	2.4	4
10	Diabetes attenuated ageâ€related aortic root dilatation in endâ€stage renal disease patients receiving peritoneal dialysis. Journal of Diabetes Investigation, 2019, 10, 1550-1557.	2.4	2
11	Subclinical endothelial dysfunction and low-grade inflammation play roles in the development of erectile dysfunction in young man with low risk of coronary heart disease. Heart, 2011, 97, A242-A242.	2.9	1
12	Benign pericardial schwannoma with massive pericardial effusion in a Chinese man: A case report. Echocardiography, 2019, 36, 1944-1946.	0.9	1
13	GENDER DIFFERENCES IN VASCULAR ENDOTHELIAL FUNCTION AND CAROTID INTIMA MEDIA THICKNESS BY FRAMINGHAM RISK SCORE. Heart, 2012, 98, E150.1-E150.	2.9	0
14	GW24-e2277â€Cardiac Diastolic Dysfunction may Hold Promise as Markers of Cardiotoxicity during Daunorubicin Treatment in Acute Nonlymphocytic Leukemia Patients. Heart, 2013, 99, A273.2-A273.	2.9	0
15	GW24-e1282â€A noninvasive sizing method to choose fitted atrial septal defect occluder by transthoracic echocardiography in adults with secundum atrial septal defects. Heart, 2013, 99, A227.3-A228.	2.9	0
16	Left atrial diameter is associated with target organ damage in patients with type 2 diabetes mellitus. International Journal of Diabetes in Developing Countries, 2019, 39, 355-361.	0.8	0
17	Mediastinal Tuberculoma Mimicking Malignant Cardiac Tumor: A Case Report. Cardiovascular Innovations and Applications, 2021, 5, 297-300.	0.3	0