

El-Hassane Nazih

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4122880/publications.pdf>

Version: 2024-02-01

27
papers

597
citations

759233

12
h-index

642732

23
g-index

27
all docs

27
docs citations

27
times ranked

1066
citing authors

#	ARTICLE	IF	CITATIONS
1	Intestinal Proportion of <i>Blautia</i> sp. is Associated with Clinical Stage and Histoprognostic Grade in Patients with Early-Stage Breast Cancer. <i>Nutrition and Cancer</i> , 2017, 69, 267-275.	2.0	124
2	Lithocholic bile acid inhibits lipogenesis and induces apoptosis in breast cancer cells. <i>Cellular Oncology (Dordrecht)</i> , 2018, 41, 13-24.	4.4	80
3	LXR agonists and ABCG1-dependent cholesterol efflux in MCF-7 breast cancer cells: relation to proliferation and apoptosis. <i>Anticancer Research</i> , 2012, 32, 3007-13.	1.1	64
4	Cholesterol, Oxysterols and LXRs in Breast Cancer Pathophysiology. <i>International Journal of Molecular Sciences</i> , 2020, 21, 1356.	4.1	42
5	Faecal Microbiota Composition Varies between Patients with Breast Cancer and Healthy Women: A Comparative Case-Control Study. <i>Nutrients</i> , 2021, 13, 2705.	4.1	32
6	Butyrate stimulates ApoA-IV-containing lipoprotein secretion in differentiated Caco-2 cells: Role in cholesterol efflux. <i>Journal of Cellular Biochemistry</i> , 2001, 83, 230-238.	2.6	31
7	Metabolomics-Driven Discovery of Meroterpenoids from a Mussel-Derived <i>Penicillium ubiquestum</i> . <i>Journal of Natural Products</i> , 2018, 81, 2501-2511.	3.0	31
8	Preventive Effects of the Marine Microalga <i>Phaeodactylum tricornutum</i> , Used as a Food Supplement, on Risk Factors Associated with Metabolic Syndrome in Wistar Rats. <i>Nutrients</i> , 2019, 11, 1069.	4.1	25
9	The Marine-Derived Fungus <i>Clonostachys rosea</i> , Source of a Rare Conjugated 4-Me-6E,8E-hexadecadienoic Acid Reducing Viability of MCF-7 Breast Cancer Cells and Gene Expression of Lipogenic Enzymes. <i>Marine Drugs</i> , 2015, 13, 4934-4948.	4.6	20
10	Macrophage apolipoprotein E and proliferation of MCF-7 breast cancer cells: role of LXR. <i>Anticancer Research</i> , 2013, 33, 3783-9.	1.1	19
11	Microalgae in Human Health. , 2018, , 211-226.		15
12	The Marine Microalga, <i>Tisochrysis lutea</i> , Protects against Metabolic Disorders Associated with Metabolic Syndrome and Obesity. <i>Nutrients</i> , 2021, 13, 430.	4.1	15
13	Steroids from Marine-Derived Fungi: Evaluation of Antiproliferative and Antimicrobial Activities of <i>Eburicol</i> . <i>Marine Drugs</i> , 2019, 17, 372.	4.6	14
14	4-cholesten-3-one decreases breast cancer cell viability and alters membrane raft-localized EGFR expression by reducing lipogenesis and enhancing LXR-dependent cholesterol transporters. <i>Lipids in Health and Disease</i> , 2019, 18, 168.	3.0	13
15	LXR Activation Down-regulates Lipid Raft Markers FLOT2 and DHH5 in MCF-7 Breast Cancer Cells. <i>Anticancer Research</i> , 2017, 37, 4067-4073.	1.1	11
16	Effect of Carotenoids from <i>Phaeodactylum tricornutum</i> on Palmitate-Treated HepG2 Cells. <i>Molecules</i> , 2020, 25, 2845.	3.8	10
17	Effects of selenium supplementation on expression of SEPP1 in mRNA and protein levels in subjects with and without metabolic syndrome suffering from coronary artery disease: Selenogene study a double-blind randomized controlled trial. <i>Journal of Cellular Biochemistry</i> , 2018, 119, 8282-8289.	2.6	9
18	Unusual sterolic mixture, and 24-isopropylcholesterol, from the sponge <i>Ciocalypta</i> sp. reduce cholesterol uptake and basolateral secretion in Caco-2 cells. <i>Journal of Cellular Biochemistry</i> , 2009, 106, 659-665.	2.6	8

#	ARTICLE	IF	CITATIONS
19	C25 steroids from the marine mussel-derived fungus <i>Penicillium ubiquestum</i> MMS330. <i>Phytochemistry Letters</i> , 2019, 34, 18-24.	1.2	6
20	Detection of ergosterol using liquid chromatography/electrospray ionization mass spectrometry: Investigation of unusual in-source reactions. <i>Rapid Communications in Mass Spectrometry</i> , 2020, 34, e8780.	1.5	6
21	Anti-proliferative and Pro-apoptotic Effect of Dichloromethane Extract of <i>Octopus vulgaris</i> By-Products on Human Breast Cancer Cell Lines. <i>Waste and Biomass Valorization</i> , 2015, 6, 237-242.	3.4	5
22	Link between Omega 3 Fatty Acids Carried by Lipoproteins and Breast Cancer Severity. <i>Nutrients</i> , 2022, 14, 2461.	4.1	4
23	Nicotinic Acid Accelerates HDL Cholesteryl Ester Turnover in Obese Insulin-Resistant Dogs. <i>PLoS ONE</i> , 2015, 10, e0136934.	2.5	3
24	High-Resolution Mass Spectrometry Unravels a Broad Range of Bioactive Lipid Species in <i>Octopus cyanea</i> and <i>Loligo</i> sp. By-products from Southwestern Madagascar. <i>Waste and Biomass Valorization</i> , 2018, 9, 1787-1793.	3.4	3
25	Basolateral Secretion from Caco-2 Cells Pretreated with Fecal Waters from Breast Cancer Patients Affects MCF7 Cell Viability. <i>Nutrients</i> , 2021, 13, 31.	4.1	3
26	Circulating HDL and Non-HDL Associated Apolipoproteins and Breast Cancer Severity. <i>Journal of Clinical Medicine</i> , 2022, 11, 1345.	2.4	3
27	Acetone Fraction of the Red Marine Alga Reduces the Expression of Bcl-2 Anti-apoptotic Marker and Flotillin-2 Lipid Raft Marker in MCF-7 Breast Cancer Cells. <i>Iranian Journal of Pharmaceutical Research</i> , 2020, 19, 321-330.	0.5	1