

# Francisco Javier Ollero

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4139442/publications.pdf>

Version: 2024-02-01

55  
papers

2,709  
citations

168829

31  
h-index

206121

51  
g-index

57  
all docs

57  
docs citations

57  
times ranked

2878  
citing authors

#	ARTICLE	IF	CITATIONS
1	Secondary Metabolites of <i>Rhizobium tropici</i> CIAT 899 Added to <i>Bradyrhizobium</i> spp. Inoculant Promote Soybean Growth and Increase Yield. <i>Journal of Soil Science and Plant Nutrition</i> , 2021, 21, 3354-3366.	1.7	1
2	Brief history of biofertilizers in Brazil: from conventional approaches to new biotechnological solutions. <i>Brazilian Journal of Microbiology</i> , 2021, 52, 2215-2232.	0.8	14
3	Plant Growth-Promoting Rhizobacteria Modulate the Concentration of Bioactive Compounds in Tomato Fruits. <i>Separations</i> , 2021, 8, 223.	1.1	2
4	OnfD, an AraC-Type Transcriptional Regulator Encoded by <i>Rhizobium tropici</i> CIAT 899 and Involved in Nod Factor Synthesis and Symbiosis. <i>Applied and Environmental Microbiology</i> , 2020, 86, .	1.4	8
5	The <i>Sinorhizobium fredii</i> HH103 type III secretion system effector NopC blocks nodulation with <i>Lotus japonicus</i> Gifu. <i>Journal of Experimental Botany</i> , 2020, 71, 6043-6056.	2.4	21
6	The non-flavonoid inducible <i>nodA3</i> and the flavonoid regulated <i>nodA1</i> genes of <i>Rhizobium tropici</i> CIAT 899 guarantee nod factor production and nodulation of different host legumes. <i>Plant and Soil</i> , 2019, 440, 185-200.	1.8	9
7	Osmotic stress activates <i>nif</i> and <i>fix</i> genes and induces the <i>Rhizobium tropici</i> CIAT 899 Nod factor production via NodD2 by up-regulation of the <i>nodA2</i> operon and the <i>nodA3</i> gene. <i>PLoS ONE</i> , 2019, 14, e0213298.	1.1	19
8	GunA of <i>Sinorhizobium</i> (Ensifer) <i>fredii</i> HH103 is a T3SS-secreted cellulase that differentially affects symbiosis with cowpea and soybean. <i>Plant and Soil</i> , 2019, 435, 15-26.	1.8	14
9	Revealing the roles of <i>y4wF</i> and <i>tidC</i> genes in <i>Rhizobium tropici</i> CIAT 899: biosynthesis of indolic compounds and impact on symbiotic properties. <i>Archives of Microbiology</i> , 2019, 201, 171-183.	1.0	13
10	Regulation of <i>hsnT</i> , <i>nodF</i> and <i>nodE</i> genes in <i>Rhizobium tropici</i> CIAT 899 and their roles in the synthesis of Nod factors and in the symbiosis. <i>Microbiology (United Kingdom)</i> , 2019, 165, 990-1000.	0.7	4
11	Structure of surface polysaccharides from <i>Aeromonas</i> sp. AMG272, a plant-growth promoting rhizobacterium isolated from rice rhizosphere. <i>Carbohydrate Research</i> , 2018, 462, 1-6.	1.1	7
12	Revealing strategies of quorum sensing in <i>Azospirillum brasilense</i> strains Ab-V5 and Ab-V6. <i>Archives of Microbiology</i> , 2018, 200, 47-56.	1.0	46
13	Transcriptomic Studies of the Effect of nod Gene-Inducing Molecules in Rhizobia: Different Weapons, One Purpose. <i>Genes</i> , 2018, 9, 1.	1.0	120
14	Co-inoculation of maize with <i>Azospirillum brasilense</i> and <i>Rhizobium tropici</i> as a strategy to mitigate salinity stress. <i>Functional Plant Biology</i> , 2018, 45, 328.	1.1	105
15	Antioxidant activity and induction of mechanisms of resistance to stresses related to the inoculation with <i>Azospirillum brasilense</i> . <i>Archives of Microbiology</i> , 2018, 200, 1191-1203.	1.0	34
16	The <i>Sinorhizobium</i> (Ensifer) <i>fredii</i> HH103 Nodulation Outer Protein NopI Is a Determinant for Efficient Nodulation of Soybean and Cowpea Plants. <i>Applied and Environmental Microbiology</i> , 2017, 83, .	1.4	43
17	Phytohormones and induction of plant-stress tolerance and defense genes by seed and foliar inoculation with <i>Azospirillum brasilense</i> cells and metabolites promote maize growth. <i>AMB Express</i> , 2017, 7, 153.	1.4	140
18	The <i>Rhizobium tropici</i> CIAT 899 NodD2 protein regulates the production of Nod factors under salt stress in a flavonoid-independent manner. <i>Scientific Reports</i> , 2017, 7, 46712.	1.6	30

#	ARTICLE	IF	CITATIONS
19	Genome of <i>Rhizobium leucaenae</i> strains CFN 299T and CPAO 29.8: searching for genes related to a successful symbiotic performance under stressful conditions. <i>BMC Genomics</i> , 2016, 17, 534.	1.2	13
20	RNA-seq analysis of the <i>Rhizobium tropici</i> CIAT 899 transcriptome shows similarities in the activation patterns of symbiotic genes in the presence of apigenin and salt. <i>BMC Genomics</i> , 2016, 17, 198.	1.2	42
21	NrcR, a New Transcriptional Regulator of <i>Rhizobium tropici</i> CIAT 899 Involved in the Legume Root-Nodule Symbiosis. <i>PLoS ONE</i> , 2016, 11, e0154029.	1.1	17
22	The <i>Sinorhizobium</i> ( <i>Ensifer</i> ) <i>fredii</i> HH103 Type 3 Secretion System Suppresses Early Defense Responses to Effectively Nodulate Soybean. <i>Molecular Plant-Microbe Interactions</i> , 2015, 28, 790-799.	1.4	38
23	Opening the "black box" of nodD3, nodD4 and nodD5 genes of <i>Rhizobium tropici</i> strain CIAT 899. <i>BMC Genomics</i> , 2015, 16, 864.	1.2	37
24	NopC Is a <i>Rhizobium</i> -Specific Type 3 Secretion System Effector Secreted by <i>Sinorhizobium</i> ( <i>Ensifer</i> ) <i>fredii</i> HH103. <i>PLoS ONE</i> , 2015, 10, e0142866.	1.1	54
25	Maize growth promotion by inoculation with <i>Azospirillum brasilense</i> and metabolites of <i>Rhizobium tropici</i> enriched on lipo-chitooligosaccharides (LCOs). <i>AMB Express</i> , 2015, 5, 71.	1.4	59
26	Regulatory nodD1 and nodD2 genes of <i>Rhizobium tropici</i> strain CIAT 899 and their roles in the early stages of molecular signaling and host-legume nodulation. <i>BMC Genomics</i> , 2015, 16, 251.	1.2	38
27	The Symbiotic Biofilm of <i>Sinorhizobium fredii</i> SMH12, Necessary for Successful Colonization and Symbiosis of <i>Glycine max</i> cv Osumi, Is Regulated by Quorum Sensing Systems and Inducing Flavonoids via NodD1. <i>PLoS ONE</i> , 2014, 9, e105901.	1.1	50
28	Plant growth promotion in cereal and leguminous agricultural important plants: From microorganism capacities to crop production. <i>Microbiological Research</i> , 2014, 169, 325-336.	2.5	504
29	Rice and bean AHL-mimic quorum-sensing signals specifically interfere with the capacity to form biofilms by plant-associated bacteria. <i>Research in Microbiology</i> , 2013, 164, 749-760.	1.0	70
30	Changes in flavonoids secreted by <i>Phaseolus vulgaris</i> roots in the presence of salt and the plant growth-promoting rhizobacterium <i>Chryseobacterium balustinum</i> . <i>Applied Soil Ecology</i> , 2012, 57, 31-38.	2.1	43
31	Genomic basis of broad host range and environmental adaptability of <i>Rhizobium tropici</i> CIAT 899 and <i>Rhizobium</i> sp. PRF 81 which are used in inoculants for common bean ( <i>Phaseolus vulgaris</i> L.). <i>BMC Genomics</i> , 2012, 13, 735.	1.2	118
32	Nodulation-gene-inducing flavonoids increase overall production of autoinducers and expression of N-acyl homoserine lactone synthesis genes in rhizobia. <i>Research in Microbiology</i> , 2011, 162, 715-723.	1.0	58
33	Effect of the presence of the plant growth promoting rhizobacterium (PGPR) <i>Chryseobacterium balustinum</i> Aur9 and salt stress in the pattern of flavonoids exuded by soybean roots. <i>Plant and Soil</i> , 2010, 328, 483-493.	1.8	129
34	The Absence of Nops Secretion in <i>Sinorhizobium fredii</i> HH103 Increases <i>GmPR1</i> Expression in Williams Soybean. <i>Molecular Plant-Microbe Interactions</i> , 2009, 22, 1445-1454.	1.4	65
35	<i>Sinorhizobium fredii</i> HH103 cgs Mutants Are Unable to Nodulate Determinate- and Indeterminate Nodule-Forming Legumes and Overproduce an Altered EPS. <i>Molecular Plant-Microbe Interactions</i> , 2009, 22, 575-588.	1.4	34
36	Regulation and symbiotic significance of nodulation outer proteins secretion in <i>Sinorhizobium fredii</i> HH103. <i>Microbiology (United Kingdom)</i> , 2008, 154, 1825-1836.	0.7	67

#	ARTICLE	IF	CITATIONS
37	NopM and NopD Are Rhizobial Nodulation Outer Proteins: Identification Using LC-MALDI and LC-ESI with a Monolithic Capillary Column. <i>Journal of Proteome Research</i> , 2007, 6, 1029-1037.	1.8	80
38	<i>Sinorhizobium fredii</i> HH103 Mutants Affected in Capsular Polysaccharide (KPS) are Impaired for Nodulation with Soybean and <i>Cajanus cajan</i> . <i>Molecular Plant-Microbe Interactions</i> , 2006, 19, 43-52.	1.4	61
39	Inactivation of the <i>Sinorhizobium fredii</i> HH103 <i>rhcJ</i> gene abolishes nodulation outer proteins (Nops) secretion and decreases the symbiotic capacity with soybean. <i>International Microbiology</i> , 2006, 9, 125-33.	1.1	52
40	The effect of FITA mutations on the symbiotic properties of <i>Sinorhizobium fredii</i> varies in a chromosomal-background-dependent manner. <i>Archives of Microbiology</i> , 2004, 181, 144-154.	1.0	35
41	NolR Regulates Diverse Symbiotic Signals of <i>Sinorhizobium fredii</i> HH103. <i>Molecular Plant-Microbe Interactions</i> , 2004, 17, 676-685.	1.4	58
42	Field assessment and genetic stability of <i>Sinorhizobium fredii</i> strain SMH12 for commercial soybean inoculants. <i>European Journal of Agronomy</i> , 2003, 19, 299-309.	1.9	23
43	A Catalogue of Molecular, Physiological and Symbiotic Properties of Soybean-Nodulating Rhizobial Strains from Different Soybean Cropping Areas of China. <i>Systematic and Applied Microbiology</i> , 2003, 26, 453-465.	1.2	21
44	Alfalfa nodulation by <i>Sinorhizobium fredii</i> does not require sulfated Nod-factors. <i>Functional Plant Biology</i> , 2003, 30, 1219.	1.1	7
45	<i>Sinorhizobium fredii</i> HH103 Has a Truncated <i>nolO</i> Gene Due to a -1 Frameshift Mutation That Is Conserved Among Other Geographically Distant <i>S. fredii</i> Strains. <i>Molecular Plant-Microbe Interactions</i> , 2002, 15, 150-159.	1.4	36
46	Soils of the Chinese Hubei Province Show a Very High Diversity of <i>Sinorhizobium fredii</i> Strains. <i>Systematic and Applied Microbiology</i> , 2002, 25, 592-602.	1.2	38
47	Effect of pH and soybean cultivars on the quantitative analyses of soybean rhizobia populations. <i>Journal of Biotechnology</i> , 2001, 91, 243-255.	1.9	58
48	Determination of the chemical structure of the capsular polysaccharide of strain B33, a fast-growing soya bean-nodulating bacterium isolated from an arid region of China. <i>Biochemical Journal</i> , 2001, 357, 505.	1.7	12
49	Determination of the chemical structure of the capsular polysaccharide of strain B33, a fast-growing soya bean-nodulating bacterium isolated from an arid region of China. <i>Biochemical Journal</i> , 2001, 357, 505-511.	1.7	18
50	Mutation in GDP-Fucose Synthesis Genes of <i>Sinorhizobium fredii</i> Alters Nod Factors and Significantly Decreases Competitiveness to Nodulate Soybeans. <i>Molecular Plant-Microbe Interactions</i> , 1999, 12, 207-217.	1.4	64
51	ISRF1, a transposable insertion sequence from <i>Sinorhizobium fredii</i> . <i>Gene</i> , 1997, 204, 63-69.	1.0	3
52	Construction of multipurpose gene cartridges based on a novel synthetic promoter for high-level gene expression in Gram-negative bacteria. <i>Gene</i> , 1994, 144, 17-24.	1.0	30
53	Experimental conditions may affect reproducibility of the beta-galactosidase assay. <i>FEMS Microbiology Letters</i> , 1992, 100, 87-90.	0.7	41
54	Selection and symbiotic properties of <i>Rhizobium leguminosarum</i> biovar <i>phaseoli</i> strains harboring pRtr5a. <i>Current Microbiology</i> , 1989, 19, 179-181.	1.0	3

#	ARTICLE	IF	CITATIONS
55	Isolation and Characterization of Plant Growth-Promotion Diazotrophic Endophytic Bacteria Associated to Sugarcane ( <i>Saccharum officinarum</i> L.) Grown in ParaÁba, Brazil. Brazilian Archives of Biology and Technology, 0, 65, .	0.5	1