

Andreas Schicho

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4142469/publications.pdf>

Version: 2024-02-01

14
papers

125
citations

1478505

6
h-index

1281871

11
g-index

14
all docs

14
docs citations

14
times ranked

201
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Emphysematous Cystitis: Mortality, Risk Factors, and Pathogens of a Rare Disease. <i>Clinics and Practice</i> , 2017, 7, 930. | 1.4 | 25 |
| 2 | Incidence of traumatic carotid and vertebral artery dissections: results of cervical vessel computed tomography angiogram as a mandatory scan component in severely injured patients. <i>Therapeutics and Clinical Risk Management</i> , 2018, Volume 14, 173-178. | 2.0 | 23 |
| 3 | Long-term survival after percutaneous irreversible electroporation of inoperable colorectal liver metastases. <i>Cancer Management and Research</i> , 2019, Volume 11, 317-322. | 1.9 | 20 |
| 4 | Safety and efficacy of transarterial chemoembolization with degradable starch microspheres (DSM-TACE) in the treatment of secondary liver malignancies. <i>OncoTargets and Therapy</i> , 2018, Volume 11, 345-350. | 2.0 | 13 |
| 5 | Color coded perfusion imaging with contrast enhanced ultrasound (CEUS) for post-interventional success control following trans-arterial chemoembolization (TACE) of hepatocellular carcinoma. <i>PLoS ONE</i> , 2019, 14, e0217599. | 2.5 | 13 |
| 6 | Color Coded Perfusion Imaging with Contrast Enhanced Ultrasound (CEUS) for Post-Interventional Success Control Following Irreversible Electroporation (IRE) of Primary and Secondary Malignant Liver Lesions. <i>Journal of Gastrointestinal and Liver Diseases</i> , 2019, 28, 311-318. | 0.9 | 11 |
| 7 | Modified Split-Scan Computed Tomography (CT) Diagnostics of Severely Injured Patients: First Results from a Level I Trauma Center Using a Dedicated Head-and-Neck CT-Angiogram for the Detection of Cervical Artery Dissections. <i>Journal of Clinical Medicine</i> , 2020, 9, 2568. | 2.4 | 5 |
| 8 | Non-penetrating traumatic injuries of the aortic arch. <i>Acta Radiologica</i> , 2018, 59, 275-279. | 1.1 | 4 |
| 9 | Vascular geometry as a risk factor for non-penetrating traumatic injuries of the aortic arch. <i>PLoS ONE</i> , 2017, 12, e0180066. | 2.5 | 3 |
| 10 | Early Detection of Local Tumor Progression after Irreversible Electroporation (IRE) of Hepatocellular Carcinoma Using Gd-EOB-DTPA-Based MR Imaging at 3T. <i>Cancers</i> , 2021, 13, 1595. | 3.7 | 2 |
| 11 | Necessity of Immediate MRI Imaging in the Acute Care of Severely Injured Patients. <i>Medicina (Lithuania)</i> , 2021, 57, 982. | 2.0 | 2 |
| 12 | Detection of Incomplete Irreversible Electroporation (IRE) and Microwave Ablation (MWA) of Hepatocellular Carcinoma (HCC) Using Iodine Quantification in Dual Energy Computed Tomography (DECT). <i>Diagnostics</i> , 2022, 12, 986. | 2.6 | 2 |
| 13 | <p>Evaluation of Alterations to Bile Ducts and Laboratory Values During the First 3 Months After Irreversible Electroporation of Malignant Hepatic Tumors</p>. <i>Cancer Management and Research</i> , 2020, Volume 12, 8425-8433. | 1.9 | 1 |
| 14 | Occurrence of spontaneous fistulas detected by contrast filling during computed tomography-guided percutaneous drainage placement of splenic and perisplenic fluid collections. <i>Acta Radiologica</i> , 2022, 63, 719-726. | 1.1 | 1 |