Francesca Biandolino

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4145516/publications.pdf

Version: 2024-02-01

40 738 16 25
papers citations h-index g-index

41 41 830 all docs docs citations times ranked citing authors

#	Article	IF	CITATIONS
1	Total lipid content and fatty acid composition of commercially important fish species from the Mediterranean, Mar Grande Sea. Food Chemistry, 2012, 131, 1233-1239.	4.2	92
2	Effect of diet on growth performance, feed efficiency and nutritional composition of Octopus vulgaris. Aquaculture, 2010, 309, 203-211.	1.7	53
3	Influence of a prepared diet and a macroalga (Ulva sp.) on the growth, nutritional and sensory qualities of gonads of the sea urchin Paracentrotus lividus. Aquaculture, 2018, 493, 240-250.	1.7	41
4	Predicting toxicity in marine sediment in Taranto Gulf (Ionian Sea, Southern Italy) using Sediment Quality Guidelines and a battery bioassay. Ecotoxicology, 2007, 16, 239-246.	1,1	33
5	A standardization of Amphibalanus (Balanus) amphitrite (Crustacea, Cirripedia) larval bioassay for ecotoxicological studies. Ecotoxicology and Environmental Safety, 2012, 79, 134-138.	2.9	32
6	Influence of natural diet on growth and biochemical composition of Octopus vulgaris Cuvier, 1797. Aquaculture International, $2010, 18, 1163-1175$.	1.1	28
7	Nutritional Quality of Edible Marine Bivalves from the Southern Coast of Italy, Mediterranean Sea. Polish Journal of Food and Nutrition Sciences, 2019, 69, 71-81.	0.6	27
8	Effect of formulated diets on the proximate composition and fatty acid profiles of sea urchin Paracentrotus lividus gonad. Aquaculture International, 2018, 26, 185-202.	1.1	26
9	Effect of Different Cooking Methods on Lipid Content and Fatty Acid Profiles of Mytilus galloprovincialis. Foods, 2021, 10, 416.	1.9	25
10	Ecotoxicological effects of sediments from Mar Piccolo, South Italy: toxicity testing with organisms from different trophic levels. Environmental Science and Pollution Research, 2016, 23, 12755-12769.	2.7	21
11	Seasonal changes of commercial traits, proximate and fatty acid compositions of the scallop <i>Flexopecten glaber</i> from the Mediterranean Sea (Southern Italy). PeerJ, 2019, 7, e5810.	0.9	21
12	Evaluation of a bioassays battery for ecotoxicological screening of marine sediments from Ionian Sea (Mediterranea Sea, Southern Italy). Environmental Monitoring and Assessment, 2012, 184, 5225-5238.	1.3	20
13	Chronic sublethal effects of ZnO nanoparticles on Tigriopus fulvus (Copepoda, Harpacticoida). Environmental Science and Pollution Research, 2020, 27, 30957-30968.	2.7	19
14	Effects of temperature on the acute toxicity of cadmium to Corophium Insidiosum. Environmental Monitoring and Assessment, 2007, 136, 161-166.	1.3	18
15	Factors influencing the sensitivity of Gammarus aequicauda population from Mar Piccolo in Taranto (Southern Italy). Ecotoxicology and Environmental Safety, 2009, 72, 770-774.	2.9	18
16	Effects of commercial formulations of glyphosate on marine crustaceans and implications for risk assessment under temperature changes. Ecotoxicology and Environmental Safety, 2021, 213, 112068.	2.9	18
17	Seasonal fluctuations of some biological traits of the invader <i>Caprella scaura</i> (Crustacea: Amphipoda: Caprellidae) in the Mar Piccolo of Taranto (Ionian Sea, southern Italy). Scientia Marina, 2013, 77, 169-178.	0.3	18
18	Bioactive fatty acids in seafood from Ionian Sea and relation to dietary recommendations. International Journal of Food Sciences and Nutrition, 2020, 71, 693-705.	1.3	17

#	Article	IF	CITATIONS
19	Preliminary assessment of <i>Ostreopsis </i> cfr. <i>ovata </i> acute toxicity by using a battery bioassay. Chemistry and Ecology, 2011, 27, 117-125.	0.6	15
20	Assessment of individual and combined toxicities of three heavy metals (Cu, Cd and Hg) by using <i>Tigriopus fulvus </i> . Chemistry and Ecology, 2013, 29, 635-642.	0.6	15
21	Gammarus aequicauda(Crustacea: Amphipoda): A potential test species in marine sediment toxicity assessment. Aquatic Ecosystem Health and Management, 2005, 8, 475-482.	0.3	14
22	Ecotoxicological evaluation of sediments by battery bioassays: application and comparison of two integrated classification systems. Chemistry and Ecology, 2015, 31, 661-678.	0.6	14
23	Tigriopus fulvus: The interlaboratory comparison of the acute toxicity test. Ecotoxicology and Environmental Safety, 2016, 124, 309-314.	2.9	14
24	Bioactive fatty acids of three commercial scallop species. International Journal of Food Properties, 2018, 21, 519-532.	1.3	14
25	Life history of the amphipod <i>Corophium insidiosum</i> (Crustacea: Amphipoda) from Mar Piccolo (Ionian Sea, Italy). Scientia Marina, 2006, 70, 355-362.	0.3	14
26	A preliminary investigation of the lipids and fatty acids composition of Gammarus aequicauda (Crustacea: Amphipoda) and its main food source. Journal of the Marine Biological Association of the United Kingdom, 2006, 86, 345-348.	0.4	13
27	Effects of short- and long-term exposures to copper on lethal and reproductive endpoints of the harpacticoid copepod Tigriopus fulvus. Ecotoxicology and Environmental Safety, 2018, 147, 327-333.	2.9	13
28	The recruitment of scallops (and beyond) by two different artificial collectors (Gulf of Taranto,) Tj ETQq0 0 0 rgB	T /Overloc	k 10 Tf 50 382
29	Comparative toxicity of ionic and nanoparticulate zinc in the species Cymodoce truncata, Gammarus aequicauda and Paracentrotus lividus. Environmental Science and Pollution Research, 2021, 28, 42891-42900.	2.7	11
30	Life history of Talorchestia deshayesii (Amphipoda, Talitridae) in the Ionian sandy beach (southern) Tj ETQq0 0 0	rgBT/Ove	rlock 10 Tf 50
31	A toxicity scoring system for the 10-day whole sediment test with Corophium insidiosum (Crawford). Environmental Monitoring and Assessment, 2015, 187, 180.	1.3	9
32	Comparison of amphipodsCorophium insidiosumandC. orientale(Crustacea: Amphipoda) in sediment toxicity testing. Journal of Environmental Science and Health - Part A Toxic/Hazardous Substances and Environmental Engineering, 2010, 45, 1461-1467.	0.9	8
33	Comparative Characteristics of Percentage Edibility, Condition Index, Biochemical Constituents and Lipids Nutritional Quality Indices of Wild and Farmed Scallops (Flexopecten glaber). Water (Switzerland), 2020, 12, 1777.	1.2	8
34	Estimation of Growth Parameters of the Black Scallop Mimachlamys Varia in the Gulf of Taranto (Ionian Sea, Southern Italy). Water (Switzerland), 2020, 12, 3342.	1.2	7
35	Can Different Body Tissues of Two Sea Cucumbers Supply a Fair Amount of Omega 3 for Health Benefit?. Journal of Aquatic Food Product Technology, 2019, 28, 821-836.	0.6	6
36	The Contribution of Fish to the Mediterranean Diet., 2015,, 165-174.		5

#	Article	IF	CITATIONS
37	Hepatopancreas mitochondria of Mytilus galloprovincialis: effect of zinc ions on mitochondrial bioenergetics and metabolism. Turkish Journal of Biology, 2013, 37, 565-572.	2.1	4
38	Strategies for Successful Scallops Spat Collection on Artificial Collectors in the Taranto Gulf (Mediterranean Sea). Water (Switzerland), 2021, 13, 462.	1.2	2
39	Multi-endpoint effects of derelict tubular mussel plastic nets on Tigriopus fulvus. Environmental Science and Pollution Research, 2022, 29, 83554-83566.	2.7	2
40	Bioassays Utilization for Toxicity Assessment of Sediments along Apulia Coast., 2006,,.		0