

Anna Enblom

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4152286/publications.pdf>

Version: 2024-02-01

21
papers

373
citations

933447

10
h-index

794594

19
g-index

21
all docs

21
docs citations

21
times ranked

371
citing authors

#	ARTICLE	IF	CITATIONS
1	Acupuncture compared with placebo acupuncture in radiotherapy-induced nausea—a randomized controlled study. <i>Annals of Oncology</i> , 2012, 23, 1353-1361.	1.2	74
2	One third of patients with radiotherapy-induced nausea consider their antiemetic treatment insufficient. <i>Supportive Care in Cancer</i> , 2009, 17, 23-32.	2.2	71
3	Getting the Grip on Nonspecific Treatment Effects: Emesis in Patients Randomized to Acupuncture or Sham Compared to Patients Receiving Standard Care. <i>PLoS ONE</i> , 2011, 6, e14766.	2.5	39
4	Can individuals identify if needling was performed with an acupuncture needle or a non-penetrating sham needle?. <i>Complementary Therapies in Medicine</i> , 2008, 16, 288-294.	2.7	33
5	Experiences of incontinence and pelvic floor muscle training after gynaecologic cancer treatment. <i>Supportive Care in Cancer</i> , 2017, 25, 157-166.	2.2	22
6	Patients' and Physiotherapists' Belief in and Use of Acupuncture for Cancer-Related Symptoms. <i>Acupuncture in Medicine</i> , 2017, 35, 251-258.	1.0	19
7	Pilot Testing of Methods for Evaluation of Acupuncture for Emesis during Radiotherapy: A Randomised Single Subject Experimental Design. <i>Acupuncture in Medicine</i> , 2011, 29, 94-102.	1.0	17
8	Emesis in patients receiving acupuncture, sham acupuncture or standard care during chemo-radiation: A randomized controlled study. <i>Complementary Therapies in Medicine</i> , 2017, 34, 16-25.	2.7	13
9	The Nonpenetrating Telescopic Sham Needle May Blind Patients with Different Characteristics and Experiences When Treated by Several Therapists. <i>Evidence-based Complementary and Alternative Medicine</i> , 2011, 2011, 1-12.	1.2	12
10	Participation in water-exercising long-term after breast cancer surgery: Experiences of significant factors for continuing exercising as a part of cancer rehabilitation. <i>European Journal of Cancer Care</i> , 2018, 27, e12736.	1.5	11
11	WATER EXERCISE COMPARED TO LAND EXERCISE OR STANDARD CARE IN FEMALE CANCER SURVIVORS WITH SECONDARY LYMPHEDEMA. <i>Lymphology</i> , 2015, 48, 64-79.	0.2	11
12	Emesis and gastrointestinal problems during radiotherapy: A comparison of performance of daily activities between patients experiencing nausea and patients free from nausea. <i>European Journal of Oncology Nursing</i> , 2010, 14, 359-366.	2.1	8
13	Type and Frequency of Side Effects during Pc6 Acupuncture: Observations from Therapists and Patients Participating in Clinical Efficacy Trials of Acupuncture. <i>Acupuncture in Medicine</i> , 2017, 35, 421-429.	1.0	8
14	Reduced Need for Rescue Antiemetics and Improved Capacity to Eat in Patients Receiving Acupuncture Compared to Patients Receiving Sham Acupuncture or Standard Care during Radiotherapy. <i>Evidence-based Complementary and Alternative Medicine</i> , 2017, 2017, 1-11.	1.2	8
15	Health-related quality of life and musculoskeletal function in patients with musculoskeletal disorders: after compared to before short-term group-based aqua-exercises. <i>European Journal of Physiotherapy</i> , 2016, 18, 218-225.	1.3	7
16	Identifying female pelvic cancer survivors with low levels of physical activity after radiotherapy: women with fecal and urinary leakage need additional support. <i>Supportive Care in Cancer</i> , 2020, 28, 2669-2681.	2.2	7
17	Complementary and alternative medicine self-care strategies for nausea in patients undergoing abdominal or pelvic irradiation for cancer: A longitudinal observational study of implementation in routine care. <i>Complementary Therapies in Medicine</i> , 2017, 34, 141-148.	2.7	5
18	Treatment expectations seem to affect bowel health when using acupuncture during radiotherapy for cancer: Secondary outcomes from a clinical randomized sham-controlled trial. <i>Complementary Therapies in Medicine</i> , 2020, 52, 102404.	2.7	3

#	ARTICLE	IF	CITATIONS
19	The Effects of Positive or Neutral Communication during Acupuncture for Relaxing Effects: A Sham-Controlled Randomized Trial. Evidence-based Complementary and Alternative Medicine, 2016, 2016, 1-11.	1.2	2
20	Frequent Stools Were Related to Reduced Quality of Life and Capacity in Daily Activities. Cancer Nursing, 2020, 43, 478-488.	1.5	2
21	Understanding rationales for acupuncture treated individuals' beliefs in acupuncture effects, to be able to maximize therapeutic results: A qualitative analysis. Complementary Therapies in Medicine, 2018, 39, 101-108.	2.7	1