

Taeko Ito

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4157880/publications.pdf>

Version: 2024-02-01

47
papers

398
citations

840776

11
h-index

839539

18
g-index

47
all docs

47
docs citations

47
times ranked

264
citing authors

#	ARTICLE	IF	CITATIONS
1	Endolymphatic space size in patients with Meniere's disease and healthy controls. <i>Acta Oto-Laryngologica</i> , 2016, 136, 879-882.	0.9	39
2	Three-Dimensional Magnetic Resonance Imaging Reveals the Relationship Between the Control of Vertigo and Decreases in Endolymphatic Hydrops After Endolymphatic Sac Drainage With Steroids for Meniere's Disease. <i>Frontiers in Neurology</i> , 2019, 10, 46.	2.4	31
3	A Two-Year Randomized Trial of Interventions to Decrease Stress Hormone Vasopressin Production in Patients with Meniere's Disease—A Pilot Study. <i>PLoS ONE</i> , 2016, 11, e0158309.	2.5	30
4	Volumetric measurements of the inner ear in patients with Meniere's disease using three-dimensional magnetic resonance imaging. <i>Acta Oto-Laryngologica</i> , 2016, 136, 888-893.	0.9	28
5	Endolymphatic volume in patients with meniere's disease and healthy controls: Three-dimensional analysis with magnetic resonance imaging. <i>Laryngoscope Investigative Otolaryngology</i> , 2019, 4, 653-658.	1.5	25
6	Magnetic resonance volumetric measurement of endolymphatic space in patients without vertiginous or cochlear symptoms. <i>Acta Oto-Laryngologica</i> , 2016, 136, 1206-1212.	0.9	21
7	Idiopathic benign paroxysmal positional vertigo with persistent vertigo/dizziness sensation is associated with latent canal paresis, endolymphatic hydrops, and osteoporosis. <i>Auris Nasus Larynx</i> , 2019, 46, 27-33.	1.2	21
8	Surgical results and psychological status in patients with intractable Ménière's disease. <i>Auris Nasus Larynx</i> , 2016, 43, 287-291.	1.2	17
9	Magnetic resonance-based volumetric measurement of the endolymphatic space in patients with Meniere's disease and other endolymphatic hydrops-related diseases. <i>Auris Nasus Larynx</i> , 2019, 46, 493-497.	1.2	16
10	Comparison of the video head impulse test results with caloric test in patients with Meniere's disease and other vestibular disorders. <i>Acta Oto-Laryngologica</i> , 2020, 140, 720-727.	0.9	15
11	Results in caloric test, video head impulse test and inner ear MRI in patients with Ménière's disease. <i>Auris Nasus Larynx</i> , 2020, 47, 71-78.	1.2	12
12	Ménière's disease with unremitting floating sensation is associated with canal paresis, gravity-sensitive dysfunction, mental illness, and bilaterality. <i>Auris Nasus Larynx</i> , 2019, 46, 186-192.	1.2	11
13	Head-Up Sleep May Cure Patients With Intractable Benign Paroxysmal Positional Vertigo: A six-Month Randomized Trial. <i>Laryngoscope Investigative Otolaryngology</i> , 2019, 4, 353-358.	1.5	11
14	Magnetic resonance imaging of the endolymphatic space in patients with acute low-tone sensorineural hearing loss. <i>Auris Nasus Larynx</i> , 2019, 46, 859-865.	1.2	10
15	Effects of Vestibular Rehabilitation on Physical Activity and Subjective Dizziness in Patients With Chronic Peripheral Vestibular Disorders: A Six-Month Randomized Trial. <i>Frontiers in Neurology</i> , 2021, 12, 656157.	2.4	9
16	Relationship between changes in hearing function and volumes of endolymphatic hydrops after endolymphatic sac drainage. <i>Acta Oto-Laryngologica</i> , 2019, 139, 739-746.	0.9	8
17	Magnetic resonance imaging of endolymphatic space in patients with sensorineural hearing loss: comparison between fluctuating and idiopathic sudden sensorineural hearing loss. <i>Acta Oto-Laryngologica</i> , 2020, 140, 345-350.	0.9	8
18	Left parietal involvement in motion sickness susceptibility revealed by multimodal magnetic resonance imaging. <i>Human Brain Mapping</i> , 2022, 43, 1103-1111.	3.6	8

#	ARTICLE	IF	CITATIONS
19	Effects of repeated snowboard exercise in virtual reality with time lags of visual scene behind body rotation on head stability and subjective slalom run performance in healthy young subjects. <i>Acta Oto-Laryngologica</i> , 2016, 136, 1121-1124.	0.9	7
20	Vestibular Compensation after Vestibular Dysfunction Induced by Arsanilic Acid in Mice. <i>Brain Sciences</i> , 2019, 9, 329.	2.3	7
21	Combination of head-up sleep and vertical recognition training may cure intractable motion-evoked dizziness with unknown origin. <i>Acta Oto-Laryngologica</i> , 2020, 140, 467-472.	0.9	7
22	Patients with vertigo/dizziness of unknown origin during follow-ups by general otolaryngologists at outpatient town clinic. <i>Auris Nasus Larynx</i> , 2021, 48, 400-407.	1.2	7
23	Novel Magnetic Resonance Imaging-Based Method for Accurate Diagnosis of Meniere's Disease. <i>Frontiers in Surgery</i> , 2021, 8, 671624.	1.4	7
24	Endolymphatic Sac Drainage Surgery and Plasma Stress Hormone Vasopressin Levels in Meniere's Disease. <i>Frontiers in Neurology</i> , 2021, 12, 722217.	2.4	7
25	Vestibular Morphological Asymmetry Associated With Motion Sickness Susceptibility. <i>Frontiers in Neuroscience</i> , 2021, 15, 763040.	2.8	6
26	A new immunohistochemical method to evaluate the development of vestibular compensation after unilateral labyrinthectomy in rats. <i>Acta Oto-Laryngologica</i> , 2019, 139, 505-510.	0.9	5
27	Effects of posterior tympanotomy with steroids at round window on hearing recovery after revision surgery for intractable Meniere's disease. <i>Acta Oto-Laryngologica</i> , 2015, 135, 667-672.	0.9	4
28	Magnetic resonance imaging of endolymphatic hydrops in patients with unilateral Meniere's disease: volume ratio and distribution rate of the endolymphatic space. <i>Acta Oto-Laryngologica</i> , 2021, 141, 1033-1037.	0.9	4
29	Magnetic Resonance 3D Measurement of the Endolymphatic Space in 100 Control Human Subjects. , 2021, 17, 536-540.		4
30	Changes in the Results of the Subjective Visual Vertical Test After Endolymphatic Sac Drainage for Intractable Meniere's Disease. <i>Journal of International Advanced Otology</i> , 2021, 17, 121-126.	1.0	3
31	Bilateral Asymmetry in Ocular Counter-Rolling Reflex Is Associated With Individual Motion Sickness Susceptibility. <i>Frontiers in Neurology</i> , 2021, 12, 759764.	2.4	3
32	Investigation of endolymphatic hydrops positivity rates in patients with recurrent audiovestibular symptoms using inner ear magnetic resonance imaging. <i>Auris Nasus Larynx</i> , 2022, 49, 188-194.	1.2	2
33	Vestibular rehabilitation at Nara Medical University. <i>MAHOROBATA</i> , Equilibrium Research, 2018, 77, 549-556.	0.1	1
34	Incidence of endolymphatic hydrops among patients with Meniere's disease attending the vertigo clinic of Nara Medical University. <i>Equilibrium Research</i> , 2018, 77, 158-164.	0.1	1
35	What diagnosis should we make for long-lasting vertiginous sensation after acute peripheral vertigo?. <i>Acta Oto-Laryngologica</i> , 2020, 140, 1001-1006.	0.9	1
36	Magnetic resonance imaging of endolymphatic hydrops in patients with unilateral Meniere's disease: a comparison between with and without herniation into the posterior and lateral semi-circular canals. <i>Acta Oto-Laryngologica</i> , 2021, 141, 671-677.	0.9	1

#	ARTICLE	IF	CITATIONS
37	Magnetic resonance imaging of the endolymphatic space in patients with benign paroxysmal positional vertigo: volume ratio and distribution rate of the endolymphatic space. <i>Acta Oto-Laryngologica</i> , 2022, 142, 113-117.	0.9	1
38	Negative prognostic factors for psychological conditions in patients with audiovestibular diseases. <i>Journal of Otolaryngology of Japan</i> , 2017, 120, 884-885.	0.1	0
39	Results in caloric test, vHIT and inner ear MRI in patients with Ménière's disease. <i>Journal of Otolaryngology of Japan</i> , 2021, 124, 930-931.	0.1	0
40	Disease statistics and abnormal clinical exam ratios for patients visiting a vertigo/dizziness center at Nara Medical University. <i>Equilibrium Research</i> , 2018, 77, 136-142.	0.1	0
41	Differences in the handicap associated with dizziness and mental state among patients at the Vertigo/Dizziness Center at Nara Medical University Hospital. <i>Equilibrium Research</i> , 2019, 78, 86-92.	0.1	0
42	Meniere's disease with unremitting floating sensation is associated with canal paresis, gravity sensitive dysfunction, mental illness and bilaterality. <i>Journal of Otolaryngology of Japan</i> , 2020, 123, 405-406.	0.1	0
43	Development and clinical application of Head Tilt SVV to evaluate gravity perception. <i>Equilibrium Research</i> , 2020, 79, 274-280.	0.1	0
44	Magnetic Resonance-Based Volumetric Measurement of the Endolymphatic Space in Patients with Inner Ear Diseases. <i>Practica Otologica</i> , 2020, 113, 667-678.	0.0	0
45	Idiopathic benign paroxysmal positional vertigo with persistent vertigo/dizziness sensation is associated with latent canal paresis, endolymphatic hydrops, and osteoporosis. <i>Journal of Otolaryngology of Japan</i> , 2020, 123, 87-88.	0.1	0
46	Magnetic resonance-based volumetric measurement of the endolymphatic space in patients with Meniere's disease and other endolymphatic hydrops-related diseases. <i>Journal of Otolaryngology of Japan</i> , 2020, 123, 1335-1336.	0.1	0
47	Differences of the findings of MRI between Meniere's disease and vestibular migraine. <i>Equilibrium Research</i> , 2021, 80, 565-571.	0.1	0