

Taeko Ito

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4157880/publications.pdf>

Version: 2024-02-01

47
papers

398
citations

840776

11
h-index

839539

18
g-index

47
all docs

47
docs citations

47
times ranked

264
citing authors

#	ARTICLE	IF	CITATIONS
1	Investigation of endolymphatic hydrops positivity rates in patients with recurrent audiovestibular symptoms using inner ear magnetic resonance imaging. <i>Auris Nasus Larynx</i> , 2022, 49, 188-194.	1.2	2
2	Left parietal involvement in motion sickness susceptibility revealed by multimodal magnetic resonance imaging. <i>Human Brain Mapping</i> , 2022, 43, 1103-1111.	3.6	8
3	Magnetic resonance imaging of the endolymphatic space in patients with benign paroxysmal positional vertigo: volume ratio and distribution rate of the endolymphatic space. <i>Acta Oto-Laryngologica</i> , 2022, 142, 113-117.	0.9	1
4	Patients with vertigo/dizziness of unknown origin during follow-ups by general otolaryngologists at outpatient town clinic. <i>Auris Nasus Larynx</i> , 2021, 48, 400-407.	1.2	7
5	Changes in the Results of the Subjective Visual Vertical Test After Endolymphatic Sac Drainage for Intractable Meniere's Disease. <i>Journal of International Advanced Otolaryngology</i> , 2021, 17, 121-126.	1.0	3
6	Effects of Vestibular Rehabilitation on Physical Activity and Subjective Dizziness in Patients With Chronic Peripheral Vestibular Disorders: A Six-Month Randomized Trial. <i>Frontiers in Neurology</i> , 2021, 12, 656157.	2.4	9
7	Magnetic resonance imaging of endolymphatic hydrops in patients with unilateral Meniere's disease: a comparison between with and without herniation into the posterior and lateral semi-circular canals. <i>Acta Oto-Laryngologica</i> , 2021, 141, 671-677.	0.9	1
8	Results in caloric test, vHIT and inner ear MRI in patients with Ménière's disease. <i>Journal of Otolaryngology of Japan</i> , 2021, 124, 930-931.	0.1	0
9	Novel Magnetic Resonance Imaging-Based Method for Accurate Diagnosis of Meniere's Disease. <i>Frontiers in Surgery</i> , 2021, 8, 671624.	1.4	7
10	Endolymphatic Sac Drainage Surgery and Plasma Stress Hormone Vasopressin Levels in Meniere's Disease. <i>Frontiers in Neurology</i> , 2021, 12, 722217.	2.4	7
11	Vestibular Morphological Asymmetry Associated With Motion Sickness Susceptibility. <i>Frontiers in Neuroscience</i> , 2021, 15, 763040.	2.8	6
12	Bilateral Asymmetry in Ocular Counter-Rolling Reflex Is Associated With Individual Motion Sickness Susceptibility. <i>Frontiers in Neurology</i> , 2021, 12, 759764.	2.4	3
13	Magnetic resonance imaging of endolymphatic hydrops in patients with unilateral Meniere's disease: volume ratio and distribution rate of the endolymphatic space. <i>Acta Oto-Laryngologica</i> , 2021, 141, 1033-1037.	0.9	4
14	Magnetic Resonance 3D Measurement of the Endolymphatic Space in 100 Control Human Subjects. , 2021, 17, 536-540.		4
15	Differences of the findings of MRI between Meniere's disease and vestibular migraine. <i>Equilibrium Research</i> , 2021, 80, 565-571.	0.1	0
16	Results in caloric test, video head impulse test and inner ear MRI in patients with Ménière's disease. <i>Auris Nasus Larynx</i> , 2020, 47, 71-78.	1.2	12
17	What diagnosis should we make for long-lasting vertiginous sensation after acute peripheral vertigo?. <i>Acta Oto-Laryngologica</i> , 2020, 140, 1001-1006.	0.9	1
18	Combination of head-up sleep and vertical recognition training may cure intractable motion-evoked dizziness with unknown origin. <i>Acta Oto-Laryngologica</i> , 2020, 140, 467-472.	0.9	7

#	ARTICLE	IF	CITATIONS
19	Magnetic resonance imaging of endolymphatic space in patients with sensorineural hearing loss: comparison between fluctuating and idiopathic sudden sensorineural hearing loss. <i>Acta Oto-Laryngologica</i> , 2020, 140, 345-350.	0.9	8
20	Comparison of the video head impulse test results with caloric test in patients with Meniere's disease and other vestibular disorders. <i>Acta Oto-Laryngologica</i> , 2020, 140, 720-727.	0.9	15
21	Meniere's disease with unremitting floating sensation is associated with canal paresis, gravity sensitive dysfunction, mental illness and bilaterality. <i>Journal of Otolaryngology of Japan</i> , 2020, 123, 405-406.	0.1	0
22	Development and clinical application of Head Tilt SVV to evaluate gravity perception. <i>Equilibrium Research</i> , 2020, 79, 274-280.	0.1	0
23	Magnetic Resonance-Based Volumetric Measurement of the Endolymphatic Space in Patients with Inner Ear Diseases. <i>Practica Otologica</i> , 2020, 113, 667-678.	0.0	0
24	Idiopathic benign paroxysmal positional vertigo with persistent vertigo/dizziness sensation is associated with latent canal paresis, endolymphatic hydrops, and osteoporosis. <i>Journal of Otolaryngology of Japan</i> , 2020, 123, 87-88.	0.1	0
25	Magnetic resonance-based volumetric measurement of the endolymphatic space in patients with Meniere's disease and other endolymphatic hydrops-related diseases. <i>Journal of Otolaryngology of Japan</i> , 2020, 123, 1335-1336.	0.1	0
26	Meniere's disease with unremitting floating sensation is associated with canal paresis, gravity-sensitive dysfunction, mental illness, and bilaterality. <i>Auris Nasus Larynx</i> , 2019, 46, 186-192.	1.2	11
27	Idiopathic benign paroxysmal positional vertigo with persistent vertigo/dizziness sensation is associated with latent canal paresis, endolymphatic hydrops, and osteoporosis. <i>Auris Nasus Larynx</i> , 2019, 46, 27-33.	1.2	21
28	Relationship between changes in hearing function and volumes of endolymphatic hydrops after endolymphatic sac drainage. <i>Acta Oto-Laryngologica</i> , 2019, 139, 739-746.	0.9	8
29	Head-Up Sleep May Cure Patients With Intractable Benign Paroxysmal Positional Vertigo: A six-Month Randomized Trial. <i>Laryngoscope Investigative Otolaryngology</i> , 2019, 4, 353-358.	1.5	11
30	Magnetic resonance imaging of the endolymphatic space in patients with acute low-tone sensorineural hearing loss. <i>Auris Nasus Larynx</i> , 2019, 46, 859-865.	1.2	10
31	A new immunohistochemical method to evaluate the development of vestibular compensation after unilateral labyrinthectomy in rats. <i>Acta Oto-Laryngologica</i> , 2019, 139, 505-510.	0.9	5
32	Three-Dimensional Magnetic Resonance Imaging Reveals the Relationship Between the Control of Vertigo and Decreases in Endolymphatic Hydrops After Endolymphatic Sac Drainage With Steroids for Meniere's Disease. <i>Frontiers in Neurology</i> , 2019, 10, 46.	2.4	31
33	Endolymphatic volume in patients with meniere's disease and healthy controls: Three-dimensional analysis with magnetic resonance imaging. <i>Laryngoscope Investigative Otolaryngology</i> , 2019, 4, 653-658.	1.5	25
34	Vestibular Compensation after Vestibular Dysfunction Induced by Arsanilic Acid in Mice. <i>Brain Sciences</i> , 2019, 9, 329.	2.3	7
35	Magnetic resonance-based volumetric measurement of the endolymphatic space in patients with Meniere's disease and other endolymphatic hydrops-related diseases. <i>Auris Nasus Larynx</i> , 2019, 46, 493-497.	1.2	16
36	Differences in the handicap associated with dizziness and mental state among patients at the Vertigo/Dizziness Center at Nara Medical University Hospital. <i>Equilibrium Research</i> , 2019, 78, 86-92.	0.1	0

#	ARTICLE	IF	CITATIONS
37	Vestibular rehabilitation at Nara Medical University. <i>MAHOROBAN</i> . Equilibrium Research, 2018, 77, 549-556.	0.1	1
38	Incidence of endolymphatic hydrops among patients with Meniere's disease attending the vertigo clinic of Nara Medical University. <i>Equilibrium Research</i> , 2018, 77, 158-164.	0.1	1
39	Disease statistics and abnormal clinical exam ratios for patients visiting a vertigo/dizziness center at Nara Medical University. <i>Equilibrium Research</i> , 2018, 77, 136-142.	0.1	0
40	Negative prognostic factors for psychological conditions in patients with audiovestibular diseases. <i>Journal of Otolaryngology of Japan</i> , 2017, 120, 884-885.	0.1	0
41	Volumetric measurements of the inner ear in patients with Meniere's disease using three-dimensional magnetic resonance imaging. <i>Acta Oto-Laryngologica</i> , 2016, 136, 888-893.	0.9	28
42	Endolymphatic space size in patients with Meniere's disease and healthy controls. <i>Acta Oto-Laryngologica</i> , 2016, 136, 879-882.	0.9	39
43	Magnetic resonance volumetric measurement of endolymphatic space in patients without vertiginous or cochlear symptoms. <i>Acta Oto-Laryngologica</i> , 2016, 136, 1206-1212.	0.9	21
44	Effects of repeated snowboard exercise in virtual reality with time lags of visual scene behind body rotation on head stability and subjective slalom run performance in healthy young subjects. <i>Acta Oto-Laryngologica</i> , 2016, 136, 1121-1124.	0.9	7
45	Surgical results and psychological status in patients with intractable Ménière's disease. <i>Auris Nasus Larynx</i> , 2016, 43, 287-291.	1.2	17
46	A Two-Year Randomized Trial of Interventions to Decrease Stress Hormone Vasopressin Production in Patients with Meniere's Disease: A Pilot Study. <i>PLoS ONE</i> , 2016, 11, e0158309.	2.5	30
47	Effects of posterior tympanotomy with steroids at round window on hearing recovery after revision surgery for intractable Meniere's disease. <i>Acta Oto-Laryngologica</i> , 2015, 135, 667-672.	0.9	4