

Yayoi Kaneko

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4161780/publications.pdf>

Version: 2024-02-01

57
papers

920
citations

623734

14
h-index

526287

27
g-index

57
all docs

57
docs citations

57
times ranked

1056
citing authors

#	ARTICLE	IF	CITATIONS
1	Biogeographical variation in the diet of Holarctic martens (genus <i>Martes</i> , Mammalia: Carnivora): Tj ETQq1 1 0.784314 rgBT /Overlock 102	3.0	102
2	Molecular Phylogeography of the Brown Bear (<i>Ursus arctos</i>) in Northeastern Asia Based on Analyses of Complete Mitochondrial DNA Sequences. <i>Molecular Biology and Evolution</i> , 2013, 30, 1644-1652.	8.9	98
3	Balancing the benefits of ecotourism and development: The effects of visitor trail-use on mammals in a Protected Area in rapidly developing China. <i>Biological Conservation</i> , 2013, 165, 18-24.	4.1	51
4	Effects of body size on estimation of mammalian area requirements. <i>Conservation Biology</i> , 2020, 34, 1017-1028.	4.7	51
5	Low Genetic Diversity in Japanese Populations of the Eurasian Badger <i>Meles meles</i> (Mustelidae,) Tj ETQq1 1 0.784314 rgBT /Overlock 39 1145-1151.	0.7	39
6	Frugivory and seed dispersal by a small carnivore, the Chinese ferret-badger, <i>Melogale moschata</i> , in a fragmented subtropical forest of central China. <i>Forest Ecology and Management</i> , 2008, 255, 1595-1603.	3.2	36
7	Contrasting Sociality in Two Widespread, Generalist, Mustelid Genera, <i>Meles</i> and <i>Martes</i> . <i>Mammal Study</i> , 2011, 36, 169-188.	0.6	36
8	Diet of an opportunistically frugivorous carnivore, <i>Martes flavigula</i> , in subtropical forest. <i>Journal of Mammalogy</i> , 2011, 92, 611-619.	1.3	32
9	Phylogeographic Sympatry and Isolation of the Eurasian Badgers (<i>Meles</i> , Mustelidae,) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Genes. <i>Zoological Science</i> , 2011, 28, 293-303.	0.7	26
10	The Reliance of the Golden Jackal (<i>Canis aureus</i>) on Anthropogenic Foods in winter in Central Bulgaria. <i>Mammal Study</i> , 2013, 38, 19-27.	0.6	24
11	Variations in Badger (<i>Meles meles</i>) Sett Microclimate: Differential Cub Survival between Main and Subsidiary Setts, with Implications for Artificial Sett Construction. <i>International Journal of Ecology</i> , 2010, 2010, 1-10.	0.8	22
12	Seasonal dietary shifts and food resource exploitation by the hog badger (<i>Arctonyx collaris</i>) in a Chinese subtropical forest. <i>European Journal of Wildlife Research</i> , 2015, 61, 125-133.	1.4	22
13	Home range of raccoon dogs in an urban green area of Tokyo, Japan. <i>Journal of Mammalogy</i> , 2018, 99, 732-740.	1.3	19
14	Spatio-temporal partitioning facilitates mesocarnivore sympatry in the Stara Planina Mountains, Bulgaria. <i>Zoology</i> , 2020, 141, 125801.	1.2	17
15	Diet of Japanese Weasels (<i>Mustela itatsi</i>) in a Sub-Urban Landscape: Implications for Year-Round Persistence of Local Populations. <i>Mammal Study</i> , 2009, 34, 97-105.	0.6	16
16	Spatial and temporal separation between the golden jackal and three sympatric carnivores in a human-modified landscape in central Bulgaria. <i>Zoology and Ecology</i> , 2018, 28, 172-179.	0.2	16
17	Molecular phylogeny of Eurasian badgers (<i>Meles</i>) around the distribution boundaries, revealed by analyses of mitochondrial DNA and Y-chromosomal genes. <i>Biochemical Systematics and Ecology</i> , 2017, 71, 121-130.	1.3	15
18	Genetic population structure of the masked palm civet <i>Paguma larvata</i> , (Carnivora: Viverridae) in Japan, revealed from analysis of newly identified compound microsatellites. <i>Conservation Genetics</i> , 2012, 13, 1095-1107.	1.5	14

#	ARTICLE	IF	CITATIONS
19	Hog badger (<i>Arctonyx collaris</i>) latrine use in relation to food abundance: evidence of the scarce factor paradox. <i>Ecosphere</i> , 2015, 6, 1-12.	2.2	14
20	Origins and Founder Effects on the Japanese Masked Palm Civet (<i>Paguma larvata</i>) (Viverridae). <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 Science</i> , 2010, 27, 499-505.	0.7	13
21	Badger setts provide thermal refugia, buffering changeable surface weather conditions. <i>Journal of Thermal Biology</i> , 2018, 74, 226-233.	2.5	13
22	Comparative Molecular Phylogeny and Evolution of Sex Chromosome DNA Sequences in the Family Canidae (Mammalia: Carnivora). <i>Zoological Science</i> , 2012, 29, 151-161.	0.7	12
23	The socio-spatial dynamics of the Japanese badger (<i>Meles anakuma</i>). <i>Journal of Mammalogy</i> , 2014, 95, 290-300.	1.3	12
24	Masked Palm Civet (<i>Paguma larvata</i>) Summer Diet Differs between Sexes in a Suburban Area of Central Japan. <i>Mammal Study</i> , 2017, 42, 185-190.	0.6	11
25	Diversity of MHC class II DRB alleles in the Eurasian population of the least weasel, <i>Mustela nivalis</i> (Mustelidae: Mammalia). <i>Biological Journal of the Linnean Society</i> , 2017, 121, 28-37.	1.6	11
26	Diversity and evolution of MHC class II DRB gene in the Eurasian badger genus <i>Meles</i> (Mammalia: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50	1.6	11
27	Thermal forest zone explains regional variations in the diet composition of the Japanese marten (<i>Martes melampus</i>). <i>Mammalian Biology</i> , 2019, 95, 173-180.	1.5	11
28	Ranging and activity patterns of the group-living ferret badger <i>Melogale moschata</i> in Central China. <i>Journal of Mammalogy</i> , 2010, 91, 101-108.	1.3	10
29	Identification and Molecular Variations of CAN-SINEs from the <i>ZFY</i> Gene Final Intron of the Eurasian Badgers (Genus <i>Meles</i>). <i>Mammal Study</i> , 2011, 36, 41-48.	0.6	10
30	Preliminary Study on Microsatellite and Mitochondrial DNA Variation of the Stone Marten (<i>Martes foina</i>) in Bulgaria. <i>Mammal Study</i> , 2012, 37, 353-358.	0.6	10
31	Human disturbance affects latrine use patterns of raccoon dogs. <i>Journal of Wildlife Management</i> , 2019, 83, 728-736.	1.8	10
32	Latrine use in a Low Density Japanese Badger (<i>Meles anakuma</i>) Population Determined by a Continuous Tracking System. <i>Mammal Study</i> , 2009, 34, 179-186.	0.6	9
33	A Comparison of Visual and Genetic Techniques for Identifying Japanese Marten Scats - Enabling Diet Examination in Relation to Seasonal Food Availability in a Sub-Alpine Area of Japan. <i>Zoological Science</i> , 2017, 34, 137-146.	0.7	9
34	Unjustified killing of badgers in Kyushu. <i>Nature</i> , 2017, 544, 161-161.	27.8	9
35	Evolution of MHC class I genes in Eurasian badgers, genus <i>Meles</i> (Carnivora, Mustelidae). <i>Heredity</i> , 2019, 122, 205-218.	2.6	9
36	Variation in the diets of Japanese martens (<i>Martes melampus</i>). <i>Mammal Review</i> , 2019, 49, 121-128.	4.8	9

#	ARTICLE	IF	CITATIONS
37	Winter Diet of the Raccoon Dog (<i>Nyctereutes procyonoides</i>) in Urban Parks, Central Tokyo. <i>Mammal Study</i> , 2018, 43, .	0.6	8
38	The "Trace Recorder", a new device for surveying mammal home ranges, and its application to raccoon dog research. <i>Mammal Study</i> , 1998, 23, 109-118.	0.6	7
39	The illegal exploitation of hog badgers (<i>Arctonyx collaris</i>) in China: genetic evidence exposes regional population impacts. <i>Conservation Genetics Resources</i> , 2015, 7, 697-704.	0.8	7
40	Population Structure of the Raccoon Dog on the Grounds of the Imperial Palace, Tokyo, Revealed by Microsatellite Analysis of Fecal DNA. <i>Zoological Science</i> , 2016, 33, 485-490.	0.7	7
41	Hybridization between the European and Asian badgers (<i>Meles</i> , Carnivora) in the Volga-Kama region, revealed by analyses of maternally, paternally and biparentally inherited genes. <i>Mammalian Biology</i> , 2019, 94, 140-148.	1.5	7
42	Genetic variations of the masked palm civet <i>Paguma larvata</i> , inferred from mitochondrial cytochrome <i>b</i> sequences. <i>Mammal Study</i> , 2008, 33, 19-24.	0.6	6
43	Genetic Diversity within the Japanese Badgers (<i>Meles anakuma</i>), as Revealed by Microsatellite Analysis. <i>Mammal Study</i> , 2010, 35, 221-226.	0.6	6
44	Molecular phylogenetic status of the Bulgarian marbled polecat (<i>Vormela peregusna</i> , Mustelidae,) <i>Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 Systematics and Ecology</i> , 2017, 70, 99-107.	1.3	6
45	Winter diet of the stone marten (<i>Martes foina</i>) in central Bulgaria. <i>Mammal Study</i> , 2013, 38, 293-298.	0.6	4
46	Broad-Leaved Forest Selection of the Japanese Marten (<i>Martes melampus</i>) in Central Japan Revealed by Camera Trapping. <i>Mammal Study</i> , 2014, 39, 163-166.	0.6	4
47	Comparative phylogeography of the endemic Japanese weasel (<i>Mustela itatsi</i>) and the continental Siberian weasel (<i>Mustela sibirica</i>) revealed by complete mitochondrial genome sequences. <i>Biological Journal of the Linnean Society</i> , 2016, , .	1.6	4
48	Phylogeography and population history of the least weasel (<i>Mustela nivalis</i>) in the Palearctic based on multilocus analysis. <i>Journal of Zoological Systematics and Evolutionary Research</i> , 2020, 58, 408-426.	1.4	4
49	Autumn Dietary Overlaps among Three Sympatric Mesocarnivores in the Central Part of Stara Planina Mountain, Bulgaria. <i>Mammal Study</i> , 2019, 44, 275.	0.6	4
50	Food Habits of the Urban Japanese Weasels <i>Mustela itatsi</i> Revealed by Faecal DNA Analysis. <i>Mammal Study</i> , 2014, 39, 155-161.	0.6	3
51	Mitochondrial haplogrouping of the ancient brown bears (<i>Ursus arctos</i>) in Bulgaria, revealed by the APLP method. <i>Mammal Research</i> , 2020, 65, 413-421.	1.3	3
52	Population genetic structure and diversity of the East Balkan Swine (<i>Sus scrofa</i>) in Bulgaria, revealed by mitochondrial DNA and microsatellite analyses. <i>Animal Science Journal</i> , 2021, 92, e13630.	1.4	3
53	Notes on Stomach Contents of Japanese Weasels (<i>Mustela itatsi</i>) in Ibaraki, Japan. <i>Mammal Study</i> , 2013, 38, 281-285.	0.6	2
54	Genetic diversity of MHC class II <i>DRB</i> alleles in the marbled polecat, <i>Vormela peregusna</i> , in Bulgaria. <i>Ethology Ecology and Evolution</i> , 2019, 31, 59-72.	1.4	2

#	ARTICLE	IF	CITATIONS
55	Existence Detection by On-off Code of Magnetic Dipole Field and Its Use in Recording Small Wild Animal's Behavior. Transactions of the Society of Instrument and Control Engineers, 1999, 35, 167-175.	0.2	2
56	Introduction and Expansion History of the Masked Palm Civet, Paguma larvata, in Japan, Revealed by Mitochondrial DNA Control Region and Cytochrome b Analysis. Mammal Study, 2020, 45, 243.	0.6	2
57	River bank role as wildlife habitats : implication for combining hydrospheric disaster prevention and ecosystem conservation at Tama River basin. Journal of the Japanese Society of Revegetation Technology, 2019, 44, 507-510.	0.1	0