
Mohammad Shahnaij

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4164305/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Geophagy is Associated with Environmental Enteropathy and Stunting in Children in Rural Bangladesh. American Journal of Tropical Medicine and Hygiene, 2015, 92, 1117-1124.	1.4	124
2	Changing Trends in the Prevalence of Shigella Species: Emergence of Multi-Drug Resistant Shigella sonnei Biotype g in Bangladesh. PLoS ONE, 2013, 8, e82601.	2.5	79
3	Enteric Infections in Young Children are Associated with Environmental Enteropathy and Impaired Growth. Tropical Medicine and International Health, 2018, 23, 26-33.	2.3	72
4	Unsafe Child Feces Disposal is Associated with Environmental Enteropathy and Impaired Growth. Journal of Pediatrics, 2016, 176, 43-49.	1.8	50
5	Fluoroquinolone Resistance Mechanisms of Shigella flexneri Isolated in Bangladesh. PLoS ONE, 2014, 9, e102533.	2.5	46
6	Resistance Pattern and Molecular Characterization of Enterotoxigenic Escherichia coli (ETEC) Strains Isolated in Bangladesh. PLoS ONE, 2016, 11, e0157415.	2.5	46
7	Mouthing of Soil Contaminated Objects is Associated with Environmental Enteropathy in Young Children. Tropical Medicine and International Health, 2017, 22, 670-678.	2.3	36
8	Vaccinomics Approach for Designing Potential Peptide Vaccine by Targeting <i>Shigella</i> spp. Serine Protease Autotransporter Subfamily Protein SigA. Journal of Immunology Research, 2017, 2017, 1-14.	2.2	27
9	<i>Shigella</i> Infections in Household Contacts of Pediatric Shigellosis Patients in Rural Bangladesh. Emerging Infectious Diseases, 2015, 21, 2006-2013.	4.3	24
10	PCR-Based Method for Shigella flexneri Serotyping: International Multicenter Validation. Journal of Clinical Microbiology, 2019, 57, .	3.9	21
11	FAS promoter polymorphisms and serum sFas level are associated with increased risk of nerve damage in Bangladeshi patients with Guillain-Barré syndrome. PLoS ONE, 2018, 13, e0192703.	2.5	14
12	Characterization of a serologically atypical Shigella flexneri Z isolated from diarrheal patients in Bangladesh and a proposed serological scheme for Shigella flexneri. PLoS ONE, 2018, 13, e0202704.	2.5	12
13	Liver-Directed AAV8 Booster Vaccine Expressing Plasmodium falciparum Antigen Following Adenovirus Vaccine Priming Elicits Sterile Protection in a Murine Model. Frontiers in Immunology, 2021, 12, 612910.	4.8	8