

Peter Å Æ³Å;

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4164496/publications.pdf>

Version: 2024-02-01

16
papers

1,030
citations

933447

10
h-index

1199594

12
g-index

18
all docs

18
docs citations

18
times ranked

1395
citing authors

#	ARTICLE	IF	CITATIONS
1	The Effect of a Single Dose of Intravenous Ketamine on Suicidal Ideation: A Systematic Review and Individual Participant Data Meta-Analysis. <i>American Journal of Psychiatry</i> , 2018, 175, 150-158.	7.2	476
2	Early reduction in prefrontal theta QEEG cordance value predicts response to venlafaxine treatment in patients with resistant depressive disorder. <i>European Psychiatry</i> , 2008, 23, 350-355.	0.2	120
3	Relationship of ketamine's antidepressant and psychotomimetic effects in unipolar depression. <i>Neuroendocrinology Letters</i> , 2013, 34, 287-93.	0.2	105
4	The change of prefrontal QEEG theta cordance as a predictor of response to bupropion treatment in patients who had failed to respond to previous antidepressant treatments. <i>European Neuropsychopharmacology</i> , 2010, 20, 459-466.	0.7	81
5	Electroencephalographic Spectral and Coherence Analysis of Ketamine in Rats: Correlation with Behavioral Effects and Pharmacokinetics. <i>Neuropsychobiology</i> , 2011, 63, 202-218.	1.9	73
6	Low frequency (1-Hz), right prefrontal repetitive transcranial magnetic stimulation (rTMS) compared with venlafaxine ER in the treatment of resistant depression: A double-blind, single-centre, randomized study. <i>Journal of Affective Disorders</i> , 2009, 118, 94-100.	4.1	53
7	QEEG Theta Cordance in the Prediction of Treatment Outcome to Prefrontal Repetitive Transcranial Magnetic Stimulation or Venlafaxine ER in Patients With Major Depressive Disorder. <i>Clinical EEG and Neuroscience</i> , 2015, 46, 73-80.	1.7	39
8	Impact of midazolam vs. saline on effect size estimates in controlled trials of ketamine as a rapid-acting antidepressant. <i>Neuropsychopharmacology</i> , 2019, 44, 1233-1238.	5.4	35
9	Discontinuation of hypnotics during cognitive behavioural therapy for insomnia. <i>BMC Psychiatry</i> , 2008, 8, 80.	2.6	22
10	Trazodone improves the results of cognitive behaviour therapy of primary insomnia in non-depressed patients. <i>Neuroendocrinology Letters</i> , 2008, 29, 895-901.	0.2	15
11	Is combined treatment more effective than switching to monotherapy in patients with resistant depression? A retrospective study. <i>Neuroendocrinology Letters</i> , 2009, 30, 723-8.	0.2	6
12	Structural and construct validity of the Czech version of the Pittsburgh Sleep Quality Index in chronic insomnia. <i>Neuroendocrinology Letters</i> , 2017, 38, 67-73.	0.2	5
13	The decrease of prefrontal theta QEEG cordance value predicts response to Venlafaxine treatment in patients with resistant depression. <i>European Psychiatry</i> , 2008, 23, S74.	0.2	0
14	Differences in brain electrical activity after the cerebellar hemisphere and the vermis rTMS. <i>European Psychiatry</i> , 2008, 23, S287-S288.	0.2	0
15	P.3.020 How ketamine changes the neurophysiology of depressive patients' brains: a randomised controlled trial. <i>European Neuropsychopharmacology</i> , 2011, 21, S75.	0.7	0
16	Is there a future for other glutamate receptor modulators in the pharmacotherapy of mood disorders?. <i>Evidence-Based Mental Health</i> , 2016, 19, 95-96.	4.5	0