

# Atsushi Ohwaki

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4165167/publications.pdf>

Version: 2024-02-01

15  
papers

146  
citations

1478505

6  
h-index

1281871

11  
g-index

15  
all docs

15  
docs citations

15  
times ranked

183  
citing authors

#	ARTICLE	IF	CITATIONS
1	Identification of source populations for reintroduction in extinct populations based on genome-wide SNPs and mtDNA sequence: a case study of the endangered subalpine grassland butterfly <i>Aporia hippia</i> (Lepidoptera; Pieridae) in Japan. <i>Journal of Insect Conservation</i> , 2022, 26, 121-130.	1.4	3
2	Different community assembly of ground beetles and spiders in subalpine forests and alpine scoria deserts of a young volcano, Mt. Fuji. <i>Ecological Research</i> , 2021, 36, 866-881.	1.5	0
3	Entire-area spring burning versus abandonment in grasslands: butterfly responses associated with hibernating traits. <i>Journal of Insect Conservation</i> , 2019, 23, 857-871.	1.4	3
4	Prevalence of Falls on Mount Fuji and Associated with Risk Factors: A Questionnaire Survey Study. <i>International Journal of Environmental Research and Public Health</i> , 2019, 16, 4234.	2.6	6
5	The role of linear mown firebreaks in conserving butterfly diversity: effects of adjacent vegetation and management. <i>Entomological Science</i> , 2018, 21, 112-123.	0.6	8
6	How should we view temperate semi-natural grasslands? Insights from butterflies in Japan. <i>Global Ecology and Conservation</i> , 2018, 16, e00482.	2.1	13
7	Evaluating forest clear-cuts as alternative grassland habitats for plants and butterflies. <i>Forest Ecology and Management</i> , 2018, 430, 337-345.	3.2	16
8	Seasonal variability in the response of ground beetles (Coleoptera: Carabidae) to a forest edge in a heterogeneous agricultural landscape in Japan. <i>European Journal of Entomology</i> , 2015, 112, 135-144.	1.2	21
9	Ground arthropod communities in paddy fields during the dry period: Comparison between different farming methods. <i>Journal of Asia-Pacific Entomology</i> , 2015, 18, 413-419.	0.9	3
10	Butterfly responses to cultivated field abandonment are related with ecological traits in a temperate Japanese agricultural landscape. <i>Landscape and Urban Planning</i> , 2014, 125, 174-182.	7.5	6
11	Differences in tree community among secondary deciduous oak forests in rural and residential areas in the Hokuriku District of Japan. <i>Landscape and Ecological Engineering</i> , 2013, 9, 99-110.	1.5	3
12	Effects of creation of open vegetation in abandoned terraced paddy fields on carabid beetle assemblages in temperate Japan. <i>Entomological Science</i> , 2013, 16, 379-389.	0.6	2
13	Effects of anthropogenic disturbances on the butterfly assemblage in an urban green area: the changes from 1990 to 2005 in Kanazawa Castle Park, Japan. <i>Ecological Research</i> , 2008, 23, 697-708.	1.5	16
14	Butterfly assemblages in a traditional agricultural landscape: importance of secondary forests for conserving diversity, life history specialists and endemics. <i>Biodiversity and Conservation</i> , 2007, 16, 1521-1539.	2.6	31
15	Associations between canopy openness, butterfly resources, butterfly richness and abundance along forest trails in planted and natural forests. <i>European Journal of Entomology</i> , 0, 114, 533-545.	1.2	15