

Andr Luiz Cazetta

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/4170692/andre-luiz-cazetta-publications-by-year.pdf>

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

44
papers

3,617
citations

25
h-index

44
g-index

44
ext. papers

4,296
ext. citations

8.8
avg, IF

5.48
L-index

#	Paper	IF	Citations
44	Sulfonated carbon: synthesis, properties and production of biodiesel. <i>Chemical Engineering and Processing: Process Intensification</i> , 2022 , 170, 108668	3.7	1
43	N-doped spherical activated carbon from dye adsorption: Bifunctional electrocatalyst for hydrazine oxidation and oxygen reduction. <i>Journal of Environmental Chemical Engineering</i> , 2022 , 10, 107458	6.8	1
42	Steam-activated carbon from malt bagasse: Optimization of preparation conditions and adsorption studies of sunset yellow food dye. <i>Arabian Journal of Chemistry</i> , 2021 , 14, 103001	5.9	14
41	The use of chemometric tools for screening and optimization of variables in the preparation and application of carbon-based materials. <i>Journal of the Taiwan Institute of Chemical Engineers</i> , 2021 , 121, 321-336	5.3	2
40	H3PO4-activated carbon fibers of high surface area from banana tree pseudo-stem fibers: Adsorption studies of methylene blue dye in batch and fixed bed systems. <i>Journal of Molecular Liquids</i> , 2021 , 324, 114771	6	20
39	Optimization of thermal conditions of sol-gel method for synthesis of TiO ₂ using RSM and its influence on photodegradation of tartrazine yellow dye. <i>Journal of Environmental Chemical Engineering</i> , 2021 , 9, 104753	6.8	6
38	Biochar from the mixture of poultry litter and charcoal fines as soil conditioner: Optimization of preparation conditions via response surface methodology. <i>Bioresource Technology Reports</i> , 2021 , 15, 100800	4.1	0
37	Caffeine adsorption on activated biochar derived from macrophytes (<i>Eichornia crassipes</i>). <i>Journal of Molecular Liquids</i> , 2021 , 340, 117206	6	2
36	Optimization of sulfonation process for the development of carbon-based catalyst from crambe meal via response surface methodology. <i>Energy Conversion and Management</i> , 2020 , 217, 112975	10.6	15
35	Sugarcane vinasse-derived nanoporous N-S-doped carbon material decorated with Co: A new and efficient multifunctional electrocatalyst. <i>International Journal of Hydrogen Energy</i> , 2020 , 45, 9669-9682	6.7	14
34	Synthesis of superparamagnetic activated carbon for paracetamol removal from aqueous solution. <i>Journal of Molecular Liquids</i> , 2020 , 300, 112282	6	16
33	Activated carbon fibers prepared from cellulose and polyester-derived residues and their application on removal of Pb ²⁺ ions from aqueous solution. <i>Journal of Molecular Liquids</i> , 2019 , 289, 111150	6	13
32	Stevia residue as new precursor of CO-activated carbon: Optimization of preparation condition and adsorption study of triclosan. <i>Ecotoxicology and Environmental Safety</i> , 2019 , 172, 403-410	7	27
31	KOH-super activated carbon from biomass waste: Insights into the paracetamol adsorption mechanism and thermal regeneration cycles. <i>Journal of Hazardous Materials</i> , 2019 , 371, 499-505	12.8	97
30	Nanoporous Heteroatom-Doped Carbons Derived From Cotton Waste: Efficient Hydrazine Oxidation Electrocatalysts. <i>ACS Applied Energy Materials</i> , 2019 , 2, 2313-2323	6.1	20
29	Metal-free ovalbumin-derived N-S-co-doped nanoporous carbon materials as efficient electrocatalysts for oxygen reduction reaction. <i>Applied Surface Science</i> , 2019 , 467-468, 75-83	6.7	22
28	Chemometric study of thermal treatment effect on the P25 photoactivity for degradation of tartrazine yellow dye. <i>Ceramics International</i> , 2018 , 44, 12292-12300	5.1	8

27	New magnetic chitosan/alginate/Fe ₃ O ₄ @SiO ₂ hydrogel composites applied for removal of Pb(II) ions from aqueous systems. <i>Chemical Engineering Journal</i> , 2018 , 337, 595-608	14.7	76
26	Porosity enhancement of spherical activated carbon: Influence and optimization of hydrothermal synthesis conditions using response surface methodology. <i>Journal of Environmental Chemical Engineering</i> , 2018 , 6, 991-999	6.8	29
25	Mesoporous activated carbon fibers synthesized from denim fabric waste: Efficient adsorbents for removal of textile dye from aqueous solutions. <i>Journal of Cleaner Production</i> , 2018 , 171, 482-490	10.3	96
24	Adsorption of caffeine on mesoporous activated carbon fibers prepared from pineapple plant leaves. <i>Ecotoxicology and Environmental Safety</i> , 2018 , 147, 64-71	7	158
23	CO ₂ -spherical activated carbon as a new adsorbent for Methylene Blue removal: Kinetic, equilibrium and thermodynamic studies. <i>Journal of Molecular Liquids</i> , 2018 , 269, 132-139	6	51
22	Bone char-derived metal-free N- and S-co-doped nanoporous carbon and its efficient electrocatalytic activity for hydrazine oxidation. <i>Applied Catalysis B: Environmental</i> , 2018 , 225, 30-39	21.8	78
21	Inexpensive Bismuth-Film Electrode Supported on Pencil-Lead Graphite for Determination of Pb(II) and Cd(II) Ions by Anodic Stripping Voltammetry. <i>International Journal of Analytical Chemistry</i> , 2018 , 2018, 1473706	1.4	15
20	Bone char prepared by CO ₂ atmosphere: Preparation optimization and adsorption studies of Remazol Brilliant Blue R. <i>Journal of Cleaner Production</i> , 2017 , 161, 288-298	10.3	28
19	Sol-gel synthesis of new TiO ₂ /activated carbon photocatalyst and its application for degradation of tetracycline. <i>Ceramics International</i> , 2017 , 43, 4411-4418	5.1	94
18	Hydrothermal carbonization of coffee husk: Optimization of experimental parameters and adsorption of methylene blue dye. <i>Journal of Environmental Chemical Engineering</i> , 2017 , 5, 4841-4849	6.8	54
17	Preparation of biosorbents from the Jatoba (<i>Hymenaea courbaril</i>) fruit shell for removal of Pb(II) and Cd(II) from aqueous solution. <i>Environmental Monitoring and Assessment</i> , 2017 , 189, 632	3.1	11
16	Percolation as new method of preparation of modified biosorbents for pollutants removal. <i>Chemical Engineering Journal</i> , 2016 , 283, 1305-1314	14.7	24
15	Mesoporous activated carbon from industrial laundry sewage sludge: Adsorption studies of reactive dye Remazol Brilliant Blue R. <i>Chemical Engineering Journal</i> , 2016 , 303, 467-476	14.7	165
14	Magnetic Activated Carbon Derived from Biomass Waste by Concurrent Synthesis: Efficient Adsorbent for Toxic Dyes. <i>ACS Sustainable Chemistry and Engineering</i> , 2016 , 4, 1058-1068	8.3	183
13	NaOH-activated carbon of high surface area produced from guava seeds as a high-efficiency adsorbent for amoxicillin removal: Kinetic, isotherm and thermodynamic studies. <i>Chemical Engineering Journal</i> , 2016 , 288, 778-788	14.7	240
12	KOH-activated carbon prepared from sucrose spherical carbon: Adsorption equilibrium, kinetic and thermodynamic studies for Methylene Blue removal. <i>Chemical Engineering Journal</i> , 2016 , 286, 476-484	14.7	329
11	Synthesis and application of N ₂ -doped mesoporous carbon obtained from nanocasting method using bone char as heteroatom precursor and template. <i>Chemical Engineering Journal</i> , 2016 , 300, 54-63	14.7	50
10	Removal of tetracycline by NaOH-activated carbon produced from macadamia nut shells: Kinetic and equilibrium studies. <i>Chemical Engineering Journal</i> , 2015 , 260, 291-299	14.7	407

9	Synthesis of ZnCl ₂ -activated carbon from macadamia nut endocarp (<i>Macadamia integrifolia</i>) by microwave-assisted pyrolysis: Optimization using RSM and methylene blue adsorption. <i>Journal of Analytical and Applied Pyrolysis</i> , 2014 , 105, 166-176	6	97
8	Thermally activated carbon from bovine bone: Optimization of synthesis conditions by response surface methodology. <i>Journal of Analytical and Applied Pyrolysis</i> , 2014 , 110, 455-462	6	12
7	N-doped ordered mesoporous carbons with improved charge storage capacity by tailoring N-dopant density with solvent-assisted synthesis. <i>Journal of Materials Chemistry A</i> , 2014 , 2, 15181-15190 ¹³		45
6	Adsorption studies of methylene blue onto ZnCl ₂ -activated carbon produced from buriti shells (<i>Mauritia flexuosa</i> L.). <i>Journal of Industrial and Engineering Chemistry</i> , 2014 , 20, 4401-4407	6.3	153
5	Thermal regeneration study of high surface area activated carbon obtained from coconut shell: Characterization and application of response surface methodology. <i>Journal of Analytical and Applied Pyrolysis</i> , 2013 , 101, 53-60	6	68
4	Kinetic and equilibrium studies: Adsorption of food dyes Acid Yellow 6, Acid Yellow 23, and Acid Red 18 on activated carbon from flamboyant pods. <i>Chemical Engineering Journal</i> , 2012 , 181-182, 243-250 ^{14.7}		96
3	NaOH-activated carbon of high surface area produced from coconut shell: Kinetics and equilibrium studies from the methylene blue adsorption. <i>Chemical Engineering Journal</i> , 2011 , 174, 117-125	14.7	365
2	Adsorption of methylene blue on activated carbon produced from flamboyant pods (<i>Delonix regia</i>): Study of adsorption isotherms and kinetic models. <i>Chemical Engineering Journal</i> , 2011 , 168, 722-730	14.7	320
1	Preparation and characterization of activated carbon from a new raw lignocellulosic material: flamboyant (<i>Delonix regia</i>) pods. <i>Journal of Environmental Management</i> , 2011 , 92, 178-84	7.9	95