

Kei Sato

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4175051/publications.pdf>

Version: 2024-02-01

88
papers

2,779
citations

196777
29
h-index

242451
47
g-index

99
all docs

99
docs citations

99
times ranked

3175
citing authors

#	ARTICLE	IF	CITATIONS
1	Secondary organic aerosol formation from gasoline and diesel vehicle exhaust under light and dark conditions. <i>Environmental Science Atmospheres</i> , 2022, 2, 46-64.	0.9	5
2	Investigation of OH-reactivity budget in the isoprene, α -pinene and m-xylene oxidation with OH under high NO _x conditions. <i>Atmospheric Environment</i> , 2022, 271, 118916.	1.9	6
3	Formation of secondary organic aerosol tracers from anthropogenic and biogenic volatile organic compounds under varied NO and oxidant conditions. <i>Atmospheric Environment: X</i> , 2022, , 100169.	0.8	0
4	Impacts of missing OH reactivity and aerosol uptake of HO ₂ radicals on tropospheric O ₃ production during the AQUAS-Kyoto summer campaign in 2018. <i>Atmospheric Environment</i> , 2022, 281, 119130.	1.9	1
5	Temperature and acidity dependence of secondary organic aerosol formation from α -pinene ozonolysis with a compact chamber system. <i>Atmospheric Chemistry and Physics</i> , 2021, 21, 5983-6003.	1.9	17
6	Source contributions to multiple toxic potentials of atmospheric organic aerosols. <i>Science of the Total Environment</i> , 2021, 773, 145614.	3.9	30
7	Kinetics and impacting factors of HO ₂ uptake onto submicron atmospheric aerosols during the 2019 Air QUALity Study (AQUAS) in Yokohama, Japan. <i>Atmospheric Chemistry and Physics</i> , 2021, 21, 12243-12260.	1.9	16
8	Effects of Metal Ions on Aqueous-Phase Decomposition of α -Hydroxyalkyl-Hydroperoxides Derived from Terpene Alcohols. <i>Environmental Science & Technology</i> , 2021, 55, 12893-12901.	4.6	5
9	A quantitative understanding of total OH reactivity and ozone production in a coastal industrial area during the Yokohama air quality study (AQUAS) campaign of summer 2019. <i>Atmospheric Environment</i> , 2021, 267, 118754.	1.9	2
10	Nitrate radical, ozone and hydroxyl radical initiated aging of limonene secondary organic aerosol. <i>Atmospheric Environment: X</i> , 2021, 9, 100102.	0.8	0
11	Structural Characterisation of Dimeric Esters in α -Pinene Secondary Organic Aerosol Using N ₂ and CO ₂ Ion Mobility Mass Spectrometry. <i>Atmosphere</i> , 2021, 12, 17.	1.0	5
12	Four- and Five-Carbon Dicarboxylic Acids Present in Secondary Organic Aerosol Produced from Anthropogenic and Biogenic Volatile Organic Compounds. <i>Atmosphere</i> , 2021, 12, 1703.	1.0	9
13	Aerosol Liquid Water Promotes the Formation of Water-Soluble Organic Nitrogen in Submicrometer Aerosols in a Suburban Forest. <i>Environmental Science & Technology</i> , 2020, 54, 1406-1414.	4.6	33
14	Mid carbon (C ₆ +C ₂₉) in refractory black carbon aerosols is a potential tracer of open burning of rice straw: Insights from atmospheric observation and emission source studies. <i>Atmospheric Environment</i> , 2020, 238, 117729.	1.9	11
15	Modeling the Effects of Dimerization and Bulk Diffusion on the Evaporative Behavior of Secondary Organic Aerosol Formed from α -Pinene and 1,3,5-Trimethylbenzene. <i>ACS Earth and Space Chemistry</i> , 2020, 4, 1931-1946.	1.2	7
16	Comparative Analysis of PM _{2.5} -Bound Polycyclic Aromatic Hydrocarbons (PAHs), Nitro-PAHs (NPAHs), and Water-Soluble Inorganic Ions (WSIIs) at Two Background Sites in Japan. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 8224.	1.2	17
17	Volatility Distribution of Organic Compounds in Sewage Incineration Emissions. <i>Environmental Science & Technology</i> , 2020, 54, 14235-14245.	4.6	10
18	Total hydroxyl radical reactivity measurements in a suburban area during AQUAS's "Tsukuba campaign in summer 2017. <i>Science of the Total Environment</i> , 2020, 740, 139897.	3.9	9

#	ARTICLE	IF	CITATIONS
19	Degradation of PAHs during long range transport based on simultaneous measurements at Tuoji Island, China, and at Fukue Island and Cape Hedo, Japan. <i>Environmental Pollution</i> , 2020, 260, 113906.	3.7	23
20	Effects of pH on Interfacial Ozonolysis of α -Terpineol. <i>Journal of Physical Chemistry A</i> , 2019, 123, 7148-7155.	1.1	21
21	Relative and Absolute Sensitivity Analysis on Ozone Production in Tsukuba, a City in Japan. <i>Environmental Science & Technology</i> , 2019, 53, 13629-13635.	4.6	17
22	Investigation of dark condition nitrate radical- and ozone-initiated aging of toluene secondary organic aerosol: Importance of nitrate radical reactions with phenolic products. <i>Atmospheric Environment</i> , 2019, 219, 117049.	1.9	14
23	Seasonal and annual changes in PAH concentrations in a remote site in the Pacific Ocean. <i>Scientific Reports</i> , 2019, 9, 12591.	1.6	30
24	A study of volatility by composition, heating, and dilution measurements of secondary organic aerosol from 1,3,5-trimethylbenzene. <i>Atmospheric Chemistry and Physics</i> , 2019, 19, 14901-14915.	1.9	16
25	Effect of Oxidation Process on Complex Refractive Index of Secondary Organic Aerosol Generated from Isoprene. <i>Environmental Science & Technology</i> , 2018, 52, 2566-2574.	4.6	19
26	Studying volatility from composition, dilution, and heating measurements of secondary organic aerosols formed during α -pinene ozonolysis. <i>Atmospheric Chemistry and Physics</i> , 2018, 18, 5455-5466.	1.9	16
27	Contributions of Condensable Particulate Matter to Atmospheric Organic Aerosol over Japan. <i>Environmental Science & Technology</i> , 2018, 52, 8456-8466.	4.6	54
28	A Comparison of Particulate-Bound Polycyclic Aromatic Hydrocarbons Long-Range Transported from the Asian Continent to the Noto Peninsula and Fukue Island, Japan. <i>Asian Journal of Atmospheric Environment</i> , 2018, 12, 369-376.	0.4	17
29	Direct observation of new particle formation during ozonolysis of isoprene and ethene competing against the growth of preexisting particles. <i>Atmospheric Environment</i> , 2017, 170, 149-155.	1.9	7
30	Missing ozone-induced potential aerosol formation in a suburban deciduous forest. <i>Atmospheric Environment</i> , 2017, 171, 91-97.	1.9	2
31	Total OH reactivity measurements for the OH-initiated oxidation of aromatic hydrocarbons in the presence of NO _x . <i>Atmospheric Environment</i> , 2017, 171, 272-278.	1.9	15
32	Sensitivities of Simulated Source Contributions and Health Impacts of PM _{2.5} to Aerosol Models. <i>Environmental Science & Technology</i> , 2017, 51, 14273-14282.	4.6	14
33	Aerosol Health Effects from Molecular to Global Scales. <i>Environmental Science & Technology</i> , 2017, 51, 13545-13567.	4.6	384
34	Analysis of Organic Aerosol in Fukuoka, Japan Using a PMF Method. <i>Aerosol and Air Quality Research</i> , 2016, 16, 314-322.	0.9	18
35	Influence of Trans-Boundary Air Pollution on the Urban Atmosphere in Fukuoka, Japan. <i>Atmosphere</i> , 2016, 7, 51.	1.0	18
36	Temperature Effects on Secondary Organic Aerosol (SOA) from the Dark Ozonolysis and Photo-Oxidation of Isoprene. <i>Environmental Science & Technology</i> , 2016, 50, 5564-5571.	4.6	37

#	ARTICLE	IF	CITATIONS
37	Dialdehyde Production during Direct Dissociation of Energy-rich Criegee Intermediates Produced by Ozonolysis of Cycloalkenes. <i>Chemistry Letters</i> , 2016, 45, 916-918.	0.7	1
38	Terpenylic acid and nine-carbon multifunctional compounds formed during the aging of β -pinene ozonolysis secondary organic aerosol. <i>Atmospheric Environment</i> , 2016, 130, 127-135.	1.9	32
39	Gas-phase Ozone Reactions with <i>Z</i> -3-Hexenal and <i>Z</i> -3-Hexen-1-ol: Formation Yields of OH Radical, Propanal, and Ethane. <i>Chemistry Letters</i> , 2015, 44, 457-458.	0.7	5
40	Transboundary secondary organic aerosol in western Japan: An observed limitation of the f44 oxidation indicator. <i>Atmospheric Environment</i> , 2015, 120, 71-75.	1.9	5
41	Evaluation of elemental quantitative values of atmospheric aerosol samples by PIXE method. <i>International Journal of PIXE</i> , 2015, 25, 13-22.	0.4	1
42	Complex refractive index of secondary organic aerosol generated from isoprene/NO _x photooxidation in the presence and absence of SO ₂ . <i>Journal of Geophysical Research D: Atmospheres</i> , 2015, 120, 7777-7787.	1.2	27
43	4-Nitrophenol, 1-nitropyrene, and 9-nitroanthracene emissions in exhaust particles from diesel vehicles with different exhaust gas treatments. <i>Atmospheric Environment</i> , 2015, 110, 93-102.	1.9	35
44	Verification of Chemical Transport Models for PM _{2.5} Chemical Composition Using Simultaneous Measurement Data over Japan. <i>Aerosol and Air Quality Research</i> , 2015, 15, 2009-2023.	0.9	28
45	Transboundary Secondary Organic Aerosol in Western Japan Indicated by the $\delta^{13}C$ of Water-Soluble Organic Carbon and the m/z 44 Signal in Organic Aerosol Mass Spectra. <i>Environmental Science & Technology</i> , 2014, 48, 6273-6281.	4.6	19
46	Analysis of secondary organic aerosols from ozonolysis of isoprene by proton transfer reaction mass spectrometry. <i>Atmospheric Environment</i> , 2014, 97, 397-405.	1.9	53
47	Impact of long-range transport of aerosols on the PM _{2.5} composition at a major metropolitan area in the northern Kyushu area of Japan. <i>Atmospheric Environment</i> , 2014, 97, 416-425.	1.9	79
48	Secondary organic aerosol model intercomparison based on secondary organic aerosol to odd oxygen ratio in Tokyo. <i>Journal of Geophysical Research D: Atmospheres</i> , 2014, 119, 13,489.	1.2	20
49	Characterization of nitromethane emission from automotive exhaust. <i>Atmospheric Environment</i> , 2013, 81, 523-531.	1.9	34
50	On-line measurements of gaseous nitro-organic compounds in diesel vehicle exhaust by proton-transfer-reaction mass spectrometry. <i>Atmospheric Environment</i> , 2013, 73, 195-203.	1.9	38
51	Effect of OH radical scavengers on secondary organic aerosol formation from reactions of isoprene with ozone. <i>Atmospheric Environment</i> , 2013, 79, 147-154.	1.9	30
52	Real-Time Study of Particle-Phase Products from β -Pinene Ozonolysis and Isoprene Photooxidation Using Particle into Liquid Sampling Directly Coupled to a Time-of-Flight Mass Spectrometer (PILS-ToF). <i>Aerosol Science and Technology</i> , 2013, 47, 1374-1382.	1.5	14
53	Wavelength and NO _x dependent complex refractive index of SOAs generated from the photooxidation of toluene. <i>Atmospheric Chemistry and Physics</i> , 2013, 13, 531-545.	1.9	129
54	Transported and Local Organic Aerosols over Fukuoka, Japan. <i>Aerosol and Air Quality Research</i> , 2013, 13, 1263-1272.	0.9	15

#	ARTICLE	IF	CITATIONS
55	AMS and LC/MS analyses of SOA from the photooxidation of benzene and 1,3,5-trimethylbenzene in the presence of NO _x : effects of chemical structure on SOA aging. <i>Atmospheric Chemistry and Physics</i> , 2012, 12, 4667-4682.	1.9	113
56	Kinetic Study of the Daytime Atmospheric Fate of (<i>Z</i>)-3-Hexenal. <i>Journal of Physical Chemistry A</i> , 2012, 116, 8523-8529.	1.1	6
57	Wavelength Dependence of Refractive Index of Secondary Organic Aerosols Generated during the Ozonolysis and Photooxidation of α -Pinene. <i>Scientific Online Letters on the Atmosphere</i> , 2012, 8, 119-123.	0.6	32
58	Secondary organic aerosol formation from phenolic compounds in the absence of NO _x . <i>Atmospheric Chemistry and Physics</i> , 2011, 11, 10649-10660.	1.9	78
59	Secondary organic aerosol formation from the photooxidation of isoprene, 1,3-butadiene, and 2,3-dimethyl-1,3-butadiene under high NO _x conditions. <i>Atmospheric Chemistry and Physics</i> , 2011, 11, 7301-7317.	1.9	40
60	Characterization of Aerosol Particles in the Tokyo Metropolitan Area using Two Different Particle Mass Spectrometers. <i>Aerosol Science and Technology</i> , 2011, 45, 315-326.	1.5	11
61	Aerial Observation of Aerosols Transported from East Asia – Chemical Composition of Aerosols and Layered Structure of an Air Mass over the East China Sea. <i>Aerosol and Air Quality Research</i> , 2011, 11, 497-507.	0.9	29
62	Mass spectrometric study of secondary organic aerosol formed from the photo-oxidation of aromatic hydrocarbons. <i>Atmospheric Environment</i> , 2010, 44, 1080-1087.	1.9	95
63	Laboratory studies on optical properties of secondary organic aerosols generated during the photooxidation of toluene and the ozonolysis of α -pinene. <i>Journal of Geophysical Research</i> , 2010, 115, .	3.3	130
64	Long-range transport of particulate polycyclic aromatic hydrocarbons at Cape Hedo remote island site in the East China Sea between 2005 and 2008. <i>Journal of Atmospheric Chemistry</i> , 2008, 61, 243-257.	1.4	22
65	Detection of nitrooxypolyols in secondary organic aerosol formed from the photooxidation of conjugated dienes under high-NO conditions. <i>Atmospheric Environment</i> , 2008, 42, 6851-6861.	1.9	57
66	Single Particle Analysis of Secondary Organic Aerosols Formed from 1,4-Cyclohexadiene Ozonolysis Using a Laser-Ionization Single-Particle Aerosol Mass Spectrometer. <i>Bulletin of the Chemical Society of Japan</i> , 2008, 81, 120-126.	2.0	7
67	Real-Time Analysis of Secondary Organic Aerosol Particles Formed from Cyclohexene Ozonolysis Using a Laser-Ionization Single-Particle Aerosol Mass Spectrometer. <i>Analytical Sciences</i> , 2007, 23, 507-512.	0.8	18
68	Secondary Organic Aerosol Formation during the Photooxidation of Toluene: Dependence of Chemical Composition. <i>Journal of Physical Chemistry A</i> , 2007, 111, 9796-9808.	1.1	166
69	Production of the radicals in the ozonolysis of ethene: A chamber study by FT-IR and PERCA. <i>Chemical Physics Letters</i> , 2006, 427, 461-465.	1.2	12
70	Chemical Compositions of Secondary Organic Aerosol from the Ozonolysis of Cyclohexene in the Absence of Seed Particles. <i>Chemistry Letters</i> , 2005, 34, 1584-1585.	0.7	9
71	Kinetic measurements for the reactions of ozone with crotonaldehyde and its methyl derivatives and calculations of transition-state theory Electronic supplementary information (ESI) available: The stationary-point geometries optimized at B3LYP/6-31G(d,p) for the reactions of ozone with nine unsaturated carbonyls. See http://www.rsc.org/suppdata/cp/b4/b402496f/ . <i>Physical Chemistry Chemical Physics</i> , 2004, 6, 3969.	1.3	31
72	Secondary Organic Aerosol Formation during the Photo-Oxidation of Toluene: Dependence on Initial Hydrocarbon Concentration. <i>Bulletin of the Chemical Society of Japan</i> , 2004, 77, 667-671.	2.0	27

#	ARTICLE	IF	CITATIONS
73	Cyanomethylene Formation from the Reaction of Excited Nitrogen Atoms with Acetylene: A Crossed Beam and ab Initio Study. <i>Journal of the American Chemical Society</i> , 2000, 122, 4443-4450.	6.6	71
74	Kinetics and mechanisms of the reactions of CH and CD with H ₂ S and D ₂ S. <i>Chemical Physics</i> , 1999, 242, 1-10.	0.9	7
75	Theoretical calculations of thermal rate constants for the N(2D)+acetylene reaction. <i>Chemical Physics Letters</i> , 1999, 312, 503-510.	1.2	9
76	Measurements of Thermal Rate Constants for the Reactions of N(2D,2P) with C ₂ H ₄ and C ₂ D ₄ between 225 and 292 K. <i>Journal of Physical Chemistry A</i> , 1999, 103, 8650-8656.	1.1	27
77	Kinetic Studies on the N(2D, 2P) + CH ₄ and CD ₄ Reactions: The Role of Nonadiabatic Transitions on Thermal Rate Constants. <i>Journal of Physical Chemistry A</i> , 1999, 103, 250-255.	1.1	29
78	Reactions of C(1) with H ₂ , HD and D ₂ : kinetic isotope effect and the CD/CH branching ratio. <i>Chemical Physics</i> , 1998, 237, 195-204.	0.9	47
79	Measurements of Thermal Rate Constants and Theoretical Calculations for the N(2D,2P) + C ₂ H ₂ and C ₂ D ₂ Reactions. <i>Journal of Physical Chemistry A</i> , 1998, 102, 6251-6258.	1.1	38
80	Ab Initio Molecular Orbital Calculations for the N(2D) + Ethylene Reaction. <i>Journal of Physical Chemistry A</i> , 1998, 102, 10391-10398.	1.1	29
81	Ab Initio Molecular Orbital Calculations of the Potential Energy Surfaces for the N(2D) + CH ₄ Reaction. <i>Journal of Physical Chemistry A</i> , 1998, 102, 254-259.	1.1	49
82	Translational energy distributions of the products of the 193 and 157 nm photodissociation of chloroethylenes. <i>Journal of Chemical Physics</i> , 1997, 106, 10123-10133.	1.2	30
83	Infrared multiphoton dissociation of 1,1-dichloroethene. <i>Chemical Physics Letters</i> , 1995, 232, 357-363.	1.2	11
84	Translational distributions of fragments produced in the photodissociation of vinyl fluoride at 157 nm. <i>Chemical Physics Letters</i> , 1995, 242, 401-406.	1.2	27
85	Rotational state distribution of HCl formed in the infrared multiphoton dissociation of trichloroethene. <i>Chemical Physics Letters</i> , 1995, 245, 432-436.	1.2	2
86	Nascent internal state distributions of ZnH(X 2 Σ^+) produced in the reactions of Zn(4 1P ₁) with some alkane hydrocarbons. <i>Journal of Chemical Physics</i> , 1994, 101, 4803-4808.	1.2	21
87	Nascent rotational state distributions of ZnH(X 2 Σ^+) produced in the reactions of Zn(4 1P ₁) with simple alkane hydrocarbons. <i>Chemical Physics Letters</i> , 1993, 214, 271-275.	1.2	8
88	The photodissociation dynamics of dichloroethenes at 214 and 220 nm. <i>Journal of Chemical Physics</i> , 1993, 99, 1703-1709.	1.2	19