## Loredana Poggesi

## List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/4176150/loredana-poggesi-publications-by-year.pdf

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

33<br/>papers760<br/>citations16<br/>h-index27<br/>g-index33<br/>ext. papers868<br/>ext. citations5<br/>avg, IF2.9<br/>L-index

| #  | Paper   | IF   | Citations |
|----|---|------|-----------|
| 33 | Use of the FebriDx point-of-care test for the exclusion of SARS-CoV-2 diagnosis in a population with acute respiratory infection during the second (COVID-19) wave in Italy. <i>International Journal of Infectious Diseases</i> , <b>2021</b> , 108, 231-236 | 10.5 | 4         |
| 32 | Sars-CoV-2 Induced Coagulopathy and Prognosis in Hospitalized Patients: A Snapshot from Italy. <i>Thrombosis and Haemostasis</i> , <b>2020</b> , 120, 1233-1236   | 7    | 10        |
| 31 | Clinical risk score to predict in-hospital mortality in COVID-19 patients: a retrospective cohort study. <i>BMJ Open</i> , <b>2020</b> , 10, e040729  | 3    | 37        |
| 30 | The HScore for secondary hemophagocytic lymphohistiocytosis, calculated without a marrow biopsy, is consistently low in patients with COVID-19. <i>International Journal of Laboratory Hematology</i> , <b>2020</b> , 42, e270-e273                           | 2.5  | 3         |
| 29 | Quantitative and qualitative alterations of circulating myeloid cells and plasmacytoid DC in SARS-CoV-2 infection. <i>Immunology</i> , <b>2020</b> , 161, 345-353   | 7.8  | 42        |
| 28 | T-Lymphocyte-Based Renin Angiotensin System in Obesity. <i>American Journal of the Medical Sciences</i> , <b>2019</b> , 358, 51-58  | 2.2  | 1         |
| 27 | Persistent and selective upregulation of renin-angiotensin system in circulating T lymphocytes in unstable angina. <i>JRAAS - Journal of the Renin-Angiotensin-Aldosterone System</i> , <b>2017</b> , 18, 147032031769  | 8849 | 2         |
| 26 | A challenging multifactorial migraine with aura. Internal and Emergency Medicine, 2014, 9, 201-5  | 3.7  | 2         |
| 25 | Framing comorbidities and co-occurrences in a migraine with aura patient: authorts reply. <i>Internal and Emergency Medicine</i> , <b>2014</b> , 9, 605-6   | 3.7  |           |
| 24 | Adherence to antihypertensive therapy affects Ambulatory Arterial Stiffness Index. <i>European Journal of Internal Medicine</i> , <b>2011</b> , 22, 93-8  | 3.9  | 19        |
| 23 | Erythema ab igne. Internal and Emergency Medicine, 2011, 6, 175-6   | 3.7  | 4         |
| 22 | Ang II Upregulation of the T-lymphocyte renin-angiotensin system is amplified by low-grade inflammation in human hypertension. <i>American Journal of Hypertension</i> , <b>2011</b> , 24, 716-23   | 2.3  | 28        |
| 21 | Serum uric acid levels and renal damage in hyperuricemic hypertensive patients treated with renin-angiotensin system blockers. <i>American Journal of Hypertension</i> , <b>2010</b> , 23, 675-80   | 2.3  | 15        |
| 20 | Non-invasive tissue Doppler imaging pulmonary capillary wedge pressure measurement improves NT-proBNP prognostic value in heart failure. <i>Acta Cardiologica</i> , <b>2009</b> , 64, 213-8   | 0.9  | 1         |
| 19 | Angiotensin II upregulates renin-angiotensin system in human isolated T lymphocytes. <i>Regulatory Peptides</i> , <b>2008</b> , 151, 1-6  |      | 20        |
| 18 | Renal resistive index early detects chronic tubulointerstitial nephropathy in normo- and hypertensive patients. <i>American Journal of Nephrology</i> , <b>2006</b> , 26, 16-21   | 4.6  | 42        |
| 17 | Exaggerated local hand sympathetic but not renin-angiotensin system activation in patients with primary Raynauds phenomenon. <i>Microvascular Research</i> , <b>2006</b> , 71, 128-34   | 3.7  | 1         |

## LIST OF PUBLICATIONS

| 16 | Thromboxane inhibition improves renal perfusion and excretory function in severe congestive heart failure. <i>Journal of the American College of Cardiology</i> , <b>2003</b> , 42, 133-9                        | 15.1 | 14  |
|----|--|------|-----|
| 15 | Activation of cardiac renin-angiotensin system in unstable angina. <i>Journal of the American College of Cardiology</i> , <b>2001</b> , 38, 49-55  | 15.1 | 18  |
| 14 | Renal endothelin in heart failure and its relation to sodium excretion. <i>American Heart Journal</i> , <b>2000</b> , 140, 617-22  | 4.9  | 24  |
| 13 | Endothelin-1 urinary excretion, but not endothelin-1 plasma concentration, is increased in renovascular hypertension. <i>Translational Research</i> , <b>1999</b> , 134, 386-91                                  |      | 9   |
| 12 | Human vascular renin-angiotensin system and its functional changes in relation to different sodium intakes. <i>Hypertension</i> , <b>1998</b> , 31, 836-42   | 8.5  | 70  |
| 11 | Detection of malignant thymoma during myocardial perfusion tomography with Tc-99m sestamibi: potential implications for tumor evaluation and staging. <i>Clinical Nuclear Medicine</i> , <b>1998</b> , 23, 842-3 | 1.7  | 2   |
| 10 | Increased renal formation of thromboxane A2 and prostaglandin F2 alpha in heart failure. <i>American Heart Journal</i> , <b>1997</b> , 133, 94-100   | 4.9  | 22  |
| 9  | Acute T-cell activation is detectable in unstable angina. <i>Circulation</i> , <b>1997</b> , 95, 1806-12   | 16.7 | 126 |
| 8  | Evidence for the existence of a functional cardiac renin-angiotensin system in humans. <i>Circulation</i> , <b>1996</b> , 94, 1886-93  | 16.7 | 50  |
| 7  | Enhanced cardiac norepinephrine release in patients with idiopathic ventricular tachycardia related to the occurrence of arrhythmias. <i>American Heart Journal</i> , <b>1994</b> , 127, 686-9                   | 4.9  | 1   |
| 6  | Evaluation of mitral stenosis by cine magnetic resonance imaging. <i>American Heart Journal</i> , <b>1992</b> , 123, 1252-60   | 4.9  | 14  |
| 5  | Detection of apical hypertrophic cardiomyopathy by magnetic resonance imaging. <i>American Heart Journal</i> , <b>1989</b> , 117, 468-72   | 4.9  | 16  |
| 4  | Acute reversible reduction of PGI2 platelet receptors after iloprost infusion in man. <i>Thrombosis Research</i> , <b>1987</b> , 48, 663-9   | 8.2  | 26  |
| 3  | ECG-gated magnetic resonance imaging in right ventricular dysplasia. <i>American Heart Journal</i> , <b>1987</b> , 113, 1245-8   | 4.9  | 55  |
| 2  | Reduced prostacyclin production in patients with different manifestations of ischemic heart disease. <i>American Journal of Cardiology</i> , <b>1982</b> , 49, 1146-51   | 3    | 53  |
| 1  | Release of prostacyclin into the bloodstream and its exhaustion in humans after local blood flow changes (ischemia and venous stasis). <i>Thrombosis Research</i> , <b>1980</b> , 17, 197-208                    | 8.2  | 29  |