

Wade H Powell

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/4180235/publications.pdf>

Version: 2024-02-01

19
papers

733
citations

759233

12
h-index

839539

18
g-index

21
all docs

21
docs citations

21
times ranked

574
citing authors

#	ARTICLE	IF	CITATIONS
1	Dioxin Disrupts Thyroid Hormone and Glucocorticoid Induction of <i>klf9</i> , a Master Regulator of Frog Metamorphosis. <i>Toxicological Sciences</i> , 2022, 187, 150-161.	3.1	2
2	An aryl hydrocarbon receptor from the caecilian <i>Gymnopsis multiplicata</i> suggests low dioxin affinity in the ancestor of all three amphibian orders. <i>General and Comparative Endocrinology</i> , 2020, 299, 113592.	1.8	3
3	Dioxin Exposure Alters Molecular and Morphological Responses to Thyroid Hormone in <i>Xenopus laevis</i> Cultured Cells and Prometamorphic Tadpoles. <i>Toxicological Sciences</i> , 2018, 161, 196-206.	3.1	12
4	Subfunctionalization of Paralogous Aryl Hydrocarbon Receptors from the Frog <i>Xenopus Laevis</i> : Distinct Target Genes and Differential Responses to Specific Agonists in a Single Cell Type. <i>Toxicological Sciences</i> , 2017, 155, 337-347.	3.1	5
5	An Aryl Hydrocarbon Receptor from the Salamander <i>Ambystoma mexicanum</i> Exhibits Low Sensitivity to 2,3,7,8-Tetrachlorodibenzo-p-dioxin. <i>Environmental Science & Technology</i> , 2015, 49, 6993-7001.	10.0	13
6	Specific Ligand Binding Domain Residues Confer Low Dioxin Responsiveness to AHR1 ² of <i>Xenopus laevis</i> . <i>Biochemistry</i> , 2013, 52, 1746-1754.	2.5	22
7	Induction of cytochrome P450 family 1 mRNAs and activities in a cell line from the frog <i>Xenopus laevis</i> . <i>Aquatic Toxicology</i> , 2012, 114-115, 165-172.	4.0	15
8	Responsiveness of a <i>Xenopus laevis</i> cell line to the aryl hydrocarbon receptor ligands 6-formylindolo[3,2-b]carbazole (FICZ) and 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). <i>Chemico-Biological Interactions</i> , 2010, 183, 202-211.	4.0	28
9	An Aryl Hydrocarbon Receptor Repressor from <i>Xenopus laevis</i> : Function, Expression, and Role in Dioxin Responsiveness during Frog Development. <i>Toxicological Sciences</i> , 2008, 104, 124-134.	3.1	17
10	Developmental differences in elimination of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) during <i>Xenopus laevis</i> development. <i>Marine Environmental Research</i> , 2006, 62, S34-S37.	2.5	8
11	Aryl Hydrocarbon Receptors in the Frog <i>Xenopus laevis</i> : Two AhR1 Paralogs Exhibit Low Affinity for 2,3,7,8-Tetrachlorodibenzo-p-Dioxin (TCDD). <i>Toxicological Sciences</i> , 2005, 88, 60-72.	3.1	58
12	Cloning and analysis of the CYP1A promoter from the atlantic killifish (<i>Fundulus heteroclitus</i>). <i>Marine Environmental Research</i> , 2004, 58, 119-124.	2.5	20
13	ARNT gene multiplicity in amphibians: Characterization of <i>arnt2</i> from the frog <i>Xenopus laevis</i> . <i>The Journal of Experimental Zoology</i> , 2003, 300B, 48-57.	1.4	17
14	Regulatory Interactions among Three Members of the Vertebrate Aryl Hydrocarbon Receptor Family: AHR Repressor, AHR1, and AHR2. <i>Journal of Biological Chemistry</i> , 2002, 277, 6949-6959.	3.4	119
15	Identification and functional characterization of hypoxia-inducible factor 2 β from the estuarine teleost, <i>Fundulus heteroclitus</i> : Interaction of HIF-2 β with two ARNT2 splice variants. <i>The Journal of Experimental Zoology</i> , 2002, 294, 17-29.	1.4	44
16	An aryl hydrocarbon receptor (AHR) homologue from the soft-shell clam, <i>Mya arenaria</i> : evidence that invertebrate AHR homologues lack 2,3,7,8-tetrachlorodibenzo-p-dioxin and 1 β -naphthoflavone binding. <i>Gene</i> , 2001, 278, 223-234.	2.2	151
17	Identification and Functional Characterization of Two Highly Divergent Aryl Hydrocarbon Receptors (AHR1 and AHR2) in the Teleost <i>Fundulus heteroclitus</i> . <i>Journal of Biological Chemistry</i> , 1999, 274, 33814-33824.	3.4	146
18	Functional Diversity of Vertebrate ARNT Proteins: Identification of ARNT2 as the Predominant Form of ARNT in the Marine Teleost, <i>Fundulus heteroclitus</i> . <i>Archives of Biochemistry and Biophysics</i> , 1999, 361, 156-163.	3.0	53

#	ARTICLE	IF	CITATIONS
19	Glutamic acid-371 of the barnase homology domain in RNA polymerase II is not required for SII-activated RNA cleavage. <i>Molecular Genetics and Genomics</i> , 1997, 253, 507-511.	2.4	0