Tomas Urbina

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/4189288/publications.pdf

Version: 2024-02-01

1040056 940533 17 941 9 16 citations h-index g-index papers 17 17 17 2539 citing authors all docs docs citations times ranked

#	Article	IF	CITATIONS
1	Effect of Tocilizumab vs Usual Care in Adults Hospitalized With COVID-19 and Moderate or Severe Pneumonia. JAMA Internal Medicine, 2021, 181, 32.	5.1	654
2	Narrative review: clinical assessment of peripheral tissue perfusion in septic shock. Annals of Intensive Care, 2019, 9, 37.	4.6	95
3	Elevated plasma IL-6 and CRP levels are associated with adverse clinical outcomes and death in critically ill SARS-CoV-2 patients: inflammatory response of SARS-CoV-2 patients. Annals of Intensive Care, 2021, 11, 9.	4.6	70
4	Iron Supplementation–Induced Phosphaturic Osteomalacia: FGF23 is the Culprit. Journal of Bone and Mineral Research, 2018, 33, 540-542.	2.8	22
5	Vitamin C improves microvascular reactivity and peripheral tissue perfusion in septic shock patients. Critical Care, 2022, 26, 25.	5.8	20
6	Antibiotics in Necrotizing Soft Tissue Infections. Antibiotics, 2021, 10, 1104.	3.7	17
7	Impact of a multidisciplinary care bundle for necrotizing skin and soft tissue infections: a retrospective cohort study. Annals of Intensive Care, 2019, 9, 123.	4.6	16
8	Understanding necrotizing soft tissue infections in the intensive care unit. Intensive Care Medicine, 2020, 46, 1739-1742.	8.2	12
9	Long-term quality of life in necrotizing soft-tissue infection survivors: a monocentric prospective cohort study. Annals of Intensive Care, 2021, 11, 102.	4.6	12
10	Early identification of patients at high risk of group A streptococcus-associated necrotizing skin and soft tissue infections: a retrospective cohort study. Critical Care, 2019, 23, 417.	5.8	6
11	Tissue perfusion alterations correlate with mortality in patients admitted to the intensive care unit for acute pulmonary embolism. Medicine (United States), 2018, 97, e11993.	1.0	5
12	SARS-CoV-2 Genomic Characteristics and Clinical Impact of SARS-CoV-2 Viral Diversity in Critically III COVID-19 Patients: A Prospective Multicenter Cohort Study. Viruses, 2022, 14, 1529.	3.3	4
13	Endotracheal intubation rate is lower in critically-ill SARS-CoV-2 patients requiring high-flow nasal oxygen receiving additional face-mask noninvasive ventilation: a retrospective bicentric cohort with propensity score analysis. Minerva Anestesiologica, 2022, , .	1.0	3
14	Earlobe Crease in Patient With Chest Pain. Annals of Emergency Medicine, 2017, 69, 664-672.	0.6	2
15	COVID-19-associated Multi-Inflammatory Syndrome in Adults (MIS-A): look into the eyes!. Intensive Care Medicine, 2022, 48, 947-948.	8.2	2
16	Delayed inflammation decrease is associated with mortality in Tocilizumab-treated critically ill SARS-CoV-2 patients: A retrospective matched-cohort analysis. Innate Immunity, 2022, 28, 3-10.	2.4	1
17	Author's reply to "Face-mask non-invasive ventilation plus high-flow nasal oxygen in COVID-19 patients: "one size fits all" or tailored approach?": always tailored, with a trial!. Minerva Anestesiologica, 2022, 88, .	1.0	0