

# Keiko Tanaka

## List of Publications by Year in Descending Order

**Source:** <https://exaly.com/author-pdf/4190587/keiko-tanaka-publications-by-year.pdf>

**Version:** 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

187  
papers

4,461  
citations

37  
h-index

54  
g-index

190  
ext. papers

5,034  
ext. citations

3.7  
avg, IF

5.3  
L-index

#	Paper	IF	Citations
187	Maternal Use of Induction Heating Cookers During Pregnancy and Birth Outcomes: The Kyushu Okinawa Maternal and Child Health Study. <i>Bioelectromagnetics</i> , <b>2021</b> , 42, 329-335	1.6	2
186	Maternal consumption of soy and isoflavones during pregnancy and risk of childhood behavioural problems: the Kyushu Okinawa Maternal and Child Health Study. <i>International Journal of Food Sciences and Nutrition</i> , <b>2021</b> , 72, 1118-1127	3.7	1
185	Hypertension and dyslipidemia are risk factors for herpes zoster in patients with rheumatoid arthritis: a retrospective analysis using a medical information database. <i>Rheumatology International</i> , <b>2021</b> , 41, 1633-1639	3.6	1
184	Calcium intake during pregnancy is associated with decreased risk of emotional and hyperactivity problems in five-year-old Japanese children. <i>Nutritional Neuroscience</i> , <b>2021</b> , 24, 762-769	3.6	3
183	Cesarean section is associated with increased risk of postpartum depressive symptoms in Japan: the Kyushu Okinawa Maternal and Child Health Study. <i>Journal of Affective Disorders</i> , <b>2021</b> , 278, 497-501	6.6	0
182	Maternal caffeine intake during pregnancy and risk of food allergy in young Japanese children. <i>Journal of Paediatrics and Child Health</i> , <b>2021</b> , 57, 903-907	1.3	0
181	Maternal metal intake during pregnancy and childhood behavioral problems in Japan: the Kyushu Okinawa Maternal and Child Health Study. <i>Nutritional Neuroscience</i> , <b>2021</b> , 1-9	3.6	1
180	Perinatal maternal depressive symptoms and risk of behavioral problems at five years. <i>Pediatric Research</i> , <b>2021</b> ,	3.2	2
179	Higher number of teeth is associated with decreased prevalence of hearing impairment in Japan. <i>Archives of Gerontology and Geriatrics</i> , <b>2021</b> , 97, 104502	4	0
178	Dietary intake of vegetables, fruit, and antioxidants and risk of ulcerative colitis: A case-control study in Japan. <i>Nutrition</i> , <b>2021</b> , 91-92, 111378	4.8	2
177	BAIAP2 rs8079781, postnatal smoking exposure, and emotional problems in Japanese children aged 5 years: the Kyushu Okinawa Maternal and Child Health Study. <i>Journal of Neural Transmission</i> , <b>2020</b> , 127, 1081-1087	4.3	
176	Maternal prenatal stress and infantile wheeze and asthma: The Osaka Maternal and Child Health Study. <i>Journal of Psychosomatic Research</i> , <b>2020</b> , 135, 110143	4.1	3
175	Associations of job type, income, and education with postpartum depressive symptoms: The Kyushu Okinawa Maternal and Child Health Study. <i>Psychiatry Research</i> , <b>2020</b> , 291, 113224	9.9	3
174	Perinatal smoking exposure and risk of asthma in the first three years of life: A prospective prebirth cohort study. <i>Allergologia Et Immunopathologia</i> , <b>2020</b> , 48, 530-536	1.9	0
173	Hearing Impairment and Prevalence of Mild Cognitive Impairment in Japan: Baseline Data From the Aikai Cohort Study in Yawatahama and Uchiko. <i>Ear and Hearing</i> , <b>2020</b> , 41, 254-258	3.4	6
172	Maternal B vitamin intake during pregnancy and childhood behavioral problems in Japan: The Kyushu Okinawa Maternal and Child Health Study. <i>Nutritional Neuroscience</i> , <b>2020</b> , 23, 706-713	3.6	12
171	Maternal consumption of vegetables, fruit, and antioxidants during pregnancy and risk for childhood behavioral problems. <i>Nutrition</i> , <b>2020</b> , 69, 110572	4.8	4

170	Preterm birth is associated with higher prevalence of wheeze and asthma in a selected population of Japanese children aged three years. <i>Allergologia Et Immunopathologia</i> , <b>2019</b> , 47, 425-430	1.9	5
169	Case-Control Study of Idiopathic Parkinson Disease in Japan. <i>Current Topics in Environmental Health and Preventive Medicine</i> , <b>2019</b> , 61-81	0.3	
168	Association of household income and education with prevalence of hearing impairment in Japan. <i>Laryngoscope</i> , <b>2019</b> , 129, 2153-2157	3.6	1
167	Maternal caffeine intake in pregnancy is inversely related to childhood peer problems in Japan: The Kyushu Okinawa Maternal and Child Health Study. <i>Nutritional Neuroscience</i> , <b>2019</b> , 22, 817-824	3.6	7
166	Maternal fat intake during pregnancy and behavioral problems in 5-y-old Japanese children. <i>Nutrition</i> , <b>2018</b> , 50, 91-96	4.8	6
165	High birthweight is associated with increased prevalence of dental caries in Japanese children. <i>International Journal of Dental Hygiene</i> , <b>2018</b> , 16, 404-410	2.6	6
164	Soy isoflavone intake and prevalence of depressive symptoms during pregnancy in Japan: baseline data from the Kyushu Okinawa Maternal and Child Health Study. <i>European Journal of Nutrition</i> , <b>2018</b> , 57, 441-450	5.2	24
163	Nocturia and Prevalence of Depressive Symptoms in Japanese Adult Patients With Type 2 Diabetes Mellitus: The Dogo Study. <i>Canadian Journal of Diabetes</i> , <b>2018</b> , 42, 51-55	2.1	5
162	Physical activity and prevalence of erectile dysfunction in Japanese patients with type 2 diabetes mellitus: The Dogo Study. <i>Journal of Diabetes Investigation</i> , <b>2018</b> , 9, 193-198	3.9	9
161	Dietary intake habits and the prevalence of nocturia in Japanese patients with type 2 diabetes mellitus. <i>Journal of Diabetes Investigation</i> , <b>2018</b> , 9, 279-285	3.9	5
160	Dietary patterns and depressive symptoms during pregnancy in Japan: Baseline data from the Kyushu Okinawa Maternal and Child Health Study. <i>Journal of Affective Disorders</i> , <b>2018</b> , 225, 552-558	6.6	9
159	Breastfeeding duration is inversely associated with asthma in Japanese children aged 3 years. <i>Journal of Asthma</i> , <b>2018</b> , 55, 511-516	1.9	5
158	ITIH3 and ITIH4 polymorphisms and depressive symptoms during pregnancy in Japan: the Kyushu Okinawa Maternal and Child Health Study. <i>Journal of Neural Transmission</i> , <b>2018</b> , 125, 1503-1509	4.3	3
157	Association between secondhand smoke exposure and early eruption of deciduous teeth: A cross-sectional study. <i>Tobacco Induced Diseases</i> , <b>2018</b> , 16, 04	3.2	0
156	Manganese intake is inversely associated with depressive symptoms during pregnancy in Japan: Baseline data from the Kyushu Okinawa Maternal and Child Health Study. <i>Journal of Affective Disorders</i> , <b>2017</b> , 211, 124-129	6.6	9
155	Vitamin D receptor gene polymorphisms, smoking, and risk of sporadic Parkinson's disease in Japan. <i>Neuroscience Letters</i> , <b>2017</b> , 643, 97-102	3.3	13
154	Letter to the Editor. <i>European Journal of Nutrition</i> , <b>2017</b> , 56, 1787	5.2	
153	Parental occupations, educational levels, and income and prevalence of dental caries in 3-year-old Japanese children. <i>Environmental Health and Preventive Medicine</i> , <b>2017</b> , 22, 80	4.2	30

152	Pre- and Postnatal Smoking Exposure and Risk of Atopic Eczema in Young Japanese Children: A Prospective Prebirth Cohort Study. <i>Nicotine and Tobacco Research</i> , <b>2017</b> , 19, 804-809	4.9	12
151	Obesity and the prevalence of nocturia in Japanese elderly patients with type 2 diabetes mellitus: The Dogo study. <i>Geriatrics and Gerontology International</i> , <b>2017</b> , 17, 2460-2465	2.9	5
150	Reply to letter to the editor to "Soy isoflavone intake and prevalence of depressive symptoms during pregnancy in Japan: baseline data from the Kyushu Okinawa Maternal and Child Health Study". <i>European Journal of Nutrition</i> , <b>2017</b> , 56, 1791-1792	5.2	
149	Self-reported sitting time and prevalence of erectile dysfunction in Japanese patients with type 2 diabetes mellitus: The Dogo Study. <i>Journal of Diabetes and Its Complications</i> , <b>2017</b> , 31, 53-57	3.2	5
148	Rate of eating in early life is positively associated with current and later body mass index among young Japanese children: the Osaka Maternal and Child Health Study. <i>Nutrition Research</i> , <b>2017</b> , 37, 20-28	4	14
147	The IL18 Promoter Polymorphism, rs1946518, Is Associated with the Risk of Periodontitis in Japanese Women: The Kyushu Okinawa Maternal and Child Health Study. <i>Tohoku Journal of Experimental Medicine</i> , <b>2017</b> , 243, 159-164	2.4	8
146	Secondhand smoke exposure and risk of wheeze in early childhood: a prospective pregnancy birth cohort study. <i>Tobacco Induced Diseases</i> , <b>2017</b> , 15, 30	3.2	10
145	Smoking and secondhand smoke exposure and prevalence of depressive symptoms during pregnancy in Japan: baseline data from the Kyushu Okinawa Maternal and Child Health Study. <i>Tobacco Induced Diseases</i> , <b>2017</b> , 15, 34	3.2	6
144	Perinatal smoking exposure and behavioral problems in Japanese children aged 5 years: The Kyushu Okinawa Maternal and Child Health Study. <i>Environmental Research</i> , <b>2016</b> , 151, 383-388	7.9	7
143	Milk intake during pregnancy is inversely associated with the risk of postpartum depressive symptoms in Japan: the Kyushu Okinawa Maternal and Child Health Study. <i>Nutrition Research</i> , <b>2016</b> , 36, 907-913	4	11
142	Nocturia and prevalence of erectile dysfunction in Japanese patients with type 2 diabetes mellitus: The Dogo Study. <i>Journal of Diabetes Investigation</i> , <b>2016</b> , 7, 786-90	3.9	10
141	PARK16 polymorphisms, interaction with smoking, and sporadic Parkinson's disease in Japan. <i>Journal of the Neurological Sciences</i> , <b>2016</b> , 362, 47-52	3.2	7
140	Feeding practices in early life and later intake of fruit and vegetables among Japanese toddlers: the Osaka Maternal and Child Health Study. <i>Public Health Nutrition</i> , <b>2016</b> , 19, 650-7	3.3	20
139	Early sugar-sweetened beverage consumption frequency is associated with poor quality of later food and nutrient intake patterns among Japanese young children: the Osaka Maternal and Child Health Study. <i>Nutrition Research</i> , <b>2016</b> , 36, 594-602	4	6
138	Serum lipoprotein(a) levels and diabetic nephropathy among Japanese patients with type 2 diabetes mellitus. <i>Journal of Diabetes and Its Complications</i> , <b>2016</b> , 30, 923-7	3.2	11
137	Microvascular Complications and Prevalence of Nocturia in Japanese Patients With Type 2 Diabetes Mellitus: The Dogo Study. <i>Urology</i> , <b>2016</b> , 93, 147-51	1.6	5
136	Age and Prevalence of Esophageal Reflux Disease in Japanese Patients with Type 2 Diabetes Mellitus: The Dogo Study. <i>Digestive Diseases and Sciences</i> , <b>2016</b> , 61, 3530-3536	4	3
135	Association Between TSLP Polymorphisms and Eczema in Japanese Women: the Kyushu Okinawa Maternal and Child Health Study. <i>Inflammation</i> , <b>2015</b> , 38, 1663-8	5.1	8

134	IL5RA polymorphisms, smoking and eczema in Japanese women: the Kyushu Okinawa Maternal and Child Health Study. <i>International Journal of Immunogenetics</i> , <b>2015</b> , 42, 52-7	2.3	0
133	Authors' reply. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , <b>2015</b> , 122, 586-7	3.7	
132	Intake of dairy products and calcium and prevalence of depressive symptoms during pregnancy in Japan: a cross-sectional study. <i>BJOG: an International Journal of Obstetrics and Gynaecology</i> , <b>2015</b> , 122, 336-43	3.7	41
131	Higher vitamin D intake during pregnancy is associated with reduced risk of dental caries in young Japanese children. <i>Annals of Epidemiology</i> , <b>2015</b> , 25, 620-5	6.4	28
130	Maternal total caffeine intake, mainly from Japanese and Chinese tea, during pregnancy was associated with risk of preterm birth: the Osaka Maternal and Child Health Study. <i>Nutrition Research</i> , <b>2015</b> , 35, 309-16	4	38
129	The linkage among ambulance transports, death and climate parameters in Asahikawa City, Japan. <i>Environmental Health and Preventive Medicine</i> , <b>2015</b> , 20, 63-7	4.2	5
128	Evaluation of psychological distress using the K6 in patients on chronic hemodialysis. <i>Environmental Health and Preventive Medicine</i> , <b>2015</b> , 20, 102-7	4.2	3
127	Dietary vitamin D intake and prevalence of depressive symptoms during pregnancy in Japan. <i>Nutrition</i> , <b>2015</b> , 31, 160-5	4.8	27
126	Association of prenatal exposure to maternal smoking and postnatal exposure to household smoking with dental caries in 3-year-old Japanese children. <i>Environmental Research</i> , <b>2015</b> , 143, 148-53	7.9	19
125	Alcohol consumption during pregnancy and birth outcomes: the Kyushu Okinawa Maternal and Child Health Study. <i>BMC Pregnancy and Childbirth</i> , <b>2014</b> , 14, 79	3.2	26
124	IL3 rs40401 polymorphism and interaction with smoking in risk of asthma in Japanese women: the Kyushu Okinawa Maternal and Child Health study. <i>Scandinavian Journal of Immunology</i> , <b>2014</b> , 79, 410-4	3.4	14
123	Relationship between IL1 gene polymorphisms and periodontal disease in Japanese women. <i>DNA and Cell Biology</i> , <b>2014</b> , 33, 227-33	3.6	13
122	Association between 17q12-21 variants and asthma in Japanese women: rs11650680 polymorphism as potential genetic marker for asthma. <i>DNA and Cell Biology</i> , <b>2014</b> , 33, 531-6	3.6	2
121	Case-control study of rhinoconjunctivitis associated with IL5RA polymorphisms in Japanese women: the Kyushu Okinawa Maternal and Child Health Study. <i>Cytokine</i> , <b>2014</b> , 65, 138-42	4	2
120	Low birth weight, preterm birth or small-for-gestational-age are not associated with dental caries in young Japanese children. <i>BMC Oral Health</i> , <b>2014</b> , 14, 38	3.7	18
119	Daily smoking may independently predict caries development in adults. <i>Journal of Evidence-based Dental Practice</i> , <b>2014</b> , 14, 151-3	1.9	4
118	Maternal consumption of dairy products, calcium, and vitamin D during pregnancy and infantile allergic disorders. <i>Annals of Allergy, Asthma and Immunology</i> , <b>2014</b> , 113, 82-7	3.2	45
117	Prognosis of primary aldosteronism in Japan: results from a nationwide epidemiological study. <i>Endocrine Journal</i> , <b>2014</b> , 61, 35-40	2.9	32

116	Seaweed consumption and prevalence of depressive symptoms during pregnancy in Japan: Baseline data from the Kyushu Okinawa Maternal and Child Health Study. <i>BMC Pregnancy and Childbirth</i> , <b>2014</b> , 14, 301	3.2	19
115	Dietary patterns in infancy and their associations with maternal socio-economic and lifestyle factors among 758 Japanese mother-child pairs: the Osaka Maternal and Child Health Study. <i>Maternal and Child Nutrition</i> , <b>2014</b> , 10, 213-25	3.4	28
114	The role of tobacco use on dental care and oral disease severity within community dental clinics in Japan. <i>Tobacco Induced Diseases</i> , <b>2013</b> , 11, 13	3.2	8
113	Salivary cotinine concentrations and prevalence of periodontal disease in young Japanese women: the Kyushu Okinawa maternal and child health study. <i>Journal of Periodontology</i> , <b>2013</b> , 84, 1724-9	4.6	3
112	IL3 SNP rs40401 variant is a risk factor for rhinoconjunctivitis in Japanese women: the Kyushu Okinawa maternal and child health study. <i>Cytokine</i> , <b>2013</b> , 64, 86-9	4	2
111	Socioeconomic status and risk of dental caries in Japanese preschool children: the Osaka Maternal and child health study. <i>Journal of Public Health Dentistry</i> , <b>2013</b> , 73, 217-23	1.6	28
110	Active and passive maternal smoking during pregnancy and birth outcomes: the Kyushu Okinawa maternal and child health study. <i>BMC Pregnancy and Childbirth</i> , <b>2013</b> , 13, 157	3.2	42
109	Maternal fat intake during pregnancy and wheeze and eczema in Japanese infants: the Kyushu Okinawa Maternal and Child Health Study. <i>Annals of Epidemiology</i> , <b>2013</b> , 23, 674-80	6.4	29
108	Fish and fat intake and prevalence of depressive symptoms during pregnancy in Japan: baseline data from the Kyushu Okinawa Maternal and Child Health Study. <i>Journal of Psychiatric Research</i> , <b>2013</b> , 47, 572-8	5.2	57
107	MDR1 C3435T polymorphism and interaction with environmental factors in risk of Parkinson's disease: a case-control study in Japan. <i>Drug Metabolism and Pharmacokinetics</i> , <b>2013</b> , 28, 138-43	2.2	14
106	Case-control study of eczema in relation to IL4R1 genetic polymorphisms in Japanese women: The Kyushu Okinawa Maternal and Child Health Study. <i>Scandinavian Journal of Immunology</i> , <b>2013</b> , 77, 413-8	3.4	7
105	Lack of relationship between birth conditions and allergic disorders in Japanese children aged 3 years. <i>Journal of Asthma</i> , <b>2013</b> , 50, 555-9	1.9	17
104	VDR gene polymorphisms, interaction with smoking and risk of periodontal disease in Japanese women: the Kyushu Okinawa maternal and child health study. <i>Scandinavian Journal of Immunology</i> , <b>2013</b> , 78, 371-7	3.4	16
103	Active and passive smoking and prevalence of periodontal disease in young Japanese women. <i>Journal of Periodontal Research</i> , <b>2013</b> , 48, 600-5	4.3	15
102	Characteristics of under- and over-reporters of energy intake among Japanese children and adolescents: The Ryukyus Child Health Study. <i>Nutrition</i> , <b>2012</b> , 28, 532-8	4.8	36
101	Dietary patterns and risk of Parkinson's disease: a case-control study in Japan. <i>European Journal of Neurology</i> , <b>2012</b> , 19, 681-8	6	29
100	Smoking and prevalence of allergic disorders in Japanese pregnant women: baseline data from the Kyushu Okinawa Maternal and Child Health Study. <i>Environmental Health</i> , <b>2012</b> , 11, 15	6	8
99	Self-reported rate of eating and risk of overweight in Japanese children: Ryukyus Child Health Study. <i>Journal of Nutritional Science and Vitaminology</i> , <b>2012</b> , 58, 247-52	1.1	24

98	Polymorphisms in the IL4 gene, smoking, and rhinoconjunctivitis in Japanese women: the Kyushu Okinawa Maternal and Child Health Study. <i>Human Immunology</i> , <b>2012</b> , 73, 1046-9	2.3	5
97	SNCA polymorphisms, smoking, and sporadic Parkinson's disease in Japanese. <i>Parkinsonism and Related Disorders</i> , <b>2012</b> , 18, 557-61	3.6	25
96	ADAM33 genetic polymorphisms, smoking, and rhinoconjunctivitis in Japanese women: the Kyushu Okinawa Maternal and Child Health Study. <i>Human Immunology</i> , <b>2012</b> , 73, 411-5	2.3	6
95	UCHL1 S18Y variant is a risk factor for Parkinson's disease in Japan. <i>BMC Neurology</i> , <b>2012</b> , 12, 62	3.1	16
94	Dietary meat and fat intake and prevalence of rhinoconjunctivitis in pregnant Japanese women: baseline data from the Kyushu Okinawa Maternal and Child Health Study. <i>Nutrition Journal</i> , <b>2012</b> , 11, 19	4.3	5
93	Dairy products and calcium intake during pregnancy and dental caries in children. <i>Nutrition Journal</i> , <b>2012</b> , 11, 33	4.3	11
92	An energy-dense diet is cross-sectionally associated with an increased risk of overweight in male children, but not in female children, male adolescents, or female adolescents in Japan: the Ryukyus Child Health Study. <i>Nutrition Research</i> , <b>2012</b> , 32, 486-94	4	9
91	Lack of association between BST1 polymorphisms and sporadic Parkinson's disease in a Japanese population. <i>Journal of the Neurological Sciences</i> , <b>2012</b> , 323, 162-6	3.2	13
90	Employment, income, and education and prevalence of depressive symptoms during pregnancy: the Kyushu Okinawa Maternal and Child Health Study. <i>BMC Psychiatry</i> , <b>2012</b> , 12, 117	4.2	14
89	Parental employment, income, education and allergic disorders in children: a prebirth cohort study in Japan. <i>International Journal of Tuberculosis and Lung Disease</i> , <b>2012</b> , 16, 756-61	2.1	11
88	Maternal dietary patterns in pregnancy and fetal growth in Japan: the Osaka Maternal and Child Health Study. <i>British Journal of Nutrition</i> , <b>2012</b> , 107, 1526-33	3.6	68
87	ADAM33 polymorphisms, smoking and asthma in Japanese women: the Kyushu Okinawa Maternal and Child Health Study. <i>International Journal of Tuberculosis and Lung Disease</i> , <b>2012</b> , 16, 974-9	2.1	12
86	Dairy food, calcium and vitamin D intake and prevalence of allergic disorders in pregnant Japanese women. <i>International Journal of Tuberculosis and Lung Disease</i> , <b>2012</b> , 16, 255-61	2.1	11
85	Association between breastfeeding and dental caries in Japanese children. <i>Journal of Epidemiology</i> , <b>2012</b> , 22, 72-7	3.4	21
84	Dietary intake of metals and risk of Parkinson's disease: a case-control study in Japan. <i>Journal of the Neurological Sciences</i> , <b>2011</b> , 306, 98-102	3.2	55
83	Lack of association of mercury with risk of wheeze and eczema in Japanese children: the Osaka Maternal and Child Health Study. <i>Environmental Research</i> , <b>2011</b> , 111, 1180-4	7.9	24
82	Lack of association of dairy food, calcium, and vitamin D intake with the risk of Parkinson's disease: a case-control study in Japan. <i>Parkinsonism and Related Disorders</i> , <b>2011</b> , 17, 112-6	3.6	33
81	Intake of Japanese and Chinese teas reduces risk of Parkinson's disease. <i>Parkinsonism and Related Disorders</i> , <b>2011</b> , 17, 446-50	3.6	53

80	Does secondhand smoke affect the development of dental caries in children? A systematic review. <i>International Journal of Environmental Research and Public Health</i> , <b>2011</b> , 8, 1503-19	4.6	37
79	Maternal B vitamin intake during pregnancy and wheeze and eczema in Japanese infants aged 16-24 months: the Osaka Maternal and Child Health Study. <i>Pediatric Allergy and Immunology</i> , <b>2011</b> , 22, 69-74	4.2	39
78	Maternal dietary patterns during pregnancy and risk of wheeze and eczema in Japanese infants aged 16-24 months: the Osaka Maternal and Child Health Study. <i>Pediatric Allergy and Immunology</i> , <b>2011</b> , 22, 734-41	4.2	46
77	Dietary intake of antioxidant vitamins and risk of Parkinson's disease: a case-control study in Japan. <i>European Journal of Neurology</i> , <b>2011</b> , 18, 106-13	6	58
76	Dietary glycemic index and glycemic load in relation to risk of overweight in Japanese children and adolescents: the Ryukyus Child Health Study. <i>International Journal of Obesity</i> , <b>2011</b> , 35, 925-36	5.5	22
75	Polyunsaturated fatty acid intake and prevalence of eczema and rhinoconjunctivitis in Japanese children: the Ryukyus Child Health Study. <i>BMC Public Health</i> , <b>2011</b> , 11, 358	4.1	15
74	Employment, income, and education and risk of postpartum depression: the Osaka Maternal and Child Health Study. <i>Journal of Affective Disorders</i> , <b>2011</b> , 130, 133-7	6.6	56
73	Organochlorine concentrations in breast milk and prevalence of allergic disorders in Japanese women. <i>Chemosphere</i> , <b>2011</b> , 85, 374-8	8.4	10
72	APOE and CYP2E1 polymorphisms, alcohol consumption, and Parkinson's disease in a Japanese population. <i>Journal of Neural Transmission</i> , <b>2011</b> , 118, 1335-44	4.3	8
71	IL13 genetic polymorphisms, smoking, and eczema in women: a case-control study in Japan. <i>BMC Medical Genetics</i> , <b>2011</b> , 12, 142	2.1	3
70	Case-control study of IL13 polymorphisms, smoking, and rhinoconjunctivitis in Japanese women: the Kyushu Okinawa Maternal and Child Health Study. <i>BMC Medical Genetics</i> , <b>2011</b> , 12, 143	2.1	5
69	Occupational risk factors for Parkinson's disease: a case-control study in Japan. <i>BMC Neurology</i> , <b>2011</b> , 11, 83	3.1	8
68	Genetic polymorphisms involved in dopaminergic neurotransmission and risk for Parkinson's disease in a Japanese population. <i>BMC Neurology</i> , <b>2011</b> , 11, 89	3.1	37
67	Causal assessment of smoking and tooth loss: a systematic review of observational studies. <i>BMC Public Health</i> , <b>2011</b> , 11, 221	4.1	52
66	Sibling number and prevalence of allergic disorders in pregnant Japanese women: baseline data from the Kyushu Okinawa Maternal and Child Health Study. <i>BMC Public Health</i> , <b>2011</b> , 11, 561	4.1	13
65	U-shaped association between body mass index and the prevalence of wheeze and asthma, but not eczema or rhinoconjunctivitis: the ryukyus child health study. <i>Journal of Asthma</i> , <b>2011</b> , 48, 804-10	1.9	24
64	Dietary patterns during pregnancy and the risk of postpartum depression in Japan: the Osaka Maternal and Child Health Study. <i>British Journal of Nutrition</i> , <b>2011</b> , 105, 1251-7	3.6	45
63	Case-control study of eczema associated with IL13 genetic polymorphisms in Japanese children. <i>International Archives of Allergy and Immunology</i> , <b>2011</b> , 154, 328-35	3.7	7

62	Nutritional adequacy of three dietary patterns defined by cluster analysis in 997 pregnant Japanese women: the Osaka Maternal and Child Health Study. <i>Public Health Nutrition</i> , <b>2011</b> , 14, 611-21	3.3	34
61	Association between prenatal and postnatal tobacco smoke exposure and allergies in young children. <i>Journal of Asthma</i> , <b>2011</b> , 48, 458-63	1.9	16
60	Active and passive smoking and risk of Parkinson's disease. <i>Acta Neurologica Scandinavica</i> , <b>2010</b> , 122, 377-82	3.8	31
59	Consumption of vegetables, fruit, and antioxidants during pregnancy and wheeze and eczema in infants. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , <b>2010</b> , 65, 758-65	9.3	103
58	Maternal meat and fat consumption during pregnancy and suspected atopic eczema in Japanese infants aged 3-4 months: the Osaka Maternal and Child Health Study. <i>Pediatric Allergy and Immunology</i> , <b>2010</b> , 21, 38-46	4.2	29
57	Residential proximity to main roads during pregnancy and the risk of allergic disorders in Japanese infants: the Osaka Maternal and Child Health Study. <i>Pediatric Allergy and Immunology</i> , <b>2010</b> , 21, 22-8	4.2	31
56	Fish and n-3 polyunsaturated fatty acid intake and depressive symptoms: Ryukyus Child Health Study. <i>Pediatrics</i> , <b>2010</b> , 126, e623-30	7.4	62
55	Dietary fat intake and risk of Parkinson's disease: a case-control study in Japan. <i>Journal of the Neurological Sciences</i> , <b>2010</b> , 288, 117-22	3.2	64
54	Case-control study of risk of Parkinson's disease in relation to hypertension, hypercholesterolemia, and diabetes in Japan. <i>Journal of the Neurological Sciences</i> , <b>2010</b> , 293, 82-6	3.2	96
53	LRRK2 Gly2385Arg polymorphism, cigarette smoking, and risk of sporadic Parkinson's disease: a case-control study in Japan. <i>Journal of the Neurological Sciences</i> , <b>2010</b> , 297, 15-8	3.2	19
52	Intake of dairy products and the prevalence of dental caries in young children. <i>Journal of Dentistry</i> , <b>2010</b> , 38, 579-83	4.8	25
51	GST polymorphisms, interaction with smoking and pesticide use, and risk for Parkinson's disease in a Japanese population. <i>Parkinsonism and Related Disorders</i> , <b>2010</b> , 16, 447-52	3.6	30
50	Dairy food, calcium and vitamin D intake in pregnancy, and wheeze and eczema in infants. <i>European Respiratory Journal</i> , <b>2010</b> , 35, 1228-34	13.6	196
49	Dietary intake of folate, vitamin B6, vitamin B12 and riboflavin and risk of Parkinson's disease: a case-control study in Japan. <i>British Journal of Nutrition</i> , <b>2010</b> , 104, 757-64	3.6	64
48	Dietary folate, riboflavin, vitamin B-6, and vitamin B-12 and depressive symptoms in early adolescence: the Ryukyus Child Health Study. <i>Psychosomatic Medicine</i> , <b>2010</b> , 72, 763-8	3.7	53
47	Dietary glycemic index is inversely associated with the risk of Parkinson's disease: a case-control study in Japan. <i>Nutrition</i> , <b>2010</b> , 26, 515-21	4.8	42
46	Alcohol drinking and risk of Parkinson's disease: a case-control study in Japan. <i>BMC Neurology</i> , <b>2010</b> , 10, 111	3.1	18
45	Household smoking and dental caries in schoolchildren: the Ryukyus Child Health Study. <i>BMC Public Health</i> , <b>2010</b> , 10, 335	4.1	17

44	Association between breastfeeding and allergic disorders in Japanese children. <i>International Journal of Tuberculosis and Lung Disease</i> , <b>2010</b> , 14, 513-8	2.1	9
43	Maternal fat consumption during pregnancy and risk of wheeze and eczema in Japanese infants aged 16-24 months: the Osaka Maternal and Child Health Study. <i>Thorax</i> , <b>2009</b> , 64, 815-21	7.3	64
42	The effect of maternal smoking during pregnancy and postnatal household smoking on dental caries in young children. <i>Journal of Pediatrics</i> , <b>2009</b> , 155, 410-5	3.6	36
41	Breastfeeding and atopic eczema in Japanese infants: The Osaka Maternal and Child Health Study. <i>Pediatric Allergy and Immunology</i> , <b>2009</b> , 20, 234-41	4.2	30
40	Sibship size and prevalence of allergic disorders in Japan: the Ryukyus Child Health Study. <i>Pediatric Allergy and Immunology</i> , <b>2009</b> , 20, 377-84	4.2	16
39	ADAM33 genetic polymorphisms and risk of atopic dermatitis among Japanese children. <i>Clinical Biochemistry</i> , <b>2009</b> , 42, 477-83	3.5	18
38	Education, but not occupation or household income, is positively related to favorable dietary intake patterns in pregnant Japanese women: the Osaka Maternal and Child Health Study. <i>Nutrition Research</i> , <b>2009</b> , 29, 164-72	4	43
37	Breastfeeding and the risk of wheeze and asthma in Japanese infants: the Osaka Maternal and Child Health Study. <i>Pediatric Allergy and Immunology</i> , <b>2008</b> , 19, 490-6	4.2	20
36	Tuberculin reactivity and allergic disorders in schoolchildren, Okinawa, Japan. <i>Clinical and Experimental Allergy</i> , <b>2008</b> , 38, 486-92	4.1	18
35	Fatty acid intake and asthma symptoms in Japanese children: the Ryukyus Child Health Study. <i>Clinical and Experimental Allergy</i> , <b>2008</b> , 38, 1644-50	4.1	42
34	Dental caries in 3-year-old children and smoking status of parents. <i>Paediatric and Perinatal Epidemiology</i> , <b>2008</b> , 22, 546-50	2.7	25
33	Relationship between soy and isoflavone intake and periodontal disease: the Freshmen in Dietetic Courses Study II. <i>BMC Public Health</i> , <b>2008</b> , 8, 39	4.1	18
32	Dietary glycemic index and load and the risk of postpartum depression in Japan: the Osaka Maternal and Child Health Study. <i>Journal of Affective Disorders</i> , <b>2008</b> , 110, 174-9	6.6	27
31	Ambient formaldehyde levels and allergic disorders among Japanese pregnant women: baseline data from the Osaka maternal and child health study. <i>Annals of Epidemiology</i> , <b>2008</b> , 18, 78-84	6.4	17
30	Dental caries and allergic disorders in Japanese children: the Ryukyus Child Health Study. <i>Journal of Asthma</i> , <b>2008</b> , 45, 795-9	1.9	18
29	Maternal smoking and environmental tobacco smoke exposure and the risk of allergic diseases in Japanese infants: the Osaka Maternal and Child Health Study. <i>Journal of Asthma</i> , <b>2008</b> , 45, 833-8	1.9	32
28	Genetic susceptibility to atopic dermatitis. <i>Allergology International</i> , <b>2008</b> , 57, 39-56	4.4	49
27	Relationship between dietary fat and fish intake and the prevalence of atopic eczema in pregnant Japanese females: baseline data from the Osaka Maternal and Child Health Study. <i>Asia Pacific Journal of Clinical Nutrition</i> , <b>2008</b> , 17, 612-9	1	12

26	Fish and fat intake and prevalence of allergic rhinitis in Japanese females: the Osaka Maternal and Child Health Study. <i>Journal of the American College of Nutrition</i> , <b>2007</b> , 26, 279-87	3.5	36
25	Environmental factors and allergic disorders. <i>Allergology International</i> , <b>2007</b> , 56, 363-96	4.4	20
24	Home environment and suspected atopic eczema in Japanese infants: the Osaka Maternal and Child Health Study. <i>Pediatric Allergy and Immunology</i> , <b>2007</b> , 18, 425-32	4.2	44
23	Cross-sectional study of allergic disorders associated with breastfeeding in Japan: the Ryukyus Child Health Study. <i>Pediatric Allergy and Immunology</i> , <b>2007</b> , 18, 433-40	4.2	20
22	Association of total tooth loss with smoking, drinking alcohol and nutrition in elderly Japanese: analysis of national database. <i>Gerodontology</i> , <b>2007</b> , 24, 87-92	2.8	41
21	Cigarette smoking and tooth loss experience among young adults: a national record linkage study. <i>BMC Public Health</i> , <b>2007</b> , 7, 313	4.1	32
20	Relationship between smoking status and tooth loss: findings from national databases in Japan. <i>Journal of Epidemiology</i> , <b>2007</b> , 17, 125-32	3.4	32
19	Relationship between intake of vegetables, fruit, and grains and the prevalence of tooth loss in Japanese women. <i>Journal of Nutritional Science and Vitaminology</i> , <b>2007</b> , 53, 522-8	1.1	9
18	Prevalence of asthma and wheeze in relation to passive smoking in Japanese children. <i>Annals of Epidemiology</i> , <b>2007</b> , 17, 1004-10	6.4	37
17	Fat and fish intake and asthma in Japanese women: baseline data from the Osaka Maternal and Child Health Study. <i>International Journal of Tuberculosis and Lung Disease</i> , <b>2007</b> , 11, 103-9	2.1	21
16	Dietary folate and vitamins B12, B6, and B2 intake and the risk of postpartum depression in Japan: the Osaka Maternal and Child Health Study. <i>Journal of Affective Disorders</i> , <b>2006</b> , 96, 133-8	6.6	70
15	Risk of postpartum depression in relation to dietary fish and fat intake in Japan: the Osaka Maternal and Child Health Study. <i>Psychological Medicine</i> , <b>2006</b> , 36, 1727-35	6.9	93
14	Association of smoking in household and dental caries in Japan. <i>Journal of Public Health Dentistry</i> , <b>2006</b> , 66, 279-81	1.6	18
13	Relationship between smoking status and periodontal conditions: findings from national databases in Japan. <i>Journal of Periodontal Research</i> , <b>2006</b> , 41, 573-9	4.3	63
12	Active and passive smoking and tooth loss in Japanese women: baseline data from the osaka maternal and child health study. <i>Annals of Epidemiology</i> , <b>2005</b> , 15, 358-64	6.4	26
11	Association between RANTES promoter polymorphism -401A and enhanced RANTES production in atopic dermatitis patients. <i>Journal of Dermatological Science</i> , <b>2005</b> , 39, 189-91	4.3	29
10	Functional SNPs in the distal promoter of the ST2 gene are associated with atopic dermatitis. <i>Human Molecular Genetics</i> , <b>2005</b> , 14, 2919-27	5.6	152
9	Inhibition of dentine demineralization by zinc oxide: in vitro and in situ studies. <i>Dental Materials</i> , <b>2005</b> , 21, 1170-7	5.7	41

8	Association of melanin pigmentation in the gingiva of children with parents who smoke. <i>Pediatrics</i> , <b>2005</b> , 116, e186-90	7.4	50
7	Heterogeneity of interleukin 5 genetic background in atopic dermatitis patients: significant difference between those with blood eosinophilia and normal eosinophil levels. <i>Journal of Dermatological Science</i> , <b>2003</b> , 33, 121-6	4.3	29
6	Mechanism of cytotoxicity of paraquat. <i>Environmental Health and Preventive Medicine</i> , <b>2002</b> , 7, 89-94	4.2	65
5	Enamel fluoride uptake from mouthrinse solutions with different NaF concentrations. <i>Community Dentistry and Oral Epidemiology</i> , <b>2002</b> , 30, 248-53	2.8	3
4	Lack of association between atopic eczema and the genetic variants of interleukin-4 and the interleukin-4 receptor alpha chain gene: heterogeneity of genetic backgrounds on immunoglobulin E production in atopic eczema patients. <i>Clinical and Experimental Allergy</i> , <b>2001</b> , 31, 1522-7	4.1	37
3	Acid resistance of human enamel in vitro after bicarbonate application during remineralization. <i>Journal of Dentistry</i> , <b>2001</b> , 29, 421-6	4.8	9
2	Association between mast cell chymase genotype and atopic eczema: comparison between patients with atopic eczema alone and those with atopic eczema and atopic respiratory disease. <i>Clinical and Experimental Allergy</i> , <b>1999</b> , 29, 800-3	4.1	48
1	Prevalence of childhood and adolescent atopic dermatitis in a Japanese population: comparison with the disease frequency examined 20 years ago. <i>Acta Dermato-Venereologica</i> , <b>1998</b> , 78, 293-4	2.2	110