Enrique Juan Sanchez Pozzi

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/4192401/enrique-juan-sanchez-pozzi-publications-by-year.pdf

Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

47 papers 1,168 19 h-index g-index

47 type 1,261 type 2,56 type 2

#	Paper	IF	Citations
47	Role of ERK1/2 in TNFIInduced internalization of Abcc2 in rat hepatocyte couplets. <i>Biochemical Pharmacology</i> , 2019 , 164, 311-320	6	3
46	Sphingosine 1-phosphate receptor 2/adenylyl cyclase/protein kinase A pathway is involved in taurolithocholate-induced internalization of Abcc2 in rats. <i>Archives of Toxicology</i> , 2019 , 93, 2279-2294	5.8	2
45	Dynamic Localization of Hepatocellular Transporters: Role in Biliary Excretion and Impairment in Cholestasis. <i>Current Medicinal Chemistry</i> , 2019 , 26, 1113-1154	4.3	5
44	Mechanisms of canalicular transporter endocytosis in the cholestatic rat liver. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2018 , 1864, 1072-1085	6.9	19
43	Activation of insulin-like growth factor 1 receptor participates downstream of GPR30 in estradiol-17[-D-glucuronide-induced cholestasis in rats. <i>Archives of Toxicology</i> , 2018 , 92, 729-744	5.8	2
42	Mitogen-activated protein kinases are involved in hepatocanalicular dysfunction and cholestasis induced by oxidative stress. <i>Archives of Toxicology</i> , 2017 , 91, 2391-2403	5.8	9
41	EGFR participates downstream of ERIIn estradiol-17I-D-glucuronide-induced impairment of Abcc2 function in isolated rat hepatocyte couplets. <i>Archives of Toxicology</i> , 2016 , 90, 891-903	5.8	8
40	Sandwich-cultured rat hepatocytes as an in vitro model to study canalicular transport alterations in cholestasis. <i>Archives of Toxicology</i> , 2015 , 89, 979-90	5.8	10
39	Physiological concentrations of unconjugated bilirubin prevent oxidative stress-induced hepatocanalicular dysfunction and cholestasis. <i>Archives of Toxicology</i> , 2014 , 88, 501-14	5.8	12
38	G-protein-coupled receptor 30/adenylyl cyclase/protein kinase A pathway is involved in estradiol 17ED-glucuronide-induced cholestasis. <i>Hepatology</i> , 2014 , 59, 1016-29	11.2	39
37	Radical Oxygen Species and Bile Secretion 2014 , 1787-1808		1
36	The Call+-calmodulin-Call+/calmodulin-dependent protein kinase II signaling pathway is involved in oxidative stress-induced mitochondrial permeability transition and apoptosis in isolated rat hepatocytes. <i>Archives of Toxicology</i> , 2014 , 88, 1695-709	5.8	39
35	Hormonal modulation of hepatic cAMP prevents estradiol 17I-D-glucuronide-induced cholestasis in perfused rat liver. <i>Digestive Diseases and Sciences</i> , 2013 , 58, 1602-14	4	10
34	ERK1/2 and p38 MAPKs are complementarily involved in estradiol 17ED-glucuronide-induced cholestasis: crosstalk with cPKC and PI3K. <i>PLoS ONE</i> , 2012 , 7, e49255	3.7	22
33	Localization status of hepatocellular transporters in cholestasis. <i>Frontiers in Bioscience - Landmark</i> , 2012 , 17, 1201-18	2.8	37
32	Sequential activation of classic PKC and estrogen receptor ls involved in estradiol 17ED-glucuronide-induced cholestasis. <i>PLoS ONE</i> , 2012 , 7, e50711	3.7	17
31	Ursodeoxycholic acid in cholestasis: linking action mechanisms to therapeutic applications. <i>Clinical Science</i> , 2011 , 121, 523-44	6.5	103

(2003-2011)

30	Prevention of estradiol 17beta-D-glucuronide-induced canalicular transporter internalization by hormonal modulation of cAMP in rat hepatocytes. <i>Molecular Biology of the Cell</i> , 2011 , 22, 3902-15	3.5	19
29	Phosphoinositide 3-kinase/protein kinase B signaling pathway is involved in estradiol 17I-D-glucuronide-induced cholestasis: complementarity with classical protein kinase C. <i>Hepatology</i> , 2010 , 52, 1465-76	11.2	43
28	Putative role for actin organization status in the dynamic localization of canalicular carriers under oxidative stress conditions. <i>American Journal of Physiology - Renal Physiology</i> , 2009 , 296, G969	5.1	1
27	Differential effects of silymarin and its active component silibinin on plasma membrane stability and hepatocellular lysis. <i>Chemico-Biological Interactions</i> , 2009 , 179, 297-303	5	46
26	Induction of intestinal multidrug resistance-associated protein 2 (Mrp2) by spironolactone in rats. <i>European Journal of Pharmacology</i> , 2009 , 623, 103-6	5.3	13
25	Dapsone induces oxidative stress and impairs antioxidant defenses in rat liver. <i>Life Sciences</i> , 2008 , 83, 155-63	6.8	18
24	Ca(2+)-dependent protein kinase C isoforms are critical to estradiol 17beta-D-glucuronide-induced cholestasis in the rat. <i>Hepatology</i> , 2008 , 48, 1885-95	11.2	58
23	Disruption of function and localization of tight junctional structures and Mrp2 in sustained estradiol-17beta-D-glucuronide-induced cholestasis. <i>American Journal of Physiology - Renal Physiology</i> , 2007 , 293, G391-402	5.1	27
22	Oxidative stress induces actin-cytoskeletal and tight-junctional alterations in hepatocytes by a Ca2+-dependent, PKC-mediated mechanism: protective effect of PKA. <i>Free Radical Biology and Medicine</i> , 2006 , 40, 2005-17	7.8	46
21	Oxidative stress induces internalization of the bile salt export pump, Bsep, and bile salt secretory failure in isolated rat hepatocyte couplets: a role for protein kinase C and prevention by protein kinase A. <i>Toxicological Sciences</i> , 2006 , 91, 150-8	4.4	66
20	Galactosamine prevents ethinylestradiol-induced cholestasis. <i>Drug Metabolism and Disposition</i> , 2006 , 34, 993-7	4	17
19	Hepatic and extrahepatic synthesis and disposition of dinitrophenyl-S-glutathione in bile duct-ligated rats. <i>Drug Metabolism and Disposition</i> , 2006 , 34, 1301-9	4	15
18	Inhibition of rat liver UDP-glucuronosyltransferase by silymarin and the metabolite silibinin-glucuronide. <i>Life Sciences</i> , 2005 , 77, 683-92	6.8	30
17	Silibinin prevents cholestasis-associated retrieval of the bile salt export pump, Bsep, in isolated rat hepatocyte couplets: possible involvement of cAMP. <i>Biochemical Pharmacology</i> , 2005 , 69, 1113-20	6	27
16	Involvement of Mrp2 in hepatic and intestinal disposition of dinitrophenyl-S-glutathione in partially hepatectomized rats. <i>Toxicological Sciences</i> , 2005 , 84, 4-11	4.4	12
15	Role of microtubules in estradiol-17beta-D-glucuronide-induced alteration of canalicular Mrp2 localization and activity. <i>American Journal of Physiology - Renal Physiology</i> , 2005 , 288, G327-36	5.1	43
14	Prevention of Mrp2 activity impairment in ethinylestradiol-induced cholestasis by ursodeoxycholate in the rat. <i>Drug Metabolism and Disposition</i> , 2005 , 33, 888-91	4	10
13	Ursodeoxycholate reduces ethinylestradiol glucuronidation in the rat: role in prevention of estrogen-induced cholestasis. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2003 , 306, 279-8	36 ^{4.7}	27

12	Estradiol-17beta-D-glucuronide induces endocytic internalization of Bsep in rats. <i>American Journal of Physiology - Renal Physiology</i> , 2003 , 285, G449-59	5.1	105
11	Effect of spironolactone on the expression of rat hepatic UDP-glucuronosyltransferase. <i>Biochemical Pharmacology</i> , 2003 , 66, 171-7	6	9
10	Preventive effect of silymarin against taurolithocholate-induced cholestasis in the rat. <i>Biochemical Pharmacology</i> , 2003 , 66, 355-64	6	19
9	Dapsone-induced cholestasis and impairment of bile salt output in the rat. <i>Biochemical Pharmacology</i> , 2002 , 63, 1553-63	6	9
8	Beneficial effects of silymarin on estrogen-induced cholestasis in the rat: a study in vivo and in isolated hepatocyte couplets. <i>Hepatology</i> , 2001 , 34, 329-39	11.2	68
7	Effect of silymarin on biliary bile salt secretion in the rat. <i>Biochemical Pharmacology</i> , 2000 , 59, 1015-22	6	20
6	Induction of phase II biotransformation reactions in rat jejunum during lactation. Possible involvement of prolactin. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 1999 , 1472, 82-92	4	7
5	Enhancement of intestinal UDP-glucuronosyltranferase activity in partially hepatectomized rats. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 1998 , 1380, 345-53	4	15
4	Gender-related differences in the amount and functional state of rat liver UDP-glucuronosyltransferase. <i>Biochemical Pharmacology</i> , 1995 , 50, 509-14	6	29
3	Analysis of p-nitrophenol glucuronidation in hepatic microsomes from lactating rats. <i>Biochemical Pharmacology</i> , 1994 , 47, 1179-85	6	19
2	Inhibition of rat liver microsomal bilirubin UDP-glucuronosyltransferase by ursodeoxycholic acid. <i>Life Sciences</i> , 1994 , 55, 111-20	6.8	9
1	Analysis of the interaction uridin 5\diphosphoglucuronic acid with intestinal bilirubin UDP-glucuronyltransferase. <i>International Journal of Biochemistry & Cell Biology</i> , 1992 , 24, 1429-34		3